

lata, interdum tantum in margine barbata plerumque extus fuscostrigosa.

Hab. in frustulis ligni dejecti, Bethlehem Amer. bor.

32. **Lachnella maculicola** (Schw.) Sacc. *Peziza maculicola* Schw. Syn. 1646
ops. pag. 174. — Gregaria, 2 mm. diam. excedens, sessilis, basi nigra nuda rugosa, versus marginem strigosissima, tomento stri-goso ferrugineo-rubo, subpulverulento; disco primum clauso (tum ascomate globoso), demum aperto, corneo-rubo; cupulis in macula subconcentricis.

Hab. in maculis determinatis albidis omnino tamen non crustaceis, in lignis decorticatis ramorum, aggregata, Bethlehem et Buffalo Americæ borealis. — A *L. flammea*, teste Berk., non satis differt.

33. **Lachnella farinosa** (Wallr.) Sacc. *Peziza farinosa* Wallr. Fl. 1647
crypt. n. 2441. — Ascomatibus sessilibus, molliter carnosulis, sic-cis clavis globosis, extus pilis propexis tenerrimis dilute cinnamomeis sparsis confertisve iisque apice farinula albida adspersis undique vestitis similiterque in stratum tenue intertextis, humectis discum concavum dilute-carneum margine inflexo cinctum reclu-dentibus.

Hab. ad lignorum cæsorum frustula humi jacentia, Fraxini, Tiliæ, Fagi in Thuringia passim. — Cum *P. rufiberbi* Alb. et Schw. et Fr. syst. II, p. 107, forsan comparanda.

34. **Lachnella ascoboloidea** (Schw.) Sacc. *Peziza ascoboloidea* Schw. 1648
Syn. p. 175. — Minutissima, sparsa, extus fibrosa, candida, turbina-ta, sed vix stipitata, disco subcarneo quasi ascis ascoboloideis punctato, subconvexulo, siccitate clauso.

Hab. in ramis corticis Vitis Labruscæ etc., Bethlehem Amer. bor. — 1 mm. circ. lat.

35. **Lachnella tumida** (Pers.) Sacc. *Peziza tumida* Pers. Eur. I, p. 269, 1649
Fr. Syst. Myc. II, p. 97. — Sessilis, subcoriacea, depresso ventri-cosa, leviter tomentosa, opaca, gilvo-pallida, disco tumido nigre-scente.

Hab. in sarm entis Vitis viniferæ rarissime (CHAILLET) et in Thuringia (WALLR.). — Ab initio tota clausa videtur, saltim to-mento involuta, in medio dein aperitur ore circulari; discus tan-dem conspicitur, qui est superficialis seu prominens, rugulosus qua-si et immarginatus; latitudo linearis.

36. **Lachnella fusco-barbata** (Schw.) Sacc. *Peziza fusco-barbata* Schw. 1650
Syn. p. 173. — Patellæformis, extus duriuscula et in basi nigri-

cans, 2-4 mm. diam., margine densim fusco-barbata, disco eleganter carneo, sparsa, siccitate clausa, extra barbam nuda rugosa.

Hab. in caulis valde corruptis Verbasci, Bethlehem Amer. borealis.

37. **Lachnella solitaria** (Schw.) Sacc. *Peziza solitaria* Schw. Syn. 1651 p. 175. — Sessilis, solitaria, majuscula ad 6 mm. lata, applanata, firma, crassa, margine tumido obtuso glabro, colore chalybeo; disco undulatim inæquali; cupula extus vestita fibris quibusdam paucis albescentibus.

Hab. in stipitibus tenerrimis, Bethlehem Amer. bor.

38. **Lachnella extricata** (B. et C.) Sacc. *Peziza extricata* B. et C. 1652 North Am. Fungi n. 719. — Erumpens, congesta margine undulato; extus pallide umbrina sericeo-tomentosa, intus albida.

Hab. in caulis emortuis Umbelliferarum, Alabama Amer. bor. (PETERS). — Pulchra species, sed immatura observata.

TRICHOPEZIZA Fuck. Symb. p. 295 (1869) [Etym. *thrix* capillus et *Peziza*], *Peziza* sect. *Dasyscypha* Fr. Syst. Myc. II, p. 89, *Lachnella* Sacc. e. p. *Hyalopeziza* Fuck. — Ascoma ceraceum, sessile vel subsessile, ex urceolato scutellatum, extus pilosum v. margine distinctius ciliatum, nunc minutum tenacellum, nunc minutissimum, tenerius. Asci tereti-fusoidei, 4-8-spori. Sporidia tereti-oblonga vel fusoidea, continua hyalina. Paraphyses saepius bacillari-fusoideæ, crassiusculæ. — Species quædam aptius ad *Pseudohelotium* et *Dasyscypham* referri possent.

A. Species fructificatione cognita.

I. Caulicolæ.

1. **Trichopeziza sulphurea** (Pers.) Fuck. Symb. p. 296, *Lachnum sulphureum* Karst. M. F. I, p. 176, *Peziza sulphurea* Pers. Disp. p. 33, Fr. S. M. II, p. 104, Pat. Tab. 280, Greg. t. 83, *P. citrinella* DC. Franc. VI, p. 24, *Lachnea* Gill. Disc. c. ic. — Sparsa vel gregaria, sessilis, hemisphærica, pallida, sulphureo-strigoso-villosa, vetustate et siccitate saepè fuscescens, ferrugineo-fusco-villosa, disco pallido vel albido, 1 mm. lat.; ascis cylindraceo-clavatis, 65-73 \approx 5-6; sporidiis fusoideo-filiformibus guttulatis vel spurie tenuiter uni-triseptatis aut eguttulatis, 20-34 \approx 2-2,5; paraphysibus apice acutiusculis 4,5 μ . cr.

Hab. ad caules putrescentes Urticæ dioicæ, Cerefolii silvestris, Solani tuberosi, Sambuci Eboli, Antirrhini.

ni in Fennia, Britannia, Belgio, Gallia, Sibiria asiatica, Amer. bor.

— Teste Fuck., stat. conidicus est *Tricholeconium roseum* Corda.

2. **Trichopeziza leucophæa** (Pers.) Rehm Ascom. n. 65, *Peziza sulphurea* var. *leucophæa* Pers. Myc. eur. I, p. 250, *P. leucophæa* Nyl. Obs. p. 31, *P. sulphurea* Auct. pr. p., *Lachnum leucophæum* Karst. M. F. I, p. 177. — Sparsa vel gregaria, sessilis, hemisphærica, pallida, albicanti- v. sulphureo-strigoso-villosa, siccitate et vetustate ferrugineo-fusco-villosa, disco albido, 1 mm. lat., pilis flavis septatis, scabris 4-5 μ . cr.; ascis cylindraceo-subclavatis, 50-60 \approx 4-5; sporidiis fusoideo-filiformibus simplicibus, 7-16 \approx 1,5-2 distichis; paraphysibus fusoideis supra acutiusculis, 3,5-4 μ . cr.

Hab. ad caules putrescentes Umbelliferarum majorum, Urticæ dioicæ, Aristolochiæ, Arunci, Artemisiæ vulgaris etc. in Fennia, Lapponia, Germania, Gallia, Italia. — Forma **alpestris** Rehm Asc. n. 308. Extus sulphurea vel viridi-sulphurea, disco flavidulo, hirta vel strigosa, 2-3 mm. diam. explanata; ascis 60 \approx 5; sporidiis 12-15 \approx 1,5. Ad caules putridos Cirsii spinosissimi, in alpibus tyrolensibus juxta glaciem æternam.

3. **Trichopeziza relicina** (Fr.) Fuck. Symb. p. 296, *Peziza relicina* 1655 Fr. Syst. Myc. II, p. 103, Pat. Tab. p. 282, *P. Atropæ* A. S. Lus. p. 320, *Excipula phæotricha* Rabenh., *Lachnum relicinum* (Fr.) Karst. M. F. I, p. 182. — Radio-fusca, sparsa vel subgregaria, sessilis v. brevissime stipitata, obconica (sicca), pilis erectis, confertis, badiis strigosa, 4-5 mm. lat.; ascis clavatis, apice incrassatis, 8-sporis, 60-70 \approx 6-8; sporidiis oblongis, obtusis, rectis vel subcurvatis, hyalinis, 1-cellularibus, 12-15 \approx 2,5-3, distichis; paraphysibus lanceolatis, apice acuminatis, c. 2,5 μ . cr., ascos superantibus.

Hab. in caulibus Artemisiæ, Urticæ, Aconiti, Chenopodiæ, Sonchi, Cacaliæ in Fennia, Suecia, Gallia, Helvetia, Germania, Italia, Hungaria, Amer. bor. — Pili distincte articulati, fusi, apice vix incrassato, 180-220 \approx 4,5-6 mm. long. A *T. Nidulo* affini cupulis minoribus, pilis erectis longioribus et tenuioribus diversa.

4. **Trichopeziza nidulans** (Quél.) Sacc. *Lachnella nidulans* Quél. 11 1656 Suppl. p. 20, t. XII, f. 5. — Ascomate sessili, ovoideo, applanato, 1-2 mm. diam., humido, tenui, ceraceo, ciliato-hirto, brunneo vel fulvo, basi subiculo e filamentis arachnoideis, albis prædito; hymenio concavo incarnato-pallescente; sporidiis fusoideis, 12-15 μ . longis, septatis.

Hab. gregatim in caulis majoribus Urticæ, Eupatorii in Jura. — Affinis *T. Nidulo* et *T. relicinæ*.

5. **Trichopeziza mollissima** (Lasch) Fuck. Symb. myc. p. 296, *Peziza 1657* *mollissima* Lasch in Reb. H. M. II, n. 708, *Lachnum mollissimum* Karst. M. F. I, p. 174, *Peziza ovina* Lib. Herb. — Sparsa v. gregaria, sessilis, albida vel pallida ex hemisphærico appanata, sicca hemisphærica vel subsphæroideo-contracta, colore immutata, albo-villosa, margine sericeo-fimbriato, 0,5 mm. lat.; ascis subclavato-cylindraceis 50-66 \approx 4-4,5; sporidiis filiformi-fusoideis, vulgo egutulatis, 8-14 \approx 1,5; paraphysibus 4-5 μ . cr., apice cuspidatis.

Hab. ad caules Urticæ dioicæ, Umbelliferarum variarum, Eupatorii, Actææ, Valerianæ et Artemisiæ vulgaris putrescentes, in Fennia meridionali et media et in Gallia, Germania, Hungaria.

6. **Trichopeziza subglobosa** (Saut.) Sacc. *Peziza subglobosa* Saut. 1658 Pilz. Salzb. p. 10, Wint. Hedw. 1881, p. 132. — Globosa, tenera, albido-pallens, margine ciliato; ascis cylindraceo-clavatis, 8-sporis, 44-50 \approx 5; sporidiis cylindraceis, utrinque rotundatis, continuis, hyalinis, 9-12 \approx 2; paraphysibus filiformibus.

Hab. in caulis Urticæ emortuis, Mittersill Salzburg.

7. **Trichopeziza urticina** (Peck) Sacc. *Peziza urticina* Peck Bull. N. Y. Mus. 1887, p. 21. — Ascomate minuto, 170-350 μ . diam., sessili, subgloboso, primo albo, udo labiis conniventibus, sicco piloso-furfuraceo, albescente; ascis subfusiformibus; sporidiis stipatis v. distichis, fusiformibus, 10-13 μ . longis; paraphysibus filiformibus.

Hab. in caulis emortuis Urticæ (La portææ) canadensis, Catskill Amer. bor. — Pili madore sunt approximati et ascoma longitudinaliter striatum videtur. Discus in sicco sæpe absconditus.

8. **Trichopeziza rubicunda** (Saut.) Sacc. *Peziza rubicunda* Saut. Flor. Salzb. p. 8, Wint. Hedw. 1881, p. 130. — Sessilis, irregularis, appanata, submembranacea, flavens, disco miniato; ascis elongato-clavatis, 8-sporis, 35-40 \approx 5; sporidiis oblique distichis, cylindraceis, utrinque rotundatis, sæpe parum curvatis, continuis, hyalinis, 6-7 \approx 2.

Hab. in caulis putridis Urticæ prope Mittersill Salzburg. — Ex Winter *Moll. chlorinellæ* Ces. affinis, sed vix, cum inter vestitas ab auctore locetur.

9. **Trichopeziza crispula** (Karst.) Sacc. *Hel. crispulum* Karst. M. F. 1661 I, p. 161. — Gregaria vel subsparsa, sessilis, urceolata, tomento-

sa, albida, 0,2-0,3 mm. lat.; ascis cylindraceis vel fusoideo-cylindraceis, 25-36 \times 4-4,5, obturaculo minutissimo jodo leviter cœrulecente; sporidiis fusoideo-elongatis, rectis, 2-4-guttulatis, 6-10 \times 1,5-2; paraphysibus gracilibus, 0,5 μ . cr.

Hab. ad caules emortuos Chenopodii Boni Henrici, Artemisiæ vulgaris, Rumicis et Cerefolii silvestris in Fennia australi et Nordcap Norvegiæ. — Pili flexuoso-curvati, inarticulati, apicem versus attenuati, 100 \times 3-6. Forma, colore et magnitudine ascomatum cum *Pseudohel. scrupuloso* convenit. Vergit ad *Mollisiam*.

10. **Trichopeziza brevipila** (Rob. et Desm.) Sacc. *Peziza brevipila* 1662
Rob. in Desm. 17 Not. pag. 36. — Caulicola, erumpens, sparsa, sessilis, minuta, orbicularis vel suboblonga, pallide grisea, externe hirsuta; pilis brevibus, nigris; disco plano vel concaviusculo, pallido; ascis tubulosis, 55 μ . longis; sporidiis elongatis, linearibus, rectis vel curvulis; guttulis 4-6 globosis.

Hab. in caulis siccis Centaureæ, in Gallia (DESMAZIÈRES).

— Discus apertus 700-750 μ . æquans.

11. **Trichopeziza Bernardiana** Sacc. et Let. Misc. mycol. I, p. 8. — 1663
Ascomatibus gregariis, sessilibus, cupularibus, totis pallide carneo-luteolis, $\frac{1}{3}$ - $\frac{1}{2}$ mm. diam., subceraceis, extus pilis exiguis asperulis lutescentibus ornatis; ascis teretiusculis, brevissime stipitatis, filiformi-paraphysatis, 50-60 \times 6, octosporis; sporidiis distichis fusoideo-cylindraceis, lenissime curvulis, utrinque obtusiusculis, 10-13 \times 1,7-2, hyalinis.

Hab. in caule emortuo Cirsii palustris prope Rouen (LE-TENDRE). — Generi *Helotio* affinis.

12. **Trichopeziza subnidulans** Rehm Ascom. Alp. n. 15. — Plerumque gregaria, sessilis, primitus globosa, extus flavidо-brunnea, stri-gosa, humida explanata, margine fimbriata, subtus nuda fusca, disco livido-pallido, c. 0,7-8 μ . diam., orbicularis, parenchymatico-contexta, pilis marginalibus mediocribus, 6-7 μ . cr., fuscis, apice subclavatis, obtusis, hyalinis, anguste septatis, simplicibus; ascis clavatis 8-sporis, 45 \times 5, apice jodo cœrulecente; sporidiis oblongatis, obtusis, 1-cellularibus, hyalinis, distichis, 6-7 \times 4,5; paraphysibus lanceolatis, apice acutatis, basi 5 μ . cr.

Hab. ad caules putridos Adenostyliidis alpinæ in monte Bauernalpkopf alpium Allgoviæ c. 1500 m. alt. (BRITZELMAYR). — *Trichopezizæ Nidulo* affinis.

13. **Trichopeziza cirrata** (Cr.) Sacc. *Peziza cirrata* Cr. Fin. p. 51. 1665

— Ascomate $\frac{1}{5}$ mm. diam., albo, sessili, subsphærico, concavo, villosa; pilis brevibus, fragilibus, continuis, sursum attenuatis tecto; ascis oblongis, 8-sporis; sporidiis subcylindraceis, hyalinis; paraphysibus filiformibus punctulatis.

Hab. in caule Eupatorii cannabini, Brest Galliæ (DE GUERNISAC).

14. **Trichopeziza carneo-rubra** (Ell.) Sacc. *Peziza carneo-rubra* 1666
Ellis in Cooke Disc. U. S. II, p. 22. — Sessilis, rosea, 1 mm. lat.; sporidiis linearibus bi-nucleatis.

Hab. in caule Erigeronis, New Jersey Amer. bor. (ELLIS).

15. **Trichopeziza spirotricha** (Oud.) Sacc. *Pez. spirotricha* Oud. mat. 1667
pour. fl. myc. Neerl. II, Hedw. 1874, p. 91, *Das. Rehm Asc. n. 510*,
Peziza asterostoma Phill. in Grev. VII, p. 140. — Gregaria, primitus globosa, dein apice applanato-globosa, sessilis, primitus punctiformiter aperta, dein stictioidea, concava, urceolata, margine albo, siccata albella, humida carnea, dein albido-flavidula, 0,3-4 mm. diam., contextu parenchymatico, hyalino; cellulis magnis tenerrimis; pilis creberrimis, gregariis, hyalinis, glabris, acutis, simplicibus, longis, basi ampliatis et spiraliter contortis 4 μ . crassis obsessa; ascis clavatis, apice incrassatis, 8-sporis, 36 \approx 6 apice jodo cœrulecente; sporidiis ellipticis vel elongato-ovoideis, rectis, hyalinis, 1-cellularibus, interdum 2 nucleolis præditis, distichis, 9 \approx 2,5; paraphysibus teneris, filiformibus, superne subincrassatis.

Hab. ad caules putridos Alliariæ, Adenostyolidis albifrontis in Hollandia, Britannia, Tyrolia. — Sporidia sec. Phill. 7 \approx 1,5; pili 70 μ . long., plerumque marginales; ascoma usque 2 mm.

16. **Trichopeziza elegantula** (Karst.) Sacc. *Lachn. elegantula* Karst. 1668
Rev. monogr. pag. 122, *Helotium elegantulum* Karst. M. F. I, p. 132. — Sparsa vel aggregata, sessilis vel brevissime stipitata, obovoidea, dein hemisphærico-applanata, margine connivente infexo, nigra, villo pururascenti-fusco, facile demum secedente, obvelata, 0,6 mm. lat.; ascis cylindraceis, 50-70 \approx 5; sporidiis fusoideo-oblongatis vel fusoideo-elongatis, rectis, 6-13 \approx 2, eguttulatis; paraphysibus filiformibus 1-1,5 μ . crass.

Hab. ad caules Epilobii angustifolii, Solidaginis, Virgaureæ et Umbelliferarum putrescentes in par. Lempäälä prov. Tavastland Fenniæ et in Lapponia orientali. — Pili cupulæ flexuosi, fusci, haud vel obsolete articulati, 3 μ . cr.

17. **Trichopeziza Hyperici** (Schröt.) Sacc. *Velutaria Hyperici* Schröt. 1669
Bericht. Schles. Ges. 1875, p. 37. — Subsessilis, sparsa vel 2-3-

aggregata, erumpens, extus brunneo-flava, farinosa; disco olivaceo-brunneo; ascis teretibus, 8-sporis; sporidiis ellipsoideis v. ovoideis, continuis, 11-12 \approx 7.

Hab. in caulis emortuis Hyperici perforati, Baden.

18. **Trichopeziza chlorospleniella** Rehm Ascom. Alp. n. 11. — Sparsa, 1670 sessilis, sphæroidea, dein urceolata, nigrella, extus nigro-pilosa, humida urceolato-concava, margine involuto sublacero, subrugosa, fusco-atra, disco hyalinello, c. 1 mm. diam., prosenchymatico-fusco-contexta, pilis brevibus, septatis, fuscis, obtusis, glabris, sparsis, c. 4 μ . cr., obsessa, in caule longe lateque æruginoso-tincto sessilis; ascis clavatis, 8-sporis, 36 \approx 5, apice jodo cœrulecente; paraphysibus ascos superantibus, lanceolato-acutatis, basi c. 6 μ . cr.

Hab. ad caulem putridum Aconiti variegati, Kartel-Gletscher in valle Moos alpium tyrolensis, c. 2000 m. alt. (BRITZELMAYR).

19. **Trichopeziza leucostoma** (Rehm) Sacc. *Dasyscypha leucostoma* 1671 Rehm Ascom. n. 205 et F. Lojk. n. 27. — Gregaria, sessilis, globosa, urceolata, extus fusco-brunnea v. fusco-nigra, hirta, ore conivente albo, ex fibris hyalinis constituto, humido aperto, disco dein conspicuo, concavo pallido, 0,4 mm. diam., prosenchymatico-fusco-contexta, pilis longis, fuscidulis, obtusis, subscabris, simplicibus, c. 3 μ . cr. obsessa; ascis cylindraceo-clavatis, apice incrassatis, 80-90 \approx 5, 8-sporis, apice jodo cœrulecente; sporidiis oblongis v. subfusiformibus, plerumque rectis, 1-cellularibus, hyalinis, distichis, 10-16 \approx 1,5-2; paraphysibus apice haud clavatis.

Hab. ad caules putridos Aconiti Napelli aliarumque herbarum circa lacum Zenoga in alpibus edit. transsylvaniae (LOJKA) in Helvetia et Tyrolia (REHM).

20. **Trichopeziza lanata** (Rehm) Sacc. *Dasyscypha lanata* Rehm 1672 Ascom. Lojk. n. 25. — Aperta, explanata, sessilis, solitaria, sicca corrugata, humida, disco flavo plano, extus hirta, alba, parenchymatico-flavido-contexta, pilis septatis, hyalinis, glabris, simplicibus, obtusis, longis, c. 4 μ . cr. obsessa, 1,2-2 mm. diam.; ascis clavatis, 8-sporis, 6 \approx 4, apice jodo cœrulecente; sporidiis bacilliformibus, obtusis, rectis v. subcurvatis, 1-2-cellularibus, 9-15 \approx 1-1,5, hyalinis, distichis; paraphysibus filiformibus, teneris, apice, subin-crassatis.

Hab. ad caules putrescentes Aconiti Napelli infra alpem Retyezát Hunyad Transsylvaniae (LOJKA). — Species videtur propria, ad *T. leucophœam* vergens.

21. **Trichopeziza Berkeleyi** (Blox.) Sacc. *Peziza Berkeleyi* Blox. in 1673

B. et Br. Ann. N. H. n. 770. — Gregaria, minuta sessilis; ascomatibus hemisphæricis, furfuraceo-floccosis, plerumque pruinosis, margine primo inflexo, hymenio concavo alutaceo; ascis clavatis; sporidiis oblongis v. subcymbiformibus, 7-10 μ . longis (Berk.), 5-8 \approx 2 (Phill.).

Hab. in caulis emortuis Umbelliferarum in Britannia.
— Ascoma juvenile ore radiato-lineolato. Pili ascom. brevissimi, squamulis amorphis intermixti.

22. **Trichopeziza Grevillei** (Berk.) Sacc. *P. Grevillei* Berk. Engl. Fl. 1674

V, p. 198, *Mollisia Grevillei* (Berk.) Phill. Disc. p. 180. — Sessilis, gregaria, minuta, ascomatibus farinoso-tomentosis (subhir-sutis), pallide umbrinis, margine pallidore fimbriato; disco concole; ascis tereti-clavatis; sporidiis oblongo-fusoideis, rectis curvulisve, 3-guttulatis, 6-8 \approx 2; paraphysibus filiformibus.

Hab. in caule emortuo Umbelliferarum pr. Edinburg et Berwick in Britannia.

23. **Trichopeziza pilimoniliformis** (Cr.) Sacc. *Peziza pilimoniliformis* 1675

Cr. Fin. p. 50. — Ascomatibus 1-2 mm. diam., grise o-brunneis, sessilibus, concavis, costatis, costis e pilis moniliformibus formatis; ascis clavatis, octosporis; sporidiis subpiriformibus; paraphysibus filiformibus.

Hab. in caulis emortuis *Œnanthes crocatæ* et *Rubi*, Finistère Galliæ.

24. **Trichopeziza setigera** (Phill.) *Peziza setigera* Phill. Grev. VII, 1676

p. 22. — Gregaria, sessilis, minuta, concava, extus pilis brunneis, septatis, rigidis tecta; disco brunneo, carneo; ascis clavatis, octosporis; sporidiis oblongis, 16 \approx 3,5; paraphysibus latis, asco longioribus, sursum acutis.

Hab. in caulis emortuis Araliæ in California Amer. bor.

25. **Trichopeziza pulveracea** Fuck. Symb. mycol. pag. 297, non *Peziza Pteridis* Alb. et Schw. 1677

— Punctiformis, ceraceo-mollis, intus flavidula, extus atro-olivacea, granuloso-puberula, madore aperta, sicca urceolata; ascis oblongis, sessilibus, 8-sporis, 24 \approx 6; sporidiis oblongo-clavatis, hyalinis, 8-10 μ . long.

Hab. in caulis siccis Spirææ Ulmariæ in Germania, Helvetia, Belgio.

26. **Trichopeziza candidata** (Cooke) Sacc. *Peziza candidata* Cooke 1678

Grev. I, p. 130. — Niveo-alba, sessilis, mollis, sparsa v. subgregaria, mox plana, margine saepe lobato atque irregulari leniter

408 Discomyceteæ, Pezizeæ, Hyalosporæ, Trichopeziza.

elevato, extus setulis mollibus tomentosis ornata; ascis subclavatis; sporidiis minutis $7 \times 1,5$, obtusis, hyalinis.

Hab. in sarmentis Rubi, Highgate Britanniæ.

27. **Trichopeziza Ulmariae** (Lasch) Lamb. Myc. Belg. II, p. 524, Pez. 1679

Ulmariae Lasch in Herb. myc. n. 1723 et in Flora 1853, p. 201.

— Sparsa, minuta; ascomate subgloboso, dein aperto umbrino, pilis brevibus villoso; disco melino; ascis cylindraceis; sporidiis cylindraceis, minutis.

Hab. in caulis emortuis Spirææ Ulmariae in Germania, Britannia et Belgio. — *T. Spirææ* Kirchn. Lot. 1856, pag. 246: affinis sed major. Cl. Phillips ut var. *Pyren. atratæ* censem. Cfr.

28. **Trichopeziza cinerella** (Cr.) Sacc. Pez. *cinerella* Cr. Fin. p. 51. 1680

— Ascomate 1-2 mm. diam., griseo, tomentoso, sessili, primo sphæroideo, dein hemisphærico; ascis clavatis, 8-sporis; sporidiis piriformibus, hyalinis; paraphysibus subcylindraceis.

Hab. in caulis emortuis Ruborum, Finistère Galliæ.

29. **Trichopeziza subochracea** (C. et P.) Sacc. *Peziza subochracea* 1681

C. et P. Grev. I, p. 6, t. I, fig. 4. — Sessilis, sparsa, subglobosa, dein expansa, margine incurvo lutescente, tomentosa, disco obscuriori vel nitide ochraceo-lutescente; ascis cylindraceis; sporidiis acute fusiformibus v. subcylindraceis, 12μ . longis.

Hab. in caulis Rubi odorati, Adirondack Amer. bor.

30. **Trichopeziza Nidulus** (S. et K.) Fuck. Symb. p. 297, *Peziza Ni-* 1682

dulus Schm. et Kunze exs. n. 72, Fr. Syst. Myc. II, pag. 104, *Lachnum Nidulus* (S. et K.) Karst. M. F. I, pag. 181, *Lachnea Nidulus* Gill. Disc. c. icon. — Sparsa aut subgregaria, sessilis, subhemisphærica, badio-strigoso-hirtella, disco pallido-flavescente, margine obtusiusculo connivente, subfimbriato, 0,6 mm. lat.; ascis cylindraceo-subclavatis, $40-60 \times 4-4,5$; sporidiis fusoideo-filiformibus v. fusoideo-elongatis, rectis, distichis, $8-12 \times 1-1,5$; paraphysibus supra acutis, $3,5-4,5 \mu$. cr.

Hab. ad caules putrescentes Spirææ Ulmariae, Rubi saxytilis Polygonati, Epilobii angustifolii in Fennia, Suecia, Germania, Gallia, Belgio, Italia, Helvetia, Britannia. — Pili crebri septati, recti, brunnei, sursum incrassati, obtusi, $70-120 \times 5-8$.

31. **Trichopeziza Tami** (Lamy) Sacc. *Peziza Tami* Lamy in Desm. 1683

Descr. Espec. Nouv. 1836, p. 10. — Sparsa, sessilis, minutissima, globoso-applanata, tenerrima, subtomentosa, primo albo-hyalina, dein fulva; ascis cylindraceis; sporidiis ovoideis.....

Hab. ad caules putrescentes Tam i communis in Gallia. — *Var. Humuli* Phill. Disc. p. 270: gregaria sessilis, hemisphærica, dein expansa, putrescens, alba v. pallide-flava; ascis clavatis; sporidiis fusoideis vel suboblongis, 8-2; paraphysibus filiformibus parcis. In caulibus Humuli Lupuli, Darenth Britanniæ. Ascoma 200-400 μ . lat. Pili brevissimi, hyalini, continui.

32. **Trichopeziza ustulata** (Boud.) Sacc. *Helotium ustulatum* Boud. 1684 in B. S. Fr. 1881, p. 94, t. III, f. 5. — Ascomate applanato, late marginato, margine inæqualiter dentato, 0,6-1 mm. lat., extus ochraceo pilisque articulatis adpressis et in costulas subfuscas conniventibus; carne grisea; disco plano, livido vel ochraceo-plumbeo; ascis minutis octosporis, subfusoideis, 76-80 \approx 10; sporidiis fusideo-elongatis, rectis, curvulisve, utrinque 5-7-guttulatis, 18-22 \approx 3-5; paraphysibus bacillaribus.

Hab. in caule putri Polygonati multiflori gregatim Montmorency Galliæ.

33. **Trichopeziza earoleuca** (B. et Br.) Sacc. *Pez. earoleuca* B. et 1685 Br. Fung. of Ceyl. n. 943. — Sessilis, e globoso aperta, extus villosa, nivea; disco carneo, demum convexo; sporidiis minimis, oblongis 4 μ . longis.

Hab. in caulibus herbaceis, ramentis in ins. Ceylon. — Minuta, gregaria.

2. Ramicolæ et corticolæ.

34. **Trichopeziza myricacea** (Peck) Sacc. *Peziza myricacea* Peck 30 1686 Rep. p. 59. — Ascomatibus minutis, siccis, 500-700 μ . diam., sessilibus v. substipitatis, alutaceo-brunneis v. subcervinis, dense pilosis, udis expansis et discum albescensem ostendentibus; ascis subcylindraceis, circiter 40 μ . longis; sporidiis minutis, spermatoideis; paraphysibus ascum latitudine æquantibus et multo longioribus, sursum acutis.

Hab. in ramis Myricæ Gales, Adirondack Amer. bor. — Afinis *T. brunneolæ* a qua magnitudine differt. Pili longi, continui apice non incrassati. Sicca, omnino clausa.

35. **Trichopeziza antarctica** (Speg.) Sacc. *Lachnella antarctica* Speg. 1687 F. Fueg. n. 319. — Densiuscula sparsa, sessilis v. vix substipitula, minutissima, uda applanato-concava, 0,5-1 mm. diam, sicca globuloso-contracta, candida, villosula, pilis longiusculis cylindraceis 150-250 \approx 4-5 μ . apice obtusiusculis basi non incrassatis parciissime septulatis levissimis flexuosulis dense vestita; disco levi

albo-ceraceo; ascis cylindraceo-subfusoideis, apice truncatis, deorsum breviuscule crasseque pedicellatis, 60-65 \approx 5, octosporis, pseudo-paraphysibus clavulatis, apice longiuscule attenuatis acutissimis non v. vix longioribus, 65-70 \approx 3-4, hyalinis parce obvallatis; sporidiis subfusoideis, utrinque acutiusculis, 10 \approx 1-1,5, rectis v. curvulis, continuis, hyalinis.

Hab. in ramulis emortuis decorticatis *Mayteni magellanicæ* in Gebel island. — Contextus excipuli subprosenchymaticus, pallide melleus; jodi vis nulla. Species *Cyphellæ albo-violascenti* simillima, partibus fructificationis inspectis tantum dignoscenda; peraffinis etiam *T. mollissimæ* Karst.

36. **Trichopeziza obscura** (Cooke) Sacc. *Peziza obscura* Cooke Grev. 1688 VII, pag. 3. — Sessilis, sparsa, atro-fuliginosa; cupula pubescens, demum applanata, margine elevato, disco aquoso-cinereo, $\frac{1}{2}$ -1 mm.; ascis cylindraceis; sporidiis linearibus, 10 μ . long.; paraphysibus filiformibus.

Hab. in ramis *Quercus*, Sierra Nevada Californiæ.

37. **Trichopeziza bellula** (Karst.) Sacc. *Lachnum bellulum* Karst. 1689 M. F. I. pag. 177. — Sparsa v. subgregaria, sessilis, planiuscula, sicca concava, sulphureo-strigoso-villosa, disco pallido, 1 mm. lat.; ascis cylindraceo-clavatis, 60-66 \approx 5,5-6; sporidiis fusoideo-filiformibus, rectis v. leviter curvulis, guttulis 2-6 majusculis fœtis, 8-18 \approx 1-2; paraphysibus apice acutiusculis, 3 μ . cr.

Hab. ad ramulos decorticatos *Fraxini excelsioris* prope Observatorium oppidi Abo in Fennia.

38. **Trichopeziza grandinea** (Quél.) Sacc. *Lachnea grandinea* Quél. 1690 10 Suppl. p. 12, t. XI, f. 3 et Grev. 1879, p. 37. — Ascomatibus turbinatis, 1 mm. diam., granuliformibus, firmis, costulatis, subtilater lanosis, albo-niveis; hymenio concavo, albo; sporidiis bacillaribus 12 μ . longis, 5-guttulatis spurieque septatis.

Hab. gregatim in corticibus *Populi tremulæ*, Jura.

39. **Trichopeziza glandulosa** (Gill.) Sacc. *Lachnea glandulosa* Gill. 1691 Disc. pag. 83, c. ic. — Sessilis, hemisphærica, orbicularis, rarius subovata, uda mox aperta, sordide alba vel subrosea, pilis apice capitato-glandulosis albidis præcipue margine obsessa, 1 mm. diam., disco pallido; ascis teretiusculis; sporidiis fusoideo-ellipsoideis, 2-3-guttulatis, hyalinis; paraphysibus filiformibus.

Hab. in ramis emortuis *Cerasi*, gregatim, in Gallia. bor. —

An *Lachnellæ papillaris* forma?

40. **Trichopeziza necrioidea** Rehm Ascom. Alp. n. 14. — Sparsa v. 1692

gregaria, sessilis, primitus globosa, dein basi applanata, extus albo-pilosa, sicca clausa, humida urceolata, margine albido, subtilissime fimbriato, primitus clausa, disco hyalino, c. 0,3-5 mm. diam., contextu prosenchymerico, flavidulo, pilis creberrimis, simplicibus, acutis, basi septatis ibique 4-5 μ . cr., rectis, glabris, hyalinis obsessa; linis obsessa; ascis clavatis, 8-sporis, 45 \approx 7; sporidiis ovatis, hyalinis, 1-cellularibus, 1-2-stichis, 6-7 \approx 3; paraphysibus filiformibus.

Hab. ad ramulum siccum Rhododendri ferruginei, Peischlkopf in alpibus tyrolensis, c. 1800 m. alt. (BRITZELMAYR).

41. **Trichopeziza fusco-hyalina** Rehm Ascom. Alp. n. 13. — Gre. 1693
garia, sessilis, primitus globoso-urceolata, dein subdilatata, strigosa, fusca, quasi villo subexplanata insidens, humida explanata orbicularis, concavula, disco et margine hyalino, 1-1 $\frac{1}{2}$ mm. diam., contextu fusco, prosenchymerico, pilis simplicibus, longis, septatis, fuscis, subundulatis, glabris, c. 4 μ . cr. obsessa; ascis clavatis, apice obtuse acuminatis, 8-sporis, 45 \approx 8; sporidiis oblongis, obtusis, rectis, 1-cellularibus, hyalinis, distichis, 9 \approx 4; paraphysibus filiformibus, simplicibus, c. 4 μ . cr., haud clavatis.

Hab. ad ramum siccum Rhododendri ferruginei, Peischlkopf in alpibus tyrolensis c. 1800 alt. (BRITZELMAYR).

42. **Trichopeziza Kalmiæ** (Peck) Sacc. *Peziza Kalmiæ* Peck 25 Rep. 1694
p. 99. — Ascomatibus minutis, sessilibus vix planis, marginatis, extus furfuraceis et obscure griseis, margine primo incurvo; disco rufo-brunneo; sporidiis ellipsoideis, saepe nucleatis, 10 \approx 5.

Hab. in ramis Kalmiæ angustifoliæ, Sandlake Amer. bor.

43. **Trichopeziza concolor** (Speg.) Sacc. *Peziza concolor* Speg. F. 1695
Arg. Pug. IV, p. 87. — Sessilis, applanato-patellaris, minor 0,5-0,8 mm. diam., margine repando, disco ceraceo-pallescidente, excipulo fusco-fulvescente, puberula, contextu celluloso-parenchymatico, in pilis concoloribus, parvulis 40-70 \approx 5, 1-2-septulatis producto; ascis cylindraceis, apice truncatis, basi longiuscule attenuato-stipitatis, 70-75 \approx 5-7; paraphysibus filiformibus obvallatis, octosporis; sporidiis distichis, cylindraceis rectis vel inaequilateralibus, utrinque obtusiusculis, granuloso-farctis vel 2-guttulatis, 12 \approx 2-3, hyalinis; obturaculo minutissimo ascorum iodi ope cœrulecente.

Hab. in ramis corticatis dejectis Cephalanthi sarandi, in dumetis uliginosis, Conchas Argentinæ. — Forte ad *Pirottæam* vergit.

44. **Trichopeziza chrysotricha** (Berk.) Sacc. *Peziza chrysotricha* 1696
Berk. Fl. New. Zeal. p. 201, t. CV, f. 7. — Cupula sessili, pri-

mum globosa, demum subhemisphærica, aurea, floccis innatis intricatis vestita subpulverulenta; ascis crasse cylindraceis; sporidiis brevibus cymbiformibus, 11–12 \approx 4.

Hab. in ramis pr. Waikare Lake Novæ Zelandiæ (COLENZO). — Ascoma 2–3 min. lat.

45. **Trichopeziza Cookii** (Pass.) Sacc. *Peziza Cookii* Pass. Diagn. 1697

Fung. nuovi p. 258. — Minuta, subgregaria, sessilis, mollis, pallide rosea; cupula demum applanata, extus margineque floccis brevibus hyalinis fasciculatis obsita, disco concavo; ascis parvis cylindrico-clavatis, saepe longe et flexuose stipitatis; sporidiis linearibus, minutis, spermatoideis.

Hab. in cortice interiore emortuo Piri Mali, prope Parma Ital. bor. — Mollitie *Mollisiam* appropinquat, sed indumenti gratia ad *Trichopezizas* adscribenda.

46. **Trichopeziza cœrulescens** (Rehm) Sacc. *Dasycephala cœrulescens* 1698

Rehm Ascom. Lojk. n. 24. — Gregaria, sessilis, primitus globosa, dein urceolata, denique applanata, margine spurio circumdata, cœrulescens, c. 0,5–6 mm. diam. prosenchymatico-contexta, pilis brevibus, scabris, obtusis, simplicibus, 4–6 μ . cr. obsessa; ascis clavatis, 24–30 \approx 3–4, 8-sporis, jodi ope apice cœrulecente; sporidiis oblongatis, obtusis, 1-cellularibus, hyalinis, 4–5 \approx 1, distichis; paraphysibus simplicibus, filiformibus, teneris, haud clavatis.

Hab. ad corticem Betulæ prope Chodowice, Stryj, Galiciæ australis (LOJKA).

47. **Trichopeziza ulmicola** (Cr.) Sacc. *Peziza ulmicola* Cr. Fin. p. 52. 1699

— Ascomate $\frac{1}{2}$ mm. diam., sessili, carneo, dein ferrugineo et parum lobato, margine pilis coloratis, uncinatis, fasciculatis ciliato; ascis subclavatis, 8-sporis, ovoideis, binucleatis; paraphysibus filiformibus, asco longioribus.

Hab. in cortice emortuo Ulmi, Finistère Galliæ.

48. **Trichopeziza cæsio-lutea** (B. et Br.) Sacc. *Peziza cæsio-lutea* 1700

B. et Br. Fung. of Ceyl. n. 948. — Sessilis, saepe congesta, crispata, e strato tenui nigro oriunda, extus cœruleo-tomentosa; hymenio flavo; ascis clavatis 25 μ . long.; sporidiis minutis, monostichis, 5 μ . long.

Hab. in corticibus, Central provincia Ceylon. — *L. Godroniae-næ* affinis dicitur.

3. Lignicolæ.

49. **Trichopeziza albo-lutea** (Pers.) Sacc. *Hel. albo-luteum* (Pers.) 1701

Karst. M. F. I, p. 160, *Peziza albo-lutea* Pers. Icon. et descr. Fung. p. 31, t. 8, f. 4 et 5, *Pez. variecolor* Fr. Syst. Myc. II, p. 100, *P. sulphureo-cæsia* Schum. Flor. Saell. p. 432, *P. hydnoides* Sow. t. 178. — Gregaria, sessilis, hemisphærica, sicca clausa, sulphureo- vel luteo-virescenti-villosa, disco concaviusculo vel demum plano, pallido vel albo-pallescidente, 0,5-1,1 mm. lat.; ascis cylindraceo-clavatis, 45-50 \approx 6; sporidiis oblongatis, eguttulatis, 7-11 \approx 2-3; paraphysibus gracilibus.

Hab. supra lignum putridum locis humidis in Fennia, Suecia, Dania, Germania, Italia, Gallia, Belgio, Amer. bor., Cuba. — Pili cupulæ flavi, 2,5-3 μ . cr. Defectu subiculi, ut opinor, a *Tapesia fusco-umbrina* distinguenda. Cfr. hanc speciem.

50. **Trichopeziza grisella** (Cooke et Phill.) Sacc. *Lachnella grisella* 1702
C. et Ph. Disc. p. 260. — Gregaria, sessilis, rarius substipitata, turbinata dein hemisphærica, dilutissime brunnea, furfuraceo-tomentosa, longitrorsum striatula; margine albido fimbriato; disco brunneo; ascis clavatis; sporidiis oblongo-fusoideis, 4-7 \approx 1-1,5; paraphysibus filiformibus.

Hab. in ligno emortuo decorticato in Britannia (CARMICHAEL). — Ascoma 100-500 μ . lat.; pili vix colorati hinc parum conspicui.

51. **Trichopeziza flavo-fuliginea** (A. et S.) Sacc. *Peziza flavo-fuliginea* A. et S. p. 319, t. XI, f. 7, Pers. Eur. p. 251, *Dasyscypha flavo-fuliginea* Fuck. Symb. App. I, pag. 137. — Sessilis, dilatata, planiuscula, mollis, strigoso-hirta, læte sulphurea, intus olivaceo-fuliginea; ascis oblique stipitatis, cylindraceis, 8-sporis, 88 \approx 8; sporidiis in asci superiori parte congregatis, cylindraceo-fusiformibus, sed utrinque obtusis, rectis, continuis, hyalinis 10 \approx 3; paraphysibus filiformibus.

Hab. ad lignorum frusta, folia etc. madida putrida dejecta in paludosis silvarum in Germania et Carolina. — Gregaria nitidula, initio cupularis, dein varie flexuosa applanata, 2-5 mm. lata, suborbicularis, oblonga etc. Colore et substantia tenera ab affinibus differt.

52. **Trichopeziza crossata** (Ellis) Sacc. *Lachnella crossata* Ellis Bull. 1704
Torr. Cl. 1881, p. 124. — Sessilis, grisea, primo sphæroidea et ore minuto aperta, dein expansa et concava, margine pilis griseis, rectis, 4-6-septatis, obtusis circiter 75 \approx 7 fimbriato; disco udo lido, sicco pallide griseo; ascis cylindraceis, in stipitem brevem abrupte desinentibus; paraphysibus linearibus, obscure septatis, fir-

mis sursum non incrassatis; sporidiis allantoideis, oblongis, hyalinis, continuis, $7-9 \approx 3$.

Hab. in ligno vetusto, New Mexico Amer. bor.

53. **Trichopeziza confusa** Sacc. *T. punctiformis* Rehm Asc. n. 309, 1705
Sacc. Mich. I, p. 66 nec II, pag. 140, *Peziza punctiformis* Pat.
Tab. f. 89. — Gregaria, punctiformis, alba, ex urceolato scutella-
ta, sessilis, vix $\frac{1}{3}$ mm. lat., villosa; pilis simplicibus, fragilibus,
acutis, rugulosis, hyalinis, 6μ . cr.; ascis clavatis, 8-sporis $18 \approx 7$;
sporidiis distichis oblongis, rectis, non guttulatis, $6 \approx 3$; paraphysi-
bus (ex Pat.) filiformibus.

Hab. in ligno putri in Germania, Gallia et Italia bor.

54. **Trichopeziza vitricolor** (B. et Br.) Sacc. *Peziza vitricolor* B. et 1706
Br. Ann. N. H. n. 1321, t. XIX, f. 14. — Ascomatibus brevissime
stipitatis vel sessilibus, carnosо-ceraceis, turbinatis, tenuiter to-
mentosis, citrinis; ascis 85-100 μ . longis; sporidiis fusiformibus
guttulis oleosis præeditis 20-25 $\approx 5-6$; paraphysibus linearibus.

Hab. in ligno putri in Britannia. — Ascoma 2 mm. lat.

55. **Trichopeziza dematiicola** (B. et Br.) Sacc. *Peziza dematiicola* 1707
B. et Br. Ann. N. H. n. 1070, t. 15, f. 20. — Gregaria, minuta;
ascomate hemisphærico, umbrino, pilis longis hyalinis puberulo ci-
liatoque; disco cinereo; ascis breviusculis; sporidiis subcym bifor-
mibus, hyalinis, $10 \approx 2,5$; paraphysibus filiformibus.

Hab. in ligno putri inter hyphas *Helminthosporii* speciei in
Britannia. — Ascoma 100-500 μ . diam.

56. **Trichopeziza viridicoma** (Peck) Sacc. *Peziza viridicoma* Peck 1708
31 Rep. p. 46. — Ascomatibus villosis, minutis, sessilibus, viridi-
lutescentibus; sporidiis stipatis vel distichis, oblongis vel subfu-
siformibus, $12-15 \approx 5-6$.

Hab. in ligno putri dejecto, Sandlake Amer. bor.

57. **Trichopeziza leuconica** (Cooke) Sacc. *Lachnella leuconica* Cooke 1709
in Phill. Disc. p. 267. — Gregaria, sessilis, expansa, submembrana-
cea, alba; pilis longis tenuibus attenuatis, flexuosis obsessa; margine
elevato-ciliato; ascis tereti-clavatis; sporidiis oblongis vel oblon-
go-ellipsoideis rectis curvulisve $8 \approx 2$; paraphysibus filiformibus.

Hab. in ligno putri, Clove Britanniæ. — Ascoma 0,5-1 mm.
lat. sinuosum tenue; pili hyalini continui.

58. **Trichopeziza fusca** (Schum.) Sacc. *Peziza Schumacheri* Fr. S. 1710
M. II, p. 98, Fl. Dan. t. 1755, *Lachnella Schumacheri* (Fr.) Phill.
Disc. p. 262, *P. fusca* Schum. Saell. p. 434 nec Pers. *P. cœru-
lescens* Schum. loc. cit. (disco cœruleo tantum recedit), Pers. Myc.

Eur. p. 252, *P. plumbea* Grev. t. 11 (disco plumbeo). — Sessilis, exigua, hemisphærica, tomentosa, fusca, disco planiusculo fusco-purpurascente; ascis tereti-clavatis; sporidiis fusoideis v. tereti-fusoideis, rectis curvulisve, 8-13 = 2-3; paraphysibus filiformibus.

Hab. in ligno putrido Betulæ etc. in Dania, Britannia. — Conferta, 1-1,5 mm. lat., margine humido elevato integro cincta. *Habitus omnino Mollisicæ cinereæ.*

59. **Trichopeziza Richonii** (Boud.) Sacc. *Urceolella Richonis* Boud. 1711

Bull. Soc. Myc. IV, pag. 82, t. 17, f. 3. — Minutissima, 0,2-0,5 mm. lata, tota rubra, pubescens, gregaria, sessilis, carnosa, urceolata, extus præcipue ad marginem villosa, pilis acutis, septatis, 50-120 μ . longis, ad basim 3-5 crassis, apice dilutioribus, hymenio et carne concoloribus; paraphysibus tenuibus, filiformibus, parum septatis, ad apicem non incrassatis, hyalinis, 2 μ . circiter crassis; ascis minoribus, late clavatis, octosporis, 40-45 = 9; sporidiis minutis, elliptico-fusoideis, intus biguttulatis 5-7 = 2-2,5.

Hab. lignum putridum Populi in Gallia (RICHON).

60. **Trichopeziza venatoria** (Speg.) Sacc. *Peziza venatoria* Speg. F. 1712

Arg. IV, n. 222. — Sessilis, uda appланato-patellaris, 0,7-1,5 mm. diam., disco concaviusculo roseo-pallescens, sicca margine inflexo albo, disco sordide rufo-fulvescente, crassiuscula, extus fuligineo-grisea, puberula; contextu excipuli celluloso-parenchymatico; pilis plus minusve elongatis, saepius brevissimis et subclavulatis, parce septulatis, pallide fuligineis; ascis octosporis, cylindraceis vel cylindraceo-clavatis, apice truncato-rotundatis, deorsum longe attenuato-stipitatis 85 = 6-7; paraphysibus filiformibus guttulatis obvallatis; sporidiis distichis, cylindraceo-ellipticis, utrinque rotundato-truncatis, eguttulatis, 8-11 = 3, continuis hyalinis.

Hab. in trunco putrescente decorticato dejecto Erythrinae Cristæ-galli in nemrosis uliginosis, Conchas Argentinæ.

61. **Trichopeziza paradoxa** (Karst.) Sacc. *Lachnum paradoxum* 1713

Karst. M. F. I, p. 168. — Gregaria, ut plurimum confluens, sessilis, tuberculosa, fulvo-vel rufo-testacea vel sub testacea, 0,2-0,6 mm. lat.; ascis fusoideo-cylindraceis 30-36 = 4; sporidiis vulgo distichis filiformibus vel bacillaribus, guttulas plures, plerumque 4 foventibus vel spurie tenuiter septatis, 6-10 = 1-1,5; paraphysibus parum notabilibus.

Hab. supra lignum Quercus vetustum in insula Ransala pr. oppid. Abo Fenniae.

62. **Trichopeziza albo-viridis** (Cooke) Sacc. *Pez. albo-viridis* Cooke 1714

in Grev. VII, p. 47. — Sparsa, sessilis, villosa; cupulis hemisphæricis, ærugineo-villosis, $\frac{1}{2}$ - $\frac{3}{4}$ mm.; margine albido, disco ochraceo, concavo; pilis asperulis, septatis, viridi-tinctis; ascis cylindraceis; sporidiis linearibus, obtusis, rectis 8-10 μ . longis; paraphysibus filiformibus.

Hab. in trunco decorticato vel corticis rimis Myricæ, Aiken Carolinæ.

4. Fructicolæ et foliicolæ.

63. **Trichopeziza Aranea** (De Not.) Sacc. Mich. I, p. 253, f. XIII, P. 171
Aranea De Not. Micr. ital. Dec. I, n. 1. — Cupulis sessilibus, minutis apertis, candidis, 200-300 μ . diam., extus pilis hyalinis, flexuosis longiusculis undique vestitis, excipulo prosenchymatico hyalino; ascis oblongo-clavatis, brevissime noduloso-stipitatis, 35-40 \approx $5\frac{1}{2}$ -6; paraphysibus filiformibus longioribus, 50-55 \approx 1, obvalbatis, octosporis; sporidiis distichis oblongis, 7-8 \approx 1-2, utrinque obtusiusculis, hyalinis.

Hab. in capsulis dejectis putrescentibus Castaneæ vescæ, præcipue ad parietes interiores in Italia bor., Gallia, Britannia et Amer. bor. — Forma americana præbet ascos 65-70 \approx 6-7; sporidia tereti-clavulata 8 \approx 2,5.

64. **Trichopeziza punctiformis** (Fr.) Fuck. Symb. myc. p. 296, Sacc. 171
 F. it. f. 1494, *Peziza punctiformis* Fr. Syst. Myc. II, p. 105, P.
dryophila Pers. M. E. I, p. 265. — Sessilis, perexigua, punctiformis, subfugax, villosa, ore subconnivente; pilis filiformibus, apice capitulato, septulato, albido; ascis clavulatis, octosporis, 25-28 \approx 4; sporidiis distichis, tereti-oblongis rectis curvulisve, 3-guttulatis, hyalinis, 5-6 \approx 1-1,5; paraphysibus filiformibus.

Hab. ad folia decidua putrescentia Myricæ, Alni, Quercus in Germania, Belgio, Italia, Amer. bor. — Gregaria, superficialis, minus persistens, semper fere clausa, alba, demum badio-fuscens. Typus speciei est foliicola.

65. **Trichopeziza pollinaria** (Cooke) Sacc. *Peziza pollinaria* Cooke 171
 Disc. Un. St. I, p. 292. — Epiphylla, subgregaria, minuta, sessilis, pallida, pilis albis pulveraceis, brevibus, farinam albam simulantibus tecta, ascomate globoso, dein poro centrali minuto pertusa; ascis cylindraceis; sporidiis ellipsoideis, minutis.

Hab. in foliis quercinis, New Jersey Amer. bor. (ELLIS).

66. **Trichopeziza marginata** (Cooke) Sacc. *Peziza marginata* Cooke 171
 Hedw. 1875, p. 82, Disc. U. S. I, p. 292. — Epiphylla sparsa v.

gregaria, subbrunnea, sessilis, cupula turbinata, dein aperta, minuta, margine floccis septatis, brunneis ornata, disco pallidore; ascis cylindraceis; sporidiis minutis linearibus, spermatoideis.

Hab. ad folia decidua *Quercus*, *Andromedæ* etc. New Jersey Amer. bor. (ELLIS).

67. **Trichopeziza capitata** (Peck) Sacc. *Peziza capitata* Peck 30 Rep. 1719 p. 60 et in M. U. n. 813. — Cupulis minutis, sessilibus, candidis, in siccitate clausis et subglobosis, madefactis apertis et explanatis, extus pilis candidis capitatis vestitis; disco albido vel flavido; ascis cylindricis, 30 μ . long.; sporidiis acicularibus 5-7 μ . long.; paraphysibus crassis, ascis longioribus, apice acuminatis.

Hab. ad folia arida *Quercus albæ*, Albany N. Y. Amer. bor.

— Pili basi incrassata septata, apice capitata.

68. **Trichopeziza acerina** (C. et E.) Sacc. *Peziza acerina* C. et E. 1720 Grev. VII, pag. 40. — Hypophylla, sparsa, punctiformis; cupulis sessilibus, tenuibus, fuscis $\frac{1}{5}$ mm.; margine albopilosus; ascis subclavatis; sporidiis linearibus, 6 μ . long.

Hab. in foliis *Aceris*, New Jersey Amer. bor. — Pili marginales minute rugulosi.

69. **Trichopeziza araneo-cincta** (Phill.) Sacc. *Lachnella araneo-cincta* Phill. Disc. p. 271. — Minuta sparsa, sessilis, concava, tenuis, pallide lutea, margine setis longis, flexuosis, acutis, hyalinis ornata; ascis late clavatis, octosporis; sporidiis distichis, stricte fusiformibus, utrinque acutis, 10-13 \approx 1-1,5.

Hab. in foliis emortuis *Betulæ*, Clifton, Bristol Britanniæ (C. BUCKNALL). — Ascoma 200-300 μ . lat.

70. **Trichopeziza Karstenii** Sacc. *Lachnum radians* Karst. M. F. I, 1722 p. 177, nec Saut. — Sparsa, sessilis, tenuis, convexula, sicca concaviuscula et subanguloso-contracta, subtus fuscescens, sulphureo-strigoso-villosa, disco pallido, 0,5-1 mm. lat.; ascis cylindraceo-subclavatis, 60-66 \approx 4-5,5; sporidiis fusoideo-elongatis, simplicibus plerumque leniter curvulis, 8-11 \approx 1,5-2; paraphysibus apice acutiusculis, 3 μ . cr.

Hab. supra folia *Populi tremulæ* semiputrida prope Mustiala Fenniæ.

71. **Trichopeziza ciliata** (Fuck.) Sacc. *Hyalopeziza ciliata* Fuck. 1723 Symb. mycol. p. 298. — Cupulis sparsis, sessilibus, carnosus, usque 1 mm. ad latis, primo globosis, demum appanatis, dilute carneo-flavis, disco parum obscuriore, glabris sed margine longissime patentem ciliatis, ciliis cupulæ diametro longioribus, hyalinis; ascis ob-

longis, curvato-stipitatis, 8-sporis, $32 \pm 6 \mu$. cr.; sporidiis oblique monostichis, oblongo-ovatis, 6 ± 3 , hyalinis.

Hab. in foliis siccis Carpin i Betuli, raro, pr. Reichartshausen Rhenogoviæ.

72. **Trichopeziza lachnobrachya** (Desm.) Sacc. *Peziza lachnobrachya* 1724

Desm. 19 Not. pag. 27. — Amphigena, sparsa aut laxe gregaria, sessilis, minutissima, tenerrima, albido-hyalina, primo concava, dein applanata, extus villo delicatissimo albo vestita, subtus puncto nigro affixa; ascis clavatis, brevibus, subcrassis, $30-35 \mu$. long.; sporidiis ellipsoideis.

Hab. in foliis exsiccatiis arborum in Gallia. — Ascoma $\frac{1}{4}$ mm. diam.

73. **Trichopeziza eximia** Mont. et Sacc. (sp. nov.). — Ascomatibus 1725

gregariis, obconico-cupulatis, subsessilibus, $\frac{1}{2}-\frac{3}{4}$ mm. diam., paleo lide citrinis, pilis capitatis rubro-aurantiis eximie ciliatis; ascis subteretibus 40 ± 3 ; sporidiis subovoideis $6-7 \pm 1,5$ hyalinis.

Hab. in acubis putrescentibus ramulorum resectorum Pin i silvestris prope Liege Belgii. — A *L. solfatara* C. et Ell. ascomatibus subcitrinis pilorumque capitulis amœne rubro-aurantiis differre videtur.

74. **Trichopeziza misella** (Rob. et Desm.) Sacc. Mich. II, p. 80, *Peziza misella* Rob. et Desm. 14 Not. p. 186. — Hypophylla, sessilis, minuta, numerosa, primo globosa, dein aperta, suborbicularis, extus pilis nigris brevioribus hirsuta, disco plano, dein convexiusculo, cinereo-pallescens; margine sinuoso ciliato albo; ascis clavatis; sporidiis oblongis.

Hab. in foliis exsiccatiis Rubi fruticosi in Gallia. — Ascomata (in exempl. pr. Rouen lect. a cl. MALBRANCHE) minutissima, subsessilia, aperta, albida dein subargillacea, prope marginem pilis fuligineis $5-6 \mu$. cr. vestita; ascii cylindracei 48 ± 5 , paraphysati, subsessiles; sporidia cylindraceo-clavulata, $7-8 \pm 1\frac{1}{2}-2$ hyalina.

75. **Trichopeziza dumorum** (Desm.) Sacc. Mich. II, p. 80, Rehm Asc. 1727 n. 611, *Peziza dumorum* Rob. et Desm. XVIII, Not. pag. 4. — Ascomatibus sparsis, minutissimis, subsessilibus, plano-umbilicatis, brunneo-lividis margine albicante, extus pilis brunneis hirtellis, disco concavo, pallido; ascis clavulatis $27-29 \pm 4-5$; sporidiis subfusoideis $5-8 \pm 1\frac{1}{2}-2$, hyalinis; paraphysibus filiformibus.

Hab. in pag. infer. foliorum emortuorum v. languidorum Rubi fruticosi in Gallia, Britannia, Germania. — Ascoma $\frac{1}{5}-\frac{1}{3}$ mm. lat. Affinis *T. misellæ*.

76. **Trichopeziza labrosa** (Phill. et Harkn.) Sacc. *Lachnella labrosa* 1728

Phill. et Harkn. Calif. p. 1. — Sparsa, sessilis, olivaceo-brunnea, pilis brevibus, brunneis, septatis tecta, margine involuto; disco pallide lutescente; ascis cylindraceo-clavatis octosporis; sporidiis ovoideis, 10-14 \approx 3-6; paraphysibus linearibus.

Hab. in utraque pagina foliorum emortuorum *Arctostaphyli pungentis* in California. — Ascoma 1 mm. diam., siccum nigrescens compressum.

77. **Trichopeziza venturioides** (E. et E.) Sacc. *Lachnella venturioides* 1729

E. et E. Journ. Myc. 88, pag. 99. — Gregaria, sessilis, $\frac{1}{4}$ mm. diam., disco albescente, extus pilis rigidis, acutis, continuis, rectis, nigris, 60-80 \approx 4-5 tecta; ascis sessilibus, sursum obtusis, circiter 30 \approx 6; paraphysibus filiformibus; sporidiis distichis, clavato-fusoideis vel clavato-oblongis, hyalinis, continuis, circiter 10-12 \approx 1,5-2.

Hab. in foliis emortuis *Gaylussaciæ dumosæ*, Newfield in America boreali.

5. In Gramineis et Cyperaceis (culmi-foliicolæ).

78. **Trichopeziza albo-testacea** (Desm.) Sacc. Misc. myc. I, p. 8, *Peziza albo-testacea* Desm. 10 Not. p. 34, *Lachnum albo-testaceum* Karst. M. F. I, p. 175. — Sparsa, brevissime stipitata aut sessilis e subsphæroideo hemisphærica, sicca clausa, albo-testacea, albido-vel testaceo-ferrugineo-tomentoso-villosa, 1 mm. lat.; ascis cylindraceo-subclavatis, 45-60 \approx 4-5; sporidiis distichis vel oblique monostichis, fusoideo-filiformibus vel fusoideo-elongatis, rectis, interdum levissime curvulis guttulatis, 8-19, plerumque 9-12 \approx 1,5-2; paraphysibus grandiusculis, 4-5 μ . cr., apice acuto.

Hab. ad culmos *Secalis Cerealis*, *Airæ cæspitosæ*, *Baldingeræ arundinaceæ* et *Calamagrostidum* etc. putrescentes in Gallia, Fennia, Germania et Britannia. — Est quasi *Das. patens* sed subsessilis.

79. **Trichopeziza Secalis** (Lib.) Sacc. Mich. II, p. 330, *Peziza Secalis* Lib. Cooke Grevill. VIII, pag. 84. — Minuta, sparsa, sessilis, globosa, strigoso-tomentosa, sulphureo-fuscescens, disco roseo; sporidiis linearibus, minutis.

Hab. in culmis *Secalis Cerealis* in Arduennis. — Cl. Speg. et Roum. ducunt (Reliq. Lib. I, n. 651) ad *Lachnellam flammeam*, quæ robustior, lignicola. Teste Cooke, *Tap. mutabili* affinis. Ipse ascos maturos non vidi.

80. **Trichopeziza pyrostoma** (Mont.) Sacc. *Peziza pyrostoma* Mont. Cent. V, n. 87, Syll. Crypt. n. 626. — Caulicola, sparsa, minuta, badia, pilis appressis obtectis subglabra; ore connivente, pilis lutescenti-rufis clauso; ascis longe cylindricis, 70-80 μ . longis; sporidiis distichis oblongis, 3-guttulatis, 12 \approx 2,5.

Hab. in culmis secalinis, in Gallia centrali, apud Lemovicem (LAMY). — Ascoma $1/2$ mm. lat. *T. Nidulo* affinis.

81. **Trichopeziza prasina** (Quél.) Sacc. *Lachnella prasina* Quél. As- 1732 soc. Franc. 1880, p. 13, t. IX, f. 5, Phill. Disc. p. 261. — Ascomatibus sessilibus, applanatis, tenuibus, ceraceis, olivaceis, pilis flexuosis, hyalinis, demum luteo-igneis, disco plano viridi-brunneo, dein amœne viridi demum pallido; ascis tereti-clavatis; sporidiis fusiformibus, curvatis, 10-15 \approx 2-3.

Hab. in culmis Glyceriæ fluitantis in Normandia (LE BRETON), Phalaridis arundinaceæ, Chedder Britanniæ (BUCKALL).

82. **Trichopeziza Winteri** (Cooke) Sacc. *Pez. Winteri* Cooke Grev. IV, 1733 pag. 67, *Dasyascypha Kneiffii* Rehm Ascom. n. 113, nec *Peziza Kneiffii* Wallr. — Ascomatibus gregariis, sessilibus urceolatis, argillaceis, $1/2$ mm. diam., extus tomentosis; pilis continuis, luteolis, 6 μ . cr.; ascis clavatis, octosporis, 45 \approx 7, apice jodo cœruleoscentibus; sporidiis hyalinis, bacillaribus, rectis, continuis, 9 \approx 1,5, paraphysibus fusiformibus, acutis, hyalinis, 5 μ . crass.

Hab. in culmis putridis Phragmitis communis, Leipzig Germaniæ (WINTER).

83. **Trichopeziza horridula** (Desm.) Sacc. Mich. II, p. 80, *Peziza horridula* Desm. 14 Not. p. 185. — Caulicola, sessilis, sparsa, minuta, subglobosa, sæpius clausa subconnivens, strigoso-hirta; pilis longis, rectis, septatis rufis, apice albis; disco pallescente; ascis parvis, cylindricis, hyalinis, 50 \approx 5.

Hab. in culmis siccis Tritici, Magny Galliæ (BOUTEILLE).

84. **Trichopeziza insularis** (Speg.) Sacc. *Lachnella insularis* Speg. 1735 Fung. Fueg. n. 321. — Hinc inde laxe gregaria, sessilis, uda applanato-concava 0,5-1,5 mm. diam. sicca subgloboso - contracta, extus sordide rufescenti- vel griseo-testacea, contextu parenchymatico hyalino, pilis longiusculis gracilibus 200-300 \approx 3 cylindraceis flexuosulis rigidulis sursum non attenuatis apice obtusis basi non incrassatis 6-12-septatis non constrictis rufo-fuligineis; disco sordide ceraceo; ascis clavulatis, sursum cuneatis apiceque subtruncatulis, deorsum longiuscule attenuato-pedicellatis p. sp. 85 \approx 5, ped.

15-20 = 2-3, octosporis, pseudoparaphysibus longiusculis subclavulatis apice acutissimis 110-120 = 2-3 hyalinis obvallatis; sporidiis recte distichis v. oblique monostichis subfusoideis, utrinque attenuato-obtusatis 9-11 = 1,5 rectis v. subinæquilateralibus, hyalinis, continuis; an per ætatem medio 1-septatis?

Hab. ad culmos vaginasque emortua putrescentia *Hierochloæ antarcticæ* prope Penguin rookery in Staten island Fuegiæ. — Obturaculum ascorum minutissimum, iodi ope dilute cœrulescens.

85. **Trichopeziza agrostina** (Peck) Sacc. *Peziza agrostina* Peck 29 Rep. 1736

p. 55. — Ascomatibus sparsis, minutis, in sicco 500-750 μ . diam., subsessilibus hemisphæricis vel subglobosis, externe setosis, extus roseo-coloratis, setis marginalibus introflexis (udis) sæpe apice subulatis, longioribus, ceteris non subulatis, sæpe rugosis plus minusve septatis, disco pallido vel lutescente; ascis cylindraceis; sporidiis subcylindraceis, 6-8 μ . longis; paraphysibus asco longioribus, sursum attenuatis.

Hab. in culmis *Calamagrostidis canadensis*, West Albany. — Paraphysibus sursum attenuatis *Das. hapalæ*, *D. brunneolæ* affinis.

86. **Trichopeziza distincta** (Peck) Sacc. *Pez. distincta* Peck 30 Rep. 1737

p. 60, t. I, f. 9-13. — Ascomatibus minutis $\frac{3}{4}$ -1 mm. diam., sessilibus vel puncto centrali affinis, extus nigrescentibus; margine tomentoso-piloso, pallide alutaceis vel olivaceis, disco udo rufo, sicco aurantiaco; ascis clavatis, sæpe tetrasporis, sporidiis oblongo-fusiformibus, rectis vel curvatis, 20-25 μ . longis.

Hab. in culmis *Andropogonis furcati*, Center Am. bor. — Margo in sicco inflexus et ascomata tum *Hysterium* simulantia.

87. **Trichopeziza nigro-cincta** (B. et C.) Sacc. *Peziza nigro-cincta* 1738

B. et C. North Am. Fung. n. 730. — Minuta punctiformis, plana coccinea, extus pilis brevibus, nigris hispidula; ascis clavatis; sporidiis oblongis curvatis.

Hab. in vaginis graminum in Carolina iuf. Americae borealis (RAVENEL).

88. **Trichopeziza fuegiana** (Speg.) Sacc. *Lachnella fuegiana* Speg. F. 1739

Fueg. n. 322. — Hinc inde laxæ gregaria superficialis sessilis vel vix subpedicellata, uda applanato-concava, 500-600 μ . diam., sicca globoso-contracta, candida, extus pilis biformibus intermixtis vestita, aliis longioribus, 200-300 = 4 deorsum cylindraceis non vel 1-2-septulatis sursum continuis attenuatis apice acutiuscule rotundatis medio densiuscule noduloso-asperulis, aliis brevioribus, 80-

100 \approx 5, cylindraceis 1-3-septatis apice obtusis non attenuatis (ubique levibus); disco albo-ceraceo; ascis cylindraceo-subclavatis apice late rotundatis non v. lenissime angulato-acutatis, deorsum breviter crasseque pedicellatis, 45-48 \approx 5-6, octosporis, pseudoparaphysibus tenuibus nonnihil longioribus, 50-60 \approx 3 apice incrassatulis et cuneato-acutissimis parce obvallatis; sporidiis recte v. oblique monostichis, cylindraceo-subfusoideis, utrinque attenuato-obtusatis, 8-10 \approx 1,5-2, inæquilateralibus, continuis, hyalinis.

Hab. ad folia emortua putrescentia Poæ Forsteri in scopulosis prope Port Cook in Staten island Fuegiæ. — Obturaculus ascorum minutissimum intense jodi ope cœrulescit.

89. **Trichopeziza hystricula** (Karst.) Sacc. *Lachnum hystriculum* 1740 Karst. M. F. I, p. 182. — Sparsa vel subsparsa, sessilis, planiuscula, v. concaviuscula, pallida, pilis erectis, fuscis v. badiis, confertis strigosa, 0,1 mm. lat.; ascis cylindraceo-clavatis, 45-50 \approx 6; sporidiis distichis, fusoideo-elongatis, rectis v. leniter curvulis, guttulatis, 10-14 \approx 2-3; paraphysibus 2-3 μ . cr., apice acutiusculis.

Hab. in foliis Caricis acutæ prope Mustiala Fenniæ. — Pili distincte articulati, fusti, apice dilutiores et subinde leviter attenuati, 130-200 \approx 8. Sicca obconica.

90. **Trichopeziza amœna** (Boud.) Sacc. *Coronella amœna* Boud. Bull. 1741 Soc. Myc. IV, p. 82, t. 17, f. 4. — Minutissima, pulchella, sparsa, 0,2-0,3 mm. lata, brevissime hirta, atrata, disco et margine late albo-cœrulecentibus, cupula hemisphærica initio turbinata, pediculo brevissimo punctiformi, extus ochracea, pilis brevissimis obtusis, continuis, nigro-olivaceis, 30-40 \approx 5-7, conspersa, margine pilis hyalinis, erectis, parallelis, subpulverulentis, continuis, longioribus, coronam latam, albo-cyanescentem formantibus; hymenio etiam albido-cœrulecente, primitus vix urceolato, denique plano aut paullulum convexo, marginem vix superante; paraphysibus crassioribus, septatis, ad apicem incrassatis et intus granulis oleosis repletis, 5-7 μ . cr.; ascis sat amplis, octosporis, clavatis, 120-130 \approx 12-13; sporidiis oblongo-fusiformibus, rectis aut vix curvulis, subclavatis, intus sporidiolis duobus et granulis oleosis donatis, 17-20 \approx 4-5.

Hab. ad folia Graminum et Caricum humi strata, Montmorency Galliæ.

91. **Trichopeziza Stipæ** Fuck. Symb. mycol. p. 297. — Cupulis gregariis sparsisve, punctiformibus, sessilibus, primo clausis, globosis, demum apertis, subturbinatis, extus rugulosis, atro-fuscis, gla-

bris, margine pallidiore, connivente, disco pallide fusco; ascis cylindraceis, sessilibus, 8-sporis, $34 \approx 6$; sporidiis subdistichis, oblongis, utrinque obtusis, simplicibus, $8 \approx 4$, hyalinis; paraphysibus simplicibus, asci longitudine.

Hab. in foliis siccis adhuc stantibus *Stipæ capillatæ* in silvis Budenheim Rhenogoviæ.

92. **Trichopeziza badiella** (Karst.) Sacc. *Lachnum badiellum* Karst. 1743

M. F. I, p. 182. — Sparsa, sessilis, subobconica, badio-strigoso-hirtella, 0,2 mm. lat.; ascis cylindraceo-clavatis, 36-44 $\approx 5-6$; sporidiis fusoideo-elongatis, rectis, guttulatis, $8-10 \approx 2$; paraphysibus gracilescensibus, 2-3 μ . cr., supra acutiusculis.

Hab. ad calamos *Scirpi lacustris* putridos prope oppid. Wasa Fenniæ. — Pili fusi v. lutescenti-fusi, distincte articulati, $80-100 \approx 5-6$, apice saepissime attenuato.

93. **Trichopeziza fugiens** (Phill.) Sacc. *Lachnella fugiens* Phill. Disc. 1744

pag. 268. — Sparsa, sessilis, globosa dein expansa, tenuis, alba, villosa; ascis tereti-clavatis v. subfusoideis, 4-8-sporis, $20 \approx 5$; sporidiis tereti-oblongis v. ellipsoideis rectis curvulisve $7 \approx 2$.

Hab. in calamis *Junci* in paludibus prope Bristol Britanniæ (BUCKNALL). — Ascoma 50 μ . lat.; pili brevissimi continui hyalini.

6. Cryptogamicolæ.

94. **Trichopeziza Pteridis** (A. et S.) Rehm Ascom. n. 411, *Lachnella* 1745

Pteridis (A. S.) Phill. Disc. p. 256, t. VIII, f. 47, *P. Pteridis* A. S. p. 338, t. 12, f. 7. Pers. Myc. Eur. p. 300, Fr. Syst. Myc. II, p. 144, *Tric. pulveracea* Fuck. et Auct. p. p. — Sessilis, minima, ceraceo-mollis, cupularis, intus sordide lutescens, extus subgranulosa olivascens, margine crenato; ascis clavatis, $30 \approx 7$; sporidiis obtuse clavulatis, continuis, hyalinis, $8 \approx 2,5$, distichis; paraphysibus filiformibus.

Hab. ad stipites *Pteridis aquilinæ*, Osmundæ in Suecia, Britannia, Germania, Gallia, Italia, Belgio, Amer. bor. — Sparsa v. gregaria, punctiformis, 200-500 μ . lat., basi subapplanata, humida mollis aperta, sicca contrahitur, rugulosa, olivaceo-nigricans. Margo subinde lacinulatus. Excipulum sec. Rehm constat ex fibrillis, 4 μ . cr., septatis, rugulosis, obtusis supra liberatis et ascoma pilosulum reddentibus.

95. **Trichopeziza Osmundæ** (C. et E.) Sacc. *Peziza Osmundæ* C. et 1746

E. Grev. VII, p. 7. — Sparsa, minuta, nivea; cupulis demum ap-

planatis, margine dentatis, subtus leniter pubescentibus; ascis clavatis; sporidiis linearibus, $5 \mu.$ long.

Hab. in petiolis foliisque Osmundæ, New Jersey Amer. bor.

— Ascoma $\frac{1}{10}$ mm. lat.

96. **Trichopeziza Aspidii** (Lib.) Fuck. Symb. p. 297, *Peziza Aspidii* 1747 Lib. Exs. n. 226, *Lachnum Aspidii* (Lib.) Karst. Symb. Myc. Fenn. XXVI, pag. 27. — Ascomatibus sparsis, sessilibus, minutis, punctiformibus, globosis, velutinis, albescensibus; ascis fusoideo-oblongis, $34-40 \approx 6$; sporidiis bacillari-fusoideis, $8-10 \approx 1-1,5$; paraphysibus filiformibus.

Hab. in stipitibus et frondibus putridis Aspidii aculeati in Arduennis, Fennia, Belgio et Rhenogovia. — Pili circ. $50 \approx 3$.

97. **Trichopeziza hexagona** Fuck. Symb. myc. pag. 297. — Sparsa, 1748 subsessilis, punctiformis, alba; cupulis globosis, diaphanis, ore constricto, hexagono-radiato-ciliato, ciliis candidis; ascis stipitatis, elongatis, 8-sporis, $24 \approx 4$; sporidiis oblongis cylindraceisve, hyalinis, $4 \approx 2$.

Hab. in stipitibus putridis Aspidii Filicis-maris prope Cœstrich Rhenogoviæ.

98. **Trichopeziza inquilina** (Karst.) Sacc. *Helot. inquininum* Karst. 1749 M. F. I, p. 147, *Lachnella* Karst. Rev. p. 132. — Sparsa v. gregaria, sessilis v. substipitata, planiuscula, sicca hemisphærico- vel subsphæroideo-contracta, alba, tomentella, 0,6 mm. lat.; ascis cylindraceo-clavatis, $36-40 \approx 5-5,5$, obturaculo minutissimo obsolete v. vix jodo cœrulecente; sporidiis fusoideo-elongatis, guttulatis, $7-8 \approx 1-2,5$; paraphysibus $2 \mu.$ cr., sursum incrassatulis, apice attenuato.

Hab. ad stipites Equiseti silvatici vetustos in Mustiala Fenniae. — Pili cupulæ brevissimi, articulati, apice obtusi, $4-6 \mu.$ cr.

99. **Trichopeziza stereicola** (Cooke) Sacc. *Pez. stereicola* Cooke Grev. 1750 I, p. 130, *Calloria stereicola* Phill. Disc. p. 328. — Gregaria, minuta; ascomate globoso, dein cupuliformi, carnosò-ceraceo, extus villoso, sicco delicate roseo; disco concolori; ascis subclavatis; sporidiis minutis, cylindraceis, obtusis, $7 \approx 1,5$.

Hab. in hymenio Sterei, Scotland, Birmingham North Wales, — Ascoma 300-700 $\mu.$ lat.

100. **Trichopeziza Cœdema** (Desm.) Sacc. *Peziza Cœdema* Desm. XVIII 1751 Not. p. 4, *Lachnella* Phill. Disc. p. 271. — Hypophylla, sessilis, minutissima, subinde stipata, globosa, leniter pubescens, albo-grisea, subtremellosa; ascomatibus 1-1,5 mm. diam., subiculo brunneo

insidentibus; ascis minutis clavatis, sœpe curvis, octosporis; spori-
diis distichis, fusiformibus, 6-9=2; paraphysibus filiformibus.

Hab. parasitica in Phragmidio, in foliis emortuis Rubi in
Britannia et Gallia. — Ascoma $\frac{1}{7}$ mm. lat.

B. Species fructificatione ignota.

101. **Trichopeziza Veronicæ** (Lasch) Sacc. *Peziza Veronicæ* Lasch 1752
in Klotzsch H. M. n. 632 et in Bot. Zeit. 1845, p. 65. — Sessilis
e turbinato subglobosa, applanata, strigoso-villosa, fusco-cervina,
villis longissimis, superne albidis, interioribus albis v. lutescenti-
bus, disco albo.

Hab. in caulis Veronicæ pr. Driesen Germaniæ.

102. **Trichopeziza Scrophulariæ** (Lasch) Sacc. *Peziza Scrophulariæ* 1753
Lasch in Kl. Herb. myc. n. 1816 et in Flora 1854, pag. 203. —
Sessilis, minuta, subtremellosa, ex applanato leniter convexa, in-
cano-pallida flavescens, subtus puberula.

Hab. ad caules Scrophulariæ, Driesen Germaniæ (LASCH).

103. **Trichopeziza Aconiti** (Saut.) Sacc. *Peziza Aconiti* Saut. in Ra- 1754
benh. D. C. Fl. p. 344, n. 2637. — Erumpens, sparsa, punctifor-
mis, subclausa, primo quasi trochiformis, dein globoso-depressa,
brunneolo-flava, secedens, rugulosa, margine involuto.

Hab. in caulis emortuis Aconiti in agro salisburgensi Ger-
maniæ (SAUTER).

104. **Trichopeziza Sauteri** Sacc. *Pez. mollissima* Saut. Salzb. p. 8 1755
nec Lasch. — Sessilis, hemisphærico-applanata, villosa, tota nivea.

Hab. in caulis emortuis putrescentibus Urticæ prope Mit-
tersill. — *T. sulphureæ* propinqua, differt a colore albo margine-
que longiore.

105. **Trichopeziza plano-umbilicata** (Grev.) Sacc. *Peziza plano-um- 1756
bilicata* Grev. Fl. Ed. p. 420. — Minuta, sessilis, gregaria, alba,
globoso-concava, demum plana, margine pilis horizontalibus, albis,
ciliato, regulari, disco umbilicato.

Hab. in caulis Urticæ emortuis in Britannia.

106. **Trichopeziza adpressa** (Wallr.) Sacc. *Pez. adpressa* Wallr. Fl. 1757
crypt. n. 2424. — Ascomatibus primum subsessilibus rotundis ocius
jam in discum planum vitellino-lutescentem pallescentem reclusis
iisque adpressis planis lenticularibus majusculis, subtus parce to-
mentosis, margine pilis brevissimis rectis parcis prominulis pallide
sulphureis fimbriatis.

Hab. ad caules *Helianthi annui* siccis in agro Northusano Germaniæ (WALLR.).

107. *Trichopeziza Eupatorii* (Schw.) Sacc. *Peziza Eupatorii* Schw. 1758

Syn. pag. 174. — Gregaria, cupulis 4–6 mm. diam., madefactis explanatis, disco subconvexo, margine fere oblitterato, siccitate conivens, sed non clausa, rufo-carnea, extus pilis nigro-fuscis, apice albescientibus fasciculatim obsita.

Hab. versus radices in maximis caulis emortuis *Eupatorii* purpurei et *E. maculati*, Bethlehem Amer. bor.

108. *Trichopeziza Absinthii* (Lasch) Sacc. *Peziza Absinthii* Lasch 1759 in Klotzsch Herb. myc. n. 1317 (nec 1417.) et in Flora 1850, p. 282. — Ex flavido fuscescens, sessilis, subrotunda, ruguloso-fibrillosa, margine inflexo albo fimbriata, disco dilutior flavido.

Hab. in caulis *Artemisiæ Absinthii* in agro Driesensi Germaniæ (LASCH.).

109. *Trichopeziza Galegæ* (Ces.) Sacc. *Pez. Galegæ* Ces. in Klotzsch 1760 Herb. myc. n. 1614 et in Flora 1852, p. 395. — Gregaria, sessilis primitus clausa, depresso-globularis, amœne cæsia, dense flocculosa, serius urceolata nigrescens, disco pallido; fructificans explanata, irregularis, disco virente luteo v. rufescente.

Hab. ad caules *Galegæ Vercelli* Italiæ borealis (CESATI). — Valde insignis dicitur et pro ætate varians.

110. *Trichopeziza floccosa* (Lasch) Sacc. *Peziza floccosa* Lasch in 1761 Klolzsch Herb. Myc. n. 1225, Bot. Zeit. 1849, p. 293 nec Schw. — Sessilis, floccosa, membranacea, tenuis, extus sericeo-floccosa, nivea, disco incano.

Hab. in caulis emortuis *Spartii scoparii*, Driesen Germaniæ.

111. *Trichopeziza rufiberbis* (Schw.) Sacc. *Peziza rufiberbis* Schw. 1762

Syn. p. 173. — Applanata patellæformis, siccitate ob cupulæ tenerritatem irregulariter contracta, ceterum orbicularis et madefacta valde explanata rubella, carnosa, extus fasciculis pilorum brevium ruforum eximie barbata, 2–5 mm. diametro.

Hab. in stipitibus plurimis teneris dejectis (gregatim crescens), Bethlehem Amer. bor.

112. *Trichopeziza subglobosa* (Saut.) Sacc. *Peziza subglobosa* Sauter 1763 in Rabenh. D. C. Fl. p. 344, n. 2638. — Erumpens, socialis, flavo-brunnea v. rufescens, primo globosa, minutissima, dein aperta, scutellata, margine erecto, passim fisso, extrorsum e tomentoso pilosa.

Hab. in caulis putrescentibus herbarum in agro salisburgensi (SAUTER).

113. **Trichopeziza varians** (Saut.) Sacc. *Peziza varians* Sauter in 1764

Flora 1845, p. 133. — Sessilis, subcarnosa, 1-1,5 mm. diam., excavata, demum applanata, extus villosa, nivea, disco pallido.

Hab. in caulis uidis vetustis in agro Urbis Stiriae. — *Trichopezizæ adpressæ* Wallr. proxima.

114. **Trichopeziza virescens** (Schum.) Sacc. *Pez. virescens* Schum. 1765

Sael. p. 433, Fl. Dan. t. 1785, f. 2, Fr. Syst. Myc. II, p. 104. — Sessilis, hemisphærica, tomentosa, fusco-olivacea, disco albido-carneo.

Hab. in caulis herbarum majorum in Saellandia. — Minuta, sparsa, plana, margine tumido elevato, carne albida.

115. **Trichopeziza pellucens** (Ces.) Sacc. *Peziza pellucens* Ces. in 1766

Klotzsch Herb. Myc. n. 1620 et in Flora 1852, p. 396. — Microscopica, hyalina, extus puberula, cito explanata.

Hab. in caulis (?) in agro vercellensi Ital. bor. (CESATI).

116. **Trichopeziza occulta** (Ces.) Sacc. *Peziza* Ces. in Klotzsch Herb. 1767

myc. n. 1922, et in Flora 1855, p. 267. — Gregaria, membranacea, hinc in sicco contracta, cyathiformis, puncto adfixa, extus furfurella, e viridi luteola, ore flocculoso albicante.

Hab. ad caules (?) Vercelli Italiæ borealis (CESATI). — Minutissima sed denso grege facile conspicua.

117. **Trichopeziza syringæ** (Wallr.) Fuck. Symb. myc. App. III, 1768

pag. 31, *P. syringæ* Wallr. Fl. crypt. n. 2443. — Ascomatibus sessilibus sparsis carnosis, extus villo propexo rigidulo candido (asbestino) vestitis, intus fuscescentibus, primum (et in siccando) coniventibus globosis punctiformibus, dein discum pallidum recludentibus deplanatis ascosque pulverulentos pallide-stramineos eructantibus.

Hab. ad ramos Syringæ vulgaris, Lonicerae Caprifoliæ, in Thuringia et Rhenogovia. — Discus pallidus, demum ascis emersis et pulveris pallide-straminei instar profusis conspersus. An *Cyphella*?

118. **Trichopeziza velutina** (Wallr.) Sacc. *Pez. velutina* Wallr. Fl. 1769

crypt. n. 2565. — Ascomatibus sessilibus exiguis, extus fibris terris rigidulis, furfuraceis spadiceo-castaneis adspersis, primum subrotundo-globosis in osculum circinatum marginibus inflexis cinctum, dein in discum subconcolorem concavum reclusis patellaribus, siccis connivendo varieque contrahendo 2-3-angularibus.

Hab. ad ramulos coryleo in Thuringia (WALLR.).

149. **Trichopeziza venustula** (Desm.) Sacc. *Peziza* (Lachnea) *venustula* Desm. in Gill. Disc. pag. 80. — Ascomate minuto, sessili, superficiali, globuloso, dein applanato, pressione mutua compresso v. anguloso, tomento niveo-albo, udo aperto, disco albo.

Hab. gregaria in ramis et ramulis exsiccatis Aceris Negundinis in Gallia. — Affinis *T. punctiformi*.

120. **Trichopeziza Tiliæ** (Peck) Sacc. *Pez. Tiliæ* Peck 24 Rep. p. 96. 1771 — Gregaria, minuta, sessilis, concava, extus dense albo-pilosa, minuta, disco palide lutescente, saepe pilis inflexis celato.

Hab. in ramis *Tiliæ americanæ*, Knowersville Amer. bor. — Ab *L. tiliacea* distincta.

121. **Trichopeziza melaxantha** (Fr.) Sacc. *P. melaxantha* Fr. S. M. II, p. 97 nec p. 150, *P. melanotheja* Fr. S. M. II, p. 611 (index). — Sessilis, minuta, extus farinaceo-villosa helvolo-lutea, disco concaviusculo nigro.

Hab. ad ramos prostratos decorticatos Fagi, in Suecia, Britannia et Carolina inf. — Inter minimas, sed pulchra. Gregaria, sicca, granuliformis, disco demum plano opaco, margine parum prominente integro.

122. **Trichopeziza involuta** (Lagg.) Sacc. *Peziza involuta* Lagger in Flora 1836, I, p. 236. — Sessilis, gregaria vix 1 mm. diam., obscure olivacea, intus albida vel subcæspitosa, cupulis subglobosis, pilis longis basi insertis strigosis.

Hab. ad ramulos fagineos et quercinos in Helvetia.

123. **Trichopeziza roseo-alba** (Schw.) Sacc. *P. roseo-alba* Schwein. 1774 Carol. n. 1244, Fr. Syst. Myc. II, pag. 114. — Sessilis, ceracea, subglobosa, rubra, margine crenato, extus albo-villosa.

Hab. frequens in cortice Corni floridæ in Carolina Amer. bor. — Minuta, sparsa. Cupula primum fere globosa, demum dilatata et difformis, 2-5 mm. lata.

124. **Trichopeziza cruenta** (Wallr.) Sacc. *Pez. cruenta* Wallr. Fl. crypt. n. 2438. — Ascomatibus sessilibus mollibus, extus pilis villosis tortuosis laxe adpressis vestitis ferrugineo-cruentis, primum clausis dein discum albo-pallidum cum compage interna concolor rem recludentibus.

Hab. ad ramulos siccios Corni masculæ agri badensis (KNEIFF), thuringiensis (WALLR.). — Habitu *L. flammeam* emulatur, villi colore ferrugineo-cruento et disci compagisque internæ pallore affatim differt.

125. **Trichopeziza Opulifoliæ** (Schw.) Sacc. *Peziza Opulifoliæ* Schw. 1776
 Syn. pag. 175. — Sessilis, sparsa, e globoso-turbinata, regularis, marginè tumido infexo, brunnea, extus fibroso-striata, marginè albescenti, disco pallido, diametro 2 mm.
Hab. frequens in ramis emortuis Spirææ opulifoliæ, Bethlehem Amer. bor.
126. **Trichopeziza penicillata** (Schw.) Sacc. *P. penicillata* Schwein. 1777
 Carol. p. 94, Fr. Syst. Myc. II, pag. 102. — Sessilis, perexigua, concava, alba, extus pilis longis cinereis.
Hab. in rimis corticis Vitis in Carolina Amer. bor. — Punctiformis, madido tempore aperta, marginè crinito, siccitate contrahitur, ita ut pili longi cinerei penicillatim, byssi instar, ligno insidere videantur. Gregaria persistens.
127. **Trichopeziza pomicolor** (B. et Rav.) Sacc. *Peziza pomicolor* 1778
 B. et Rav. in Cooke Disc. Un. St. I, p. 294. — Sparsa, subhemisphærica, mox aperta, extus pomicolor furfuracea; disco olivaceo.
Hab. in cortice Taxodii in Car. inf. (RAVENEL).
128. **Trichopeziza radians** (Saut.) Sacc. *Peziza radians* Saut. Salzb. 1779
 p. 8. — Exigua sparsa, sessilis, alba, globosa, tomentosa, basi villis radiantibus cincta, disco poriformi.
Hab. in ramis Berberidis vulgaris prope Mittersill. — Ascoma $\frac{1}{2}$ μ. long. A *L. Berberidis* distincta.
129. **Trichopeziza tryblidioides** (Rabenh.) Sacc. *Peziza tryblidioides* 1780
 Rabenh. in Kl. Herb. myc. n. 1130 et in Flora 1848, p. 508, Str. in Flora 1850, p. 111. — Sessilis, globosa, applanata, villosa-to-mentosa, connivens, fusca, disco albida.
Hab. ad Berberidis ramulos in Stiria (SAUTER).
130. **Trichopeziza roseola** (Fr.) Sacc. *P. roseola* Fr. Syst. Myc. II, 1781
 p. 101. — Sessilis, concava, flexuosa, mollis, albido-puberula, disco incarnato-roseo.
Hab. ad cortices ramulorum dejectorum Juglandis etc. rarer in Suecia. — Substantia tenera, fragilis, quare in statu exsiccatum difformis et passim lacerata, sed margo non dentatus. Villus tenuissimus, demum evanescens.
131. **Trichopeziza Vitis** (Schw.) Sacc. *Pez. Vitis* Schw. Syn. p. 173. 1782
 — Minuta, difformiter globosa, clausa, carneus, sed villo albo extus laxius involuta, marginè crenulato-plicata; disco pallido punctiformi.
Hab. in rimis corticis Vitis, Bethlehem Amer. bor.
132. **Trichopeziza abnormis** Mont. Fl. J. Fern. n. 26 et Fl. Chil. VII, 1783

- 430 *Discomyceteæ, Pezizeæ, Hyalosporæ, Trichopeziza.*
p. 401. — *Sessilis, applanata, confluens, albida, subtus hirta; disco villosulo; ascis..... ignotis.*
Hab. ad cortices arborum in montibus, Juan Fernandez (BERTERO). — An *Hymenomycetis* species?
133. **Trichopeziza umbilicata** (B. et C.) Sacc. *Peziza umbilicata* B. 1784
et C. North Am. Fungi n. 734, *Peziza Curtissii* Cooke Disc. U.
S. II, p. 22. — *Depresse subglobosa; ore minuto umbilicata umbrino-gilva.*
Hab. in ramis decorticatis Rosarum, Pennsylvania (MICHERNER). — Ascoma 1–2 mm. lat. Ascii ignoti.
134. **Trichopeziza leonina** (Schw.) Sacc. *Peziza leonina* Schwein. 1785
Carol. p. 93, Fr. Syst. Myc. II, p. 101. — *Sessilis, concava, hirsuta, utrinque fulva, margine crenulato.*
Hab. in ligno Ulmi maxime putrido Carolinæ. — Cupulæ 2–3 mm. latæ, siccitate minutissimæ fiunt; sæpe tomento fulvo insident; pili sparsi, breves, colore magis rubro.
135. **Trichopeziza Gmelini** (Weinm.) Sacc. *Peziza Gmelini* Weinm. 1786
Hym. Gaster. p. 447. — *Sparsa confertaque, sessilis, exigua; ascomate carnosulo, pallide stramineo, extus tomento albo, brevissimo obducto, margine elevato integro.*
Hab. ad ligna decorticata Alni incanæ in Rossia. — Ascoma primo exacte globosum, apertum, album, mox pallide stramineum, dein discus appланatus, dilutissime cinerescens.
136. **Trichopeziza triformis** (Fr.) Sacc. *Peziza triformis* Fr. in Fl. 1787
Dan. t. 1620, fig. sup. et S. M. II, p. 101. — *Sessilis, primo granuliformis, villosa alba, dein aperta, disco concavo fulvo, tandem convexo aurantiaco marginem obliterate.*
Hab. ad ligna putrida in Dania. — Inter minimas, gregaria, non satis nota.
137. **Trichopeziza inconspicua** (B. et C.) Sacc. *Peziza inconspicua* 1788
B. et C. North Pac. Exp. n. 153. — *Subsessilis; cupula hemisphærica, margine incurvo, extus albo-pruinoso, initio luteolo.*
Hab. in ligno putri, Bonin Islands.
138. **Trichopeziza gracillima** (Weinm.) Sacc. *Pez. gracillima* Weinm. 1789
Hym. Gaster. p. 445. — *Gregaria, minuta, subsessilis; ascomate globoso-turbinato, aquose albido, extus pilis tenerrimis vestito.*
Hab. in lignis carie consumptis post largas pluvias æstivales in Rossia. — Ascoma regulare, excavatum, margine semper involuto. — Stipes fere papillæformis.
139. **Trichopeziza fulvo-cana** (Schw.) Sacc. *Peziza fulvo-cana* Schw. 1790

Carol. p. 94, Fr. Syst. Myc. II, p. 101. — Semiimmersa, conicoglobosa, tomentosa, luteo-fulva, ore subconnivente incano.

Hab. rarer, sed distinctissima in frustulis succosis in arborum cæsarum disco in Carolina Amer. bor. — 2 mm. lata, intus incana; tomento intense piloso extus tecta, pilis crassioribus in lignum irrepentibus.

140. **Trichopeziza Martini** (Pers.) Sacc. *Peziza Martini* Pers. Myc. 1791

Eur. p. 251, Fr. Syst. Myc. II, p. 100. — Sessilis, depressa, orbicularis, cervina, disco applanato lutescente.

Hab. ad ligna prope Taurinum Ital. bor. — Gregaria, majuscula, 4 mm. fere lata. An a *T. variicolore* vere diversa?

141. **Trichopeziza comata** (Schw.) Sacc. *Peziza comata* Schw. Syn. 1792

p. 173. — Minutissima, sessilis, candidissima, extus pilis longis densis candidis eleganter comata, sparsa.

Hab. epiphylla in foliis quercinis dejectis, Bethlehem Americæ borealis.

142. **Trichopeziza jucundissima** (Desm.) Sacc. *Peziza jucundissima* 1793

Desm. 14 Not. p. 186. — Amphigena, sparsa, sessilis, minuta, utrinque nivea, membranacea, primo infundibuliformis, tandem plana friabilis, extus pilis longissimis, rigidulis vestita, margine flexuoso.

Hab. ad folia dejecta populina in Gallia bor. — Ascoma 1 mm. lat. Asci ignoti. An *Cyphella*?

143. **Trichopeziza Platani** (Pers.) Sacc. *Peziza Platani* Pers. Obs. 1794

2, p. 87, Fr. Syst. Myc. II, p. 103. — Sessilis, pusilla, orbicularis, setosa, nigra, disco plano albido.

Hab. rarissime in foliis deciduis Aceris Platanoidis in Germania. — Cupula sicca connivet, humida folio adpressa; setæ strictæ marginem præprimis occupant.

144. **Trichopeziza Piceæ** (Pers.) Sacc. *Peziza Piceæ* Pers. Myc. Eur. 1795

p. 269, Fr. Syst. Myc. II, p. 97, *Cenangium regulare* Fr. Elench. II, p. 22. — Sessilis, minuta, suburceolata, pulveraceo-pubescent grisea, disco piano nigrescente.

Hab. in Pini Piceæ foliis in montibus Juræ (CHAILLET). — Sparsa, dilute cinerea.

145. **Trichopeziza Arundinis** (Fr.) Sacc. *Peziza Arundinis* Fr. Syst. 1796

Myc. II, p. 105. — Sessilis, per exigua, punctiformis, fuscescens, subferrugineo-pilosa, disco fuligineo-pallente.

Hab. in culmis exsiccatis Phragmitis, Arundinis in Germania (KUNZE). — Gregaria, tenuis, non innata; primo sistit puncta minuta globosa, dein aperta cupulata, demum pili secedunt;

432 Discomyceteæ, Pezizeæ, Hyalosporæ, Dasyscypha.

color in siccis totus fucus, sed humectatus expallens, præcipue discus.

146. **Trichopeziza cyathiformis** (Wallr.) Sacc. *Peziza cyathiformis* 1791 Wallr. Fl. crypt. n. 2568. — Ascomatibus tenerime membranaceis concoloribus subsessilibus, sparsis, obconico-campanulatis, extus fibris furfuraceis stipatis vestitis, in discum subconcolorem patulum levem, marginibus inflexis cinctum reclusis.

Hab. ad gramina putria in tectorum grumis nec non inter Populi ligni fibras in Thuringia (WALLR.).

147. **Trichopeziza Struthiopteridis** (Saut.) Sacc. *Peziza Struthiopteridis* 1798 Saut. Pilz. Salzb. p. 8. — Perexigua, sessilis punctiformis, subvillosa; ascomatibus subglobosis, ore connivente, ciliato.

Hab. in stipitibus senescentibus Struthiopteris prope Mittersill. — *P. punctiformi* affinis.

148. **Trichopeziza episphæria** (Mart.) Lamb. Myc. Belg. II, p. 524, 1799 *Peziza episphæria* Mart. Erl. p. 465, Fr. Syst. Myc. II, p. 100. — Sessilis, plana, extus alba, pilis longis ciliato-pilosissima, disco dilute lutescente.

Hab. in Ustulina, Hypoxylo multiformi, Diatrypella verruciformi in Germania, Britannia, Belgio. — Minuta, gregaria, conferta.

149. **Trichopeziza pulverulenta** (Saut.) Sacc. *Peziza pulverulenta* 1800 Saut. Salz. p. 8. — Sessilis minuta hemisphærica, suburceolata, albo-pulverulenta, disco concaviusculo fulvo.

Hab. ad viam (Amtshausschlucht) prope Mittersill. — 1 mm. lata. *T. melaxanthæ* affinis.

DASYSCYPHA Fr. Syst. Myc. II, p. 89 (ut sect.) [1822] emend. Fuck. Symb. mycol. p. 304 (Etym. *dasy* pilosus et *scyphos* poculum), *Lachnella* Sacc. e. p. — Ascoma ceraceum, distincte stipitatum, ex urceolato scutellatum, extus pilosum v. margine distinctius ciliatum nunc minutum tenacellum, nunc minutissimum tenerius (*Hyalopeziza*, si mavis). Asci tereti-fusoidei, 4-8-spori. Sporidia tereti-oblonga vel fusoidea, continua, hyalina. Paraphyses saepius bacillari-fusoideæ, crassiusculæ, apice acutæ, plerumque ascos superantes.

A. Pili ascomatis hyalini vel pallidi.

I. In truncis, lignis, ramis.

1. **Dasyscypha virginea** (Batsch) Fuck. Symb. p. 305, Sacc. F. it. 1801

fig. 1314 (f. *candida* et f. *longipes*) et fig. 1315 (f. *brevipes*), *Fungoides minimum* Mich. gen. p. 205, t. 36, f. 15, *Lachnum agaricinum* Retz Scand. p. 329, *Peziza parvula* Fl. Dan. t. 1016, f. 4, *P. virginea* Batsch Elench. p. 125, Holmsk. Ot. II, p. 31, t. 14, *P. nivea* Sowerb. fung. t. 65. — Stipitata, candida, cupula hemisphærica, extus pilis confertis patulis obsessa; ascis cylindraceis $42 \approx 4-5$, octosporis; sporidiis fusoideis, hyalinis, continuis, $6-7 \approx 2$ (Rehm), $10 \approx 1$ (Fuck.).

Hab. ad ligna, cortices, pericarpia, folia etc. in fagetis alibi-que in Suecia, Britannia, Fennia, Gallia, Belgio, Germania, Italia, Amer. bor., Fuegia, Tasmania. — Gregaria, 2 mm. alta discusque demum totus explanatus æque latus. Variat lutescens et flavus (Holmsk. loc. cit. f. dextr.). Stipes tenuis, æqualis, etiam subvillo-sus, raro oblitteratus. Pili longi, strigosi, in margine ciliares. — ***D. carpophila** Pers. et Fr. subturbinata minor, magis villosa. Iisdem fere locis et tempore. — ***D. testacea** Sacc. F. it. f. 131: cupulis quam in typo amplioribus, usque $1\frac{1}{4}$ mm. diam., disco sa-turate testaceo; ascis cylindraceis, $50 \approx 5$; sporidiis fusoideis vel clavulatis, $7-8 \approx 2$, eguttulatis hyalinis. In lignis cariosis, Padova.

2. **Dasyscypha chamæleontina** (Peck) Sacc. *Peziza chamæleontina* 1802
Peck 30 Rep. pag. 60. — Ascomatibus minutis, $160-375 \mu$. diam., primo globosis, dein expansis, disco plano, vix furfuraceo, albo, vix tacto flavescentia dein rubescens, stipite brevi; ascis clavatis, $25-35 \mu$. longis; sporidiis stipatis, oblongo-ovatis vel subclavatis, 5μ . long.

Hab. in ligno Abietis canadensis dejectis, Sandlake Amer. bor. — Pubescentia parcissima, hinc ad *Phialeam* abit.

3. **Dasyscypha pudibunda** (Quél.) Sacc. *Erinella pudibunda* Quél. 1803
XIV Suppl. p. 9, tab. XII, fig. 23. — Ascomatibus cyathiformibus, albo-niveis, dein roseis, subtiliter velutinis; pilis mollibus, albis tectis, tenuibus, flexuosis, 2-3 mm. diam., stipite tenui 1-1,5 mm. longo; hymenio demum roseo-amethysteo; sporidiis bacillaribus, 8μ . longis.

Hab. gregarie in ligno saligno in turfosis Morvan Juræ. — Affinis *D. virginea* et *D. nivea*.

4. **Dasyscypha lachnoderma** (Berk.) Rehm Ascom. n. 303, *Peziza lachnoderma* Berk. Fl. Tasm. p. 274. — Cupula subhemisphæri-ca breviter stipitata extus nivea tomentosa intus miniata; ascis longis clavatis; sporidiis fusoideo-filiformibus curvatis, $25-26 \mu$.

Hab. in cortice emortuo in Tasmania (ARCHER), Africa austral.

(MAC-OWAN) et Amer. bor. (ELLIS). — Habitus *D. calycinæ* sed sporidia omnino diversa. Ascoma 4 mm. lat. Sec. Rehm asci 60 \approx 7; sporidia 18 \approx 3; paraphyses filiformes asco longiores; pili simplices, asperi, 6 μ . cr. luteoli, apice hyalini.

5. ***Dasyscypha distinguenda*** (Karst.) Sacc. *Lachnella* Karst. Rev. 1805 monogr. p. 131, *Helotium* Karst. M. F. I, p. 156. — Gregaria vel subsparsa, stipitata, alba, villosula; cupula planiuscula, sicca hemisphærico-contracta, disco albo, sicco sublutescente, 1-1,5 mm. lat.; stipite firmulo, æquali vel saepius sursum incrassato, brevi, 0,4-0,6 mm. alt.; ascis subclavato-cylindraceis 60-70 \approx 4,5-5,5; sporidiis fusoideo-oblongatis, rectis, eguttulatis, 6-12 \approx 2-3; paraphysibus filiformibus, 1,5 μ . cr.

Hab. supra lignum vetustum pineum in Mustiala Fenniæ.

6. ***Dasyscypha latebrosa*** (Ellis) Sacc. *Lachnella latebrosa* Ellis Torr. 1806 Bot. Cl. 1882, pag. 18. — Stipitata, luteo-alba, 400 μ . alta, pilis glandularibus, brevibus tecta; disco concavo, 250 μ . lat., obscure albescente vel pallide roseo, margine incurvato, pilis rectis, rugosis, sursum obscure septatis, apice obtusis vel leniter inflatis, 50 μ . longis ciliato; ascis cylindraceo-clavatis, 25-35 \approx 3,5; paraphysibus linearibus, sursum non acutis; sporidiis distichis, simplicibus, hyalinis, 3-5,5 μ . longis.

Hab. ad ligna pinea fabrefacta mucida in cellis in America boreali. — *D. virginea* similis, sed pilis non capitatis et paraphysibus apice non acutis sporidiisque minoribus præcipue distincta.

7. ***Dasyscypha papyracea*** (Karst.) Sacc. *Lachnum papyraceum* 1807 Karst. M. F. I, p. 169. — Gregaria, stipitata, alba vel albida; cupula tenui, planiuscula, sicca concaviuscula, tomentella vel villosula, disco lutescente, 1-3 mm. lat.; stipite tenui æquali, pilis patulis obpresso, 0,7-1 mm. alt.; ascis subclavato-cylindraceis, 35-50 \approx 3-4; sporidiis fusoideo-elongatis, rectis, 3-7 \approx 1-1,5; paraphysibus 3-4 μ . cr., apice acuto.

Hab. supra aseres pineos putrescentes locis stercoratis in Mustiala Fenniæ. — *D. virginea* affinis, sed cupulis majusculis, tomentellis sporidiisque minutis diversa; pili cupulæ 3-4 μ . cr., apice subclavato-obtuso.

8. ***Dasyscypha globuligera*** Fuck. Symb. myc. App. II, p. 60. — Cupulis demum aut in globulorum conidiophorum singulorum latere, aut in globulorum confluentium centro ortis, stipitatis, stipitibus 1-2 mm. altis, cupulis, primo subclausis, demum magis ex-

planatis, neconon concavis, 1-3 mm. latis, orbicularibus, integris, albis, extus stipiteque albo striato-hirtis, margine distincto, disco albo, demum lutescente; ascis cylindraceis, substipitatis, 8-sporis, $72 \div 8$; sporidiis oblique monostichis, clavato-fusciformibus, rectis, continuis, $10-12 \div 4$.

Hab. in ligno putri Salicis ad Rhenum pr. Estrich Rhenogoviae. — Fungus conidiophorus adest: globulis sessilibus, gelatinosis, mollibus, diaphanis, sordidis, $\frac{1}{2}$ -1 mm. diam., sparsis seu 2-4 confluentibus, intus filamentosis, hyphis ramosis conidiophoris; conidiis numerosis, cylindraceis, rectis, continuis, 8-10 \div 2.

9. **Dasyscypha decolorans** (Wallr.) Sacc. *P. decolorans* Wallr. Fl. 1809 crypt. n. 2578. — Ascomatibus gregariis stipato-carnosis, intus croceo-rubentibus, extus fibrillis teneris rigidulis adpressis, marginem versus seriatis albis vestitis, primum turbinatis, dein in discum pallide vitellinum, mox flavo-croceum incusum, margine inflexo discole re cinctum reclusis, hypocrateriformibus abrupte stipitatis.

Hab. ad ligna putrida Salicis capreæ in Thuringia (WALLR.).

10. **Dasyscypha dryina** (Karst.) Sacc. *Lachnella* Karst. Rev. mon. 1810 p. 131, *Helotium* Karst. M. F. I, p. 155. — Sparsa, stipitata, sublutea, albido-tomentella; cupula cyathoidea, sicca obovoideo-sphæroidea, ore connivente, circiter 1 mm. lat.; stipite firmulo, cupula breviore; ascis cylindraceo-subclavatis, $60-75 \div 4-5$, apice jodo non tinctis; sporidiis fusoideo-elongatis, rectis vel leviter curvulis, eguttulatis vel guttulas paucas minutas foventibus, $7-11 \div 1,5-2,5$; paraphysibus filiformibus, $1,5-2 \mu$. cr.

Hab. in ligno betulino in Fennia.

11. **Dasyscypha rorida** (Wallr.) Sacc. *Pez. rorida* Wallr. Fl. crypt. 1811 n. 2451, Fuck. Symb. p. 306. — Ascomatibus semiglobosis parvis albo-pallescensibus in discum concolorem v. lutescentem concavum reclusis, extus cum stipite brevi in basin tenuato pilis teneris apice crystallino-granulosis adspersis; ascis linearibus, substipitatis, 8-sporis, $28 \div 4$; sporidiis distichis, cylindraceis, subrectis, $6 \div 2$ hyalinis.

Hab. ad ligna sicca faginea in Thuringia et Rhenogovia. — Ascomata persistenter alba. *D. crystallinæ* affinis.

12. **Dasyscypha torva** (Wallr.) Sacc. *P. torva* Wallr. Fl. crypt. 1812 n. 2446. — Ascomatibus minutissimis, albis, primum e basi tenuata stipitiformi adpresso-pilosa penicilliformibus, dein subglobosis stipitem occultantibus, in discum urceolatum lutescentem reclusis, extus pilis porrectis rigidulis patulis torvis.

Hab. ad ligna Piri communis arida passim in Germania.

13. **Dasyscypha striatula** (Wallr.) Sacc. *Pez. striatula* Wallr. Fl. 1813
crypt. n. 2452. — Ascomatibus stipitatis, obconico-turbinatis, par-
vis albis dein stramineis, vertice depresso in osculum angustum
marginibus inflexis cinctum reclusis, extus cum stipite filiformi
striato firmo producto, pilis rigidulis patulis brevibus fragilibus ca-
ducis torvis.

Hab. ad ligna Cratægi Oxyacanthæ in Thuringia rarius.

— Habitus *Soleniæ anomala*, at stipite striato et pilis mollioribus,
laxioribus et fragilioribus diversa.

14. **Dasyscypha lactea** (E. et E.) Sacc. *Helotium lacteum* E. et E. 1814
in Journ. of Myc. 1888, p. 56. — Ascomatibus gregariis, stipita-
tis, tomentosis, albo-lacteis, plano-convexis, subimmarginatis, extus
minute tomentosis, 0,5 mm. crass., stipite cylindrico, æquali, brevi
170 μ . long. et cr., tomentoso; ascis tereti-clavatis, 75 \pm 6-7, fili-
formi-paraphysatis; sporidiis distichis, fusoideis, acutis, curvulis,
15-17 \pm 2,5-3, hyalinis.

Hab. in ligno putri, Cazenovia N. Y. Amer. bor.

15. **Dasyscypha soleniiformis** (E. et E.) Sacc. *Lachnella soleniifor-* 1815
mis E. et E. Journ. Myc. 1888 p. 55. — Gregaria, brunnea, breviter
stipitata, primo hemisphærica, cupuliformis et fere clausa, ob mar-
ginem incurvatum, dein expansa et vix plana, extus setis rectis,
albis, sursum rugosis, disco sordide albo, lutescente vel aquose
carneo, $\frac{1}{2}$ -1 mm. diam.; ascis circiter 40 \pm 3-3,5, clavato-cylin-
draceis, rectis; paraphysibus superantibus, obtuse lanceolatis; spo-
ridiis subdistichis, clavato-oblongis, hyalinis, continuis, 5-8 \pm 2,5
(subinde 10-12 μ . longis).

Hab. in ligno emortuo Cazenovia New York Amer. bor. —
Pili marginales 70-80 \pm 3, luteo-brunnei.

16. **Dasyscypha pygmæa** (Fr.) Sacc. *Peziza pygmæa* Fr. Syst. Myc. 1816
II, p. 79, *Helotium* Karst. M. F. I, p. 153, *Lachnella* Phill. Disc.
p. 242. — Sparsa vel subgregaria, stipitata, flocculoso-surfurella
lutescens, sicca subferruginea, cupula patelliformi, disco lateo sub-
inde demum convexiusculo, 1,5-3 mm. lat.; stipite subæquali, 1-7
mm. alt.; ascis cylindraceo-clavatis, 65-75 \pm 5-6; sporidiis mono-
stichis, fusoideo-elongatis, rectis, plerumque guttulatis, 5-10 \pm 1,5-2
(Karst.), 11 \pm 2,5 (Phill.); paraphysibus gracilibus.

Hab. ad truncos, ramos infossos, ad terram arenosam inter
muscos et radices terra tectas, locis umbrosis in Suecia, Fennia,
Britannia. — Forma **prolifera** Berk. Som. New. Fung. tab. 55,

- f. 7-13: cupulis subinde disco cupulas minores generantibus armeniacis, margine pallidiore. In Britannia.
17. **Dasyscypha calicioides** (DC.) Sacc. *Pez. calicioides* DC. Fr. 2, 1817 p. 81, Fr. Syst. Myc. II, p. 115 nec Rehm. — Cupula planiuscula, margine fimbriato, extus stipiteque gracili sursum dilatato rugoso-scabris.
Hab. ad cortices Galliæ. — Variat alba et sulphurea.
18. **Dasyscypha Crucibulum** (Fr.) Sacc. *P. Crucibulum* Fr. Syst. Myc. 1818 II, p. 112, Fl. Dan. t. 1784, f. 1? (nec Batsch), *P. crucibulifor-mis* Fr. II, p. 610. — Turbinata, albido-pallida, extus pilis confertis adpressis fibrosa, margine lato inflexo, stipite brevi firmo.
Hab. in trunco putrido fagineo in Suecia. — Sec. Rostrup icon Fl. Dan. citata potius *Cyathi* speciem sistit. Solitaria, major, 2 mm. alta, firmula subpiriformis, villo non soluto tecta, ad marginem in velum submembranaceo-contextum extenso. Stipes vix 2 mm. longus, rugosus.
19. **Dasyscypha viridula** (Schrad.) Sacc. *P. viridula* Schrad. Journ. 1819 p. 63, Fr. syst. II, p. 91. — Ascomatibus stipitatis cyathiformibus viridulis, extus albido-villosis.
Hab. ad truncos putres nemoris Sollingensis (SCHRADER).
20. **Dasyscypha inspersa** (B. et C.) Sacc. *Peziza inspersa* B. et C. 1820 Cub. Fung. n. 684. — Cupula subglobosa, extus stipiteque brevi crassiuscula dense albo-farinaceis; hymenio carneo.
Hab. in ligno emortuo in insula Cuba. — Minuta, gregaria.
21. **Dasyscypha nivea** (Hedw.) Sacc. *Pez. nivea* Fr. syst. II, 90, *P. 1821 virginea* Flor. dan. tab. 1440, fig. 1, *Octospora nivea* Hedw. fil. t. VIII, f. B, *Trichopeziza* Fuck. Sacc. F. ital. f. 1436, *Lachnella* Phill. Disc. p. 245, Pat. Tab. f. 169. — Ascomatibus breviter stipitatis candidis villoso-tomentosis, turbinatis, pilis tenuibus, hyalinis, septatis, obtusis; ascis cylindraceis; sporidiis fusoideis v. oblongo-fusoideis, distichis, 8-10 \times 2; paraphysibus filiformibus.
Hab. ad ramulos locis uidis dejectos et ligna putrida in Suecia, Britannia, Gallia, Belgio, Italia, Amer. bor. Nov. Zeland.
22. **Dasyscypha calycina** (Schum.) Fuck. Symb. p. 305, *Pez. calycina* 1822 Schum. Sael. p. 424, saltem p. p. nec Nyl. Karst. *P. calycina* var. *Laricis* Cooke Handb. pag. 685, *Peziza Willkomii* Hartig Lehrb. p. 118, t. XI, f. 10-19. — Gregaria v. sparsa, særissime cæspitosa, stipitata, albo-vilosula; cupula expansa, sicca concava, disco aurantiaco-luteo v. aurantiaco, concaviusculo vel planiusculo, 1-2 mm. lat.; stipite brevi crassiusculo, in cupulam dilatato; ascis sub-

cylindraceis, 120 \approx 9, octosporis; sporidiis elongatis, obtusis, continuis, monostichis, 16-22 \approx 6-7, hyalinis; paraphysibus filiformibus, hyalinis, sursum incrassatulis, pilis hyalinis, 2,5 μ . crassis.

Hab. in cortice Laricis et Pini silvestris in Germania, Britannia, Gallia, Italia. — Var. **Trevelyanii** Cooke Grev. III, p. 121: sporidiis majoribus, 30-35 μ . long., pseudoseptatis. In Britannia.

23. Dasyscypha subtilissima (Cooke) Sacc. *Peziza subtilissima* Cooke 1823

Grev. III, p. 121, f. 167, *Peziza calycina* Fr. ScL Suec. n. 360, Nyl. Pez. p. 24, e. p. nec Schum. Fuck. Rehm, *Helotium* Karst. M. F. I, pag. 154, *Erinella* Pat. Tab. 592, *Dasys. bruyeriensis* Sacc. Mich. II, p. 330, (sp. 7-1,5-2). — *Gregaria* v. *sparsa*, stipitata, initio globosa, dein hemisphærica, demum expansa, ceracea, extus albo-villosa; disco aurantio-flavo, margine integro; stipite brevi; ascis subclavatis; sporidiis cylindraceis, curvis, minutis, hyalinis, 10-12 \approx 2.

Hab. in cortice Abietis et Pini silvestris in Scotia, Sæcia, Fennia, Gallia et West Albany Amer. bor.

24. Dasyscypha Abietis (Karst.) Sacc. *Hel. Abietis* Karst. M. F. I, 1824

p. 154, *Peziza calycina* Auctt. pr. p. *Peziza calycina* β . *Abietis* Fr. Syst. Myc. l. c. saltem pr. p. — *D. subtilissimæ* sat similis sed ascomatibus magis villosis v. tomentosis; ascis clavatis, 70 \approx 8-10; sporidiis distichis, fusoideo-oblongatis v. fusoideo-elongatis, rectis v. curvulis, guttulis 2 majusculis præditis v. spurie tenuiter uniseptatis, 11-14 \approx 2,5-3,5.

Hab. ad ramulos Abietis excelsæ siccis in Fennia saltem meridionali autumno-vere sat frequens.

25. Dasyscypha Agassizii (B. et C.) Sacc. *Peziza Agassizii* B. et C. 1825

North Am. Fungi n. 713. — Fasciculata, erumpens; cupula extus tomentosa candida, intus aurantia, breviter stipitata; margine undulato; ascis subcylindraceis; sporidiis ellipso ideo-ovatis v. subfusiformibus, levibus, minutis, 6,5-7,5 \approx 4.

Hab. in corticibus Abietis Lake superior. (AGASSIZ), New England (OAKES) Amer. bor. — Subinde stipes longiusculus. Affinis *D. subtilissimæ*. — Var. **rufipes** Phill. Grev.: stipite rufescente. In Pino contorta, California.

26. Dasyscypha resinaria (C. et Phill.) Rehm Ascom. Lojk. n. 30, 1826

Lachnella C. et Phill. Disc. p. 242. — *Gregaria*, stipitata; cupula initio turbinata, dein aperta, $\frac{1}{2}$ -1 mm. lat., extus stipiteque brevi pilis albo-hyalinis scabris simplicibus, 4 μ . cr. obsessa; margine

inflexo; disco pallide flavo-aurantio; ascis cylindraceo-clavatis, 30-36 \approx 3-4; sporidiis ellipsoideis, 5 \approx 2,5; paraphysibus filiformibus.

Hab. in resina Abietis excelsæ, North Wales Britanniæ et in cortice abietino prope Teplicska Hungariæ. — Sec. Rehm sporidia subrotunda, nempe 2,5 \approx 2: an species diversa et ad *Lachnellulam* referenda?

27. **Dasyseypha bicolor** (Bull.) Fuck. Symb. mycol. p. 305, *Peziza* 1827 Bull. Champ. pag. 243, tab. 410, f. 3, *Lachnum* Karst. M. F. I, p. 172, *P. minuta* Fl. Dan. t. 779, fig. 2. — Sparsa v. conferta, stipitata v. subsessilis, albo-tomentosa v. dilutissime roseo-villosa, cupula humida planiuscula, sicca sphæroideo-contracta, disco aurantiaco-luteo v. aurantiaco v. subcinnabarino, 1-2 mm. lat.; stipite tenui, brevi vel indistincto; ascis cylindraceo-subclavatis, 45-60 \approx 4,5-5; sporidiis fusoideo-elongatis, 6-9 \approx 1,5-2; paraphysibus apice acutis, 3-4 μ . cr.

Hab. ad ramulos, rarius ligna arborum frondosarum, præsertim *Coryli*, *Quercus*, *Betulæ*, *Rubi*, *Myricæ*, *Crataegi* in Suecia, Britannia, Gallia, Italia, Belgio, Germania, Amer. bor. — Cellulæ excipuli dilutissime fuscidulæ, ellipticæ, 8 \approx 5, pili cylindracei subechinulati, hinc illinc septato-constricti, 100-200 \approx 5. — Variat: a. disco coccineo aurantiaco, *P. Oxyacanthæ* Pers. Obs. I, p. 41, *P. pulchella* Myc. Eur. p. 260: b. disco luteo, *Pez. transparens* Pers. Obs. 2, p. 84, *P. quercina* Myc. Eur. p. 260: c. disco sordide pallido, Nees Syst. f. 204. — Var. **Rhododendri** Rehm Ascom. n. 110: cupulis subglobosis, dein breviter stipitatis, erumpentibus, dein sessilibus, pilosellis, extus hyalino-rosellis, disco rubicundo, 2-3 congregatis, 0,8 mm. diam., senilibus disco evanido persistentibus; pilis cupulæ brevibus, scabris, simplicibus, flavidulis, obtusis, septatis, 3,5 μ . diam.; ascis clavatis, 8-sporis, 50 \approx 7; sporidiis ellipticis, rectis v. subcurvatis, 1-cellularibus, primitus nucleis 2 parvulis præeditis, 8 \approx 2,5, distichis; paraphysibus sublanceolatis, ascos superantibus, apice obtusis, c. 3-4 μ . diam. Ad ramulos siccos Rhododendri ferruginei imprimis in alpibus editis tyrolensis. — Var. **alpina** Rehm Ascom. n. 568: sporidiis ellipsoideis obtusulis, rectis, 6-8 \approx 2,5, pilis ascomatis sublevibus; paraphysibus flavidis. In ramis Alni viridis. In Tyrolia.

28. **Dasyseypha translucida** (B. et C.) Sacc. *Peziza translucida* B. 1828 et C. North Am. Fungi n. 728. — Minuta, hyalina, gregaria; cupulis hemisphæricis margine inflexo; stipite brevissimo.

Hab. in ramis Castaneæ in Pensylvania (MICHENER).

29. **Dasyscypha crystallina** (Fuck.) Sacc. *P. crystallina* Fuck. Symb. 1829 p. 306. — Cupulis gregariis, pallide fuscis, in stipitem crassum, cupulam æquantem, sensim attenuatis, 2 mm. latis, primo clausis, demum apertis, margine infexo, extus stipiteque tenuissime candido-villosis, intermixtis multis pilis glanduliferis, firmioribus, glandulis hyalino-crystallinis, demum lucido-aureis, disco concavo, pallido; ascis linearibus, stipitatis, 8-sporis, 62 ± 4 ; sporidiis distichis in asci superiore parte, cylindraceis, rectis, hyalinis, 8 ± 2 .

Hab. in ramis putridis *Quercus* pr. Estrich Rhenogoviæ. — Habitus *Phialeæ cyathoideæ*, speciosissima. — Subiculum delicatissimum, ex hyphis ramosis, fuscis contextum, quandoque adest.

30. **Dasyscypha luteola** (Curr.) Sacc. *Helotium* Curr. Linn. Trans. 1830 XXIV, p. 153, t. 25, f. 11-12, 18, *Lachnella* Phill. Disc. p. 248. — Cæspitosa, pallide lutea; stipite circiter 4-5 mm. longo, disco 1-2 mm. lat., primo cavo, dein expanso et convexo, margine pilis parallelis, minutis, albis tecto et subsulcato; sporidiis rectis v. vix curvulis, hyalinis, $10-12 \pm 2$ longis; paraphysibus clavato-elongatis, acuminatis, subinde septatis.

Hab. in ramo *Ulicis* partim limo palustri immerso, Paul's Cray Britanniæ. — Habitum cæspitoso et subramoso distincta species.

31. **Dasyscypha Ulicis** (De Guern.) Sacc. *Pez. Ulicis* De Guern. in 1831 Cr. Finist. p. 52 et Rev. mycol. vol. VI, p. 99. — Ascomate stipitato 5 mm. usque 1 mm., intus amoëne luteo, extus pallidiore, albo-tomentoso, margine integro parum lobato, subconcavo vel convexo; ascis subcylindraceis, octosporis; sporidiis oblongis; paraphysibus subfusiformibus.

Hab. in ramis emortuis *Ulicis* et *Salicis* repentis Brest et Morlaix Galliæ. — Contextus excipuli e cellulis rotundis et pilis tenuissimis anastomosantibus formatus.

32. **Dasyscypha Comitissæ** (Cooke) Sacc. *P. Comitissæ* Cooke Grev. 1832 IV, p. 111. — Cæspitosa vel gregaria, nitide aureo-lutea, extus pallescente et tomentosa, stipitata, stipite communi, sursum ramoso; ascomate primo clavato, dein expanso et cupulato; ascis cylindraceis, $50-60 \mu$. long.; sporidiis minutis, fusoideo-linearibus rectis vel curvulis, $6-7,5 \pm 2,5$; paraphysibus filiformibus.

Hab. in ramis putridis Dupplin Castle, Perth Britanniæ. — Afinis, stipite sursum ramoso, *Dasys. pygmaea*.

33. **Dasyscypha crucifera** (Phill.) Sacc. *Peziza crucifera* Phill. Gard. 1833 Chron. 1878, p. 397, f. 71. — Minuta, gregaria, stipitata, alba; ascomatibus primo globosis, dein expansis, setulis hyalinis, septa-

tis, brevibus, apice clavatis, crystallis cruciformibus coronatis tectis; stipite longo, basi villoso; ascis cylindraceo-clavatis octosporis; sporidiis cylindraceis vel stricte fusiformibus, rectis, 6-8 \approx 1; paraphysibus asco longioribus sursum acuminatis.

Hab. in ramulis emortuis Myricæ Gales, Capel Curig, North Wales, Norfolk et Scotia. — Totus fungus 5 mm. alt.

34. **Dasyscypha lætior** (Karst.) Sacc. *Lachnum lætius* Karst. M. F. 1834

I, p. 174. — *D. acutipilæ* affinis; cupulis sessilibus, paullo majoribus, 0,7-1 mm. lat., siccis difformiter contractis, disco dilute luteo, sicco luteo; sporidiis nonnihil majoribus 10-20 \approx 1-2.

Hab. in sarmenitis Rubi idæi emortuis in Mustiala Fenniæ.

35. **Dasyscypha albipes** (Wallr.) Sacc. *P. albipes* Wallr. Fr. Crypt. 1835

n. 2574. — Ascomatibus minutissimis, e basi stipitiformi crassa brevissima glabra alba hemisphæricis, osculo angusto circinato hiantibus, pallide griseis, extus scobe farinosa cum fibrillulis intermixtis vestitis.

Hab. ad ramulos decorticatos Ligustri vulgaris in agro northusano Germaniæ, rarius (WALLR.).

II. In Dicotyledoneis herbaceis.

36. **Dasyscypha gracilescens** (Karst.) Sacc. *Lachnum gracilescens* 1836

Karst. M. F. I, p. 171. — Sparsa vel subgregaria, stipitata, alba, pilosula; cupula hemisphærica vel humida planiuscula, sicca subsphæroideo-contracta, pilis sat sparsis patulis brevibus obsita, circa 0,6 mm. lat.; stipite tenui, æquali, 0,6 mm. alt.; ascis subclavato-cylindraceis, 45-50 \approx 4; sporidiis fusoideo-filiformibus, vulgo rectis 10-16 \approx 1-1,5; paraphysibus superne acutiusculis, 3 μ . cr.

Hab. supra caules herbarum putrescentes in Mustiala Fenniæ.

— Pili cupulæ 4 μ . cr., apice subclavato-obtuso; discus in statu sicco saepè lutescens; sporidiis majoribus a *D. virginea* mox dignoscitur. — Var. β . **longicaudata** Karst. loc. cit. A *D. gracilescens* recedit stipite longiore (1-2 mm. alt.), sporidiis 6-14 \approx 1-1,5 et paraphysibus crassioribus (5-6 μ . cr.). Ad caules Ranunculi acris putrescentes prope Mustiala Fenniæ.

37. **Dasyscypha aurorina** (Quél.) Sacc. *Erinella aurorina* Quél. Esp. 1837

nouv. Champ. Fr. 1878, p. 291 et X Suppl. p. 12, tab. IX, f. 2.

— Ascomatibus planis, 1 mm. diam., puberulis incarnatisque, stipite piliformi pubescente, concolori, 3-4 mm. longo; hymenio roseo-aurantiaco; sporidiis fusiformibus, acicularibus, 8 μ . longis.

Hab. in frustulis herbaceis putrescentibus, Jura.

38. **Dasyscypha nudipes** (Fuck.) Sacc. *Pez. nudipes* Fuck. Symb. 1838 mycol. p. 306, *Helotium* Sacc. Mich. II, p. 612. — Gregaria, 3 mm. alta; stipite tenui, gracili, sordido cupulaque dense hyalino-puberulo, basi nudo, obscuriore; cupulis late turbinatis, ex cavatis, marginatis, disco flavescente, concavo, siccis margine connivente; ascis linear-clavatis; sporidiis 8, distichis, cylindraceis, hyalinis $10 \approx 2$.

Hab. ad caules sicclos Spirææ Ulmariae, rarius, pr. Estrich Rhenogoviae et in Helvetia. — Affinis *Ph. crystallinæ*, statura *Ph. cyathoideæ*.

39. **Dasyscypha spiræicola** (Karst.) Sacc. *Lachnum spiræicola* Karst. 1839 M. F. I, pag. 170. — *Das. virginea* sat similis, sed cupula alba, villosula, disco lutescente vel albido, demum saepe pallido, siccо subluteo vel subaurantiaco-luteo, 0,5-1,5 mm. lat., stipite tenui, 0,3-1 mm. alt.; ascis 48-55 \approx 4; sporidiis 6-12 \approx 1-1,5; paraphysibus 4-5 μ . cr.

Hab. ad caules Spirææ Ulmariae putrescentes in Fennia saltem australi frequentissima.

40. **Dasyscypha labiata** (R. et D.) Sacc. *Peziza labiata* Rob. et Desm. 1840 14 Not. p. 184. — Sparsa vel vix gregaria, minima, stipitata, junior globosa, adulta plana, siccо plicato-labiata, extus fufuracea, grisea, substricta disco helvolo, margine integro, stipite brevi glabro crassiusculo; ascis subclavatis, 50 μ . longis; sporidiis oblongis, 10 μ . long.

Hab. in foliis aridis, pedunculis, paleis etc. variarum plantarum, praesertim Eryngii campestris, Carlinæ in Gallia. — Stipes $\frac{1}{2}$ mm. alt., ascoma $\frac{2}{3}$ - $\frac{3}{4}$ mm. lat.

41. **Dasyscypha montana** (Quél. et Pat.) Sacc. *Erinella montana* 1841 Quél. et Pat. Tab. An. II, p. 35, fig. 589. — Ascomate turbinato, stipitato, extus albescente et lanoso, 5-7 mm. diam.; disco carneo-ochraceo; ascis elongatis; paraphysibus linearibus obtusis, apice parum incrassatis; sporidiis ovoideis, 7-10 \approx 3, hyalinis.

Hab. gregarie in capitulis Cirsii, Montée de la Dole Jura Galliæ.

42. **Dasyscypha vernalis** (Schum.) Sacc. *Pez. vernalis* Schum. Saell. 1842 p. 425, Fr. S. M. II, pag. 89. — Firmula, cinereo-pallida, cupula hemisphærica extus stipiteque elongato filiformi flexuoso villosis.

Hab. in stipitibus dejectis in Saellandia. — Majuscula in hac tribu, stipite 18-20 mm. long., cupula 5 mm. fere lata. Margo primo involutus, demum patulo-erectus.

III. Foliicolæ.

43. **Dasyscypha ciliaris** (Schrad.) Sacc. *Peziza* Schrad. Journ. 1799, 1843

II, p. 63, Fr. Syst. Myc. II, p. 89, Pat. Tab. 88 (*patula*), *Hyalopeziza* Fuck. Symb. p. 298, *Trichopeziza* Rehm Ascom. n. 258. — Stipitata, cyathiformis, nivea, extrinsecus pilis sparsis longis capitatis obsessa, disco albido; ascis clavato-oblongis, oblique stipitatis, 8-sporis, 50-56 = 6-8; sporidiis subdistichis vel oblique monostichis, fusiformibus, rectis vel curvatis, hyalinis, 18 = 2,5-3; paraphysibus filiformibus acutis, 2 μ . cr.

Hab. supra folia dejecta quercina, faginea et Myricæ Gales in Suecia, Gallia, Belgio, Italia, Austria, Britannia. — Facie convenit cum *P. virginea*, sed multo gracilior. Pili septulati, 80-100 = 7, capitulo crystalligero terminati. *Das. echinulatae* et *D. capitatae* et *patulae* peraffinis. — ***H. pulchella** Sacc. Mich. I, pag. 428, ascomatibus ut in typo sed disco amœne flavo; setulis asperulis, articulatis, 70-80 = 5-7; ascis cylindraceo-clavatis 60-70 = 7-8, paraphysatis, 8-sporis; sporidiis distichis, fusoideis, utrinque acutiusculis 24-27 = 2-3, 2-guttulatis, hyalinis. In foliis dejectis quercinis, Conegliano Ital. bor. (SPEGAZZINI).

44. **Dasyscypha patula** (Pers.) Sacc. *Peziza patula* Pers. syn. 654, 1844

Nees syst. f. 285, Fr. syst. II, 91, Pat. Tab. f. 87 (*ciliaris*), *Lachnella* Phill. Disc. pag. 151 *Hyalopeziza* Fuck. Symb. pag. 298, Sacc. F. ital. t. 1370. — Ascomatibus exiguis stipitatis extus dense villosis albis, primum hemisphæricis dein in discum planum lutescentem apertis, pilis longis tenuibus hyalinis, continuis, acutis, patulis; ascis cylindraceo-clavatis, 50-60 = 7, octosporis; sporidiis oblongo-clavulatis v. subfusoideis, 8-10 = 2-3, hyalinis; paraphysibus lanceolatis, ascis superantibus.

Hab. ad folia quercina, populina et betulina dejecta in Suecia, Germania, Britannia, Italia, Gallia, Belgio, Amer. bor. — Ascoma $\frac{2}{3}$ - $\frac{1}{2}$ mm. lat., brevissime stipitatum.

45. **Dasyscypha acuum** (A. et S.) Sacc. *Peziza* A. et S. Lus. p. 330, 1845

Helotium Fr. Summa p. 355, Karst. M. F. I, p. 147, *Lachnella* Phill. Disc. pag. 246. — Gregaria v. sparsa, stipitata, tenera, albida, cupula planiuscula, obsolete villosula v. pilosula, circiter 0,2 mm. lat., stipite tenui, plus minus elongato, subinde 1 mm. æquante; ascis cylindraceo-clavatis, 26-30 = 4-5; sporidiis oblique monostichis v. distichis, fusoideo-elongatis v. fusoideo-oblongatis, 4-6 = 1,5.

Hab. ad acus Pini silvestris, Abietis excelsæ et pe-

ctinatae locis umbrosis humidis in Fennia, Suecia, Germania, Britannia, in foliis Sequoiae in California. — Var. **carnea** Fuck. Symb. pag. 315: sporidiis angustioribus, teretibus, 4-5 μ . Cum typo.

46. **Dasyscypha serinella** (Quél.) Sacc. *Erinella serinella* Quél. XIV, 1846 Suppl. pag. 9, t. XII, f. 24. — Ascomatibus cyathiformibus, pilis granulosis ornatis, velutinis, nitide luteis, urceolatis, 500-700 μ . diam., margine luteo v. albo, stipite tenui, 600-800 μ . longo, hymenio pallide citrino, dein aureo; sporidiis acicularibus, fusiformibus, 10 μ . longis, guttulatis.

Hab. in acubus abietinis, Vosges Galliæ. — Affinis *D. patulæ*.

47. **Dasyscypha puberula** (Quél.) Sacc. *Erinella puberula* Quél. XIII 1847 Suppl. p. 9, t. VIII, fig. 21. — Ascomatibus cyathiformibus, 100-200 μ . diam., breviter stipitatis, ceraceis, luteis, velutinis, hyalino-ciliatis; hymenio plano, citrino; sporidiis fusiformibus, acicularibus, 5 μ . longis (?), guttulatis, hyalinis.

Hab. in foliis putrescentibus Pinii maritimæ, Saintonge Galliæ (P. BRUNAUD). — Videtur affinis *D. eurotioidi* et *D. Galbulæ*.

48. **Dasyscypha echinulata** (Rehm) Sacc. *Trichopeziza echinulata* 1848 Rehm Ascom. n. 259 (non Auersw.). — Stipitata, solitaria, sparsa, *D. ciliari* simillima, fere hyalina, parenchymatica, pilis hyalinis, fere ascomatis longitudinis, 3-4 μ . cr., simplicibus, apice obtuso crystallis oxalatis calcis conglomeratis obsesso, minutissimo; ascis clavatis, 8-sporis, 27 \approx 4, apice jodo cœrulecente; sporidiis ellipticis, hyalinis, 1-cellularibus, rectis, 3-4,5 \approx 1-1,5, distichis; paraphysibus apice acutatis, ascos multo superantibus, c. 5 μ . cr.

Hab. ad folia putrida Tiliæ in silva Harth prope Lipsiam (DR. G. WINTER).

49. **Dasyscypha virginella** (Cooke) Sacc. *Peziza virginella* Cooke 1849 Grev. IV, p. 178 (nomen), Ellis et Everh. N. A. Fungi n. 2144. — Ascomatibus sparsis, niveis tomentellis, longiuscule stipitatis; pilis hyalinis crystallino-capitatis, 40-50 μ . longis; ascis tereti-clavatis, 40 \approx 4-5, breve stipitatis; sporidiis fusoideis v. fusoideo-clavulatis, exiguis, 6-7 \approx 1 (an immaturis?); paraphysibus lanceolatis acutis.

Hab. in foliis dejectis Vaccinii, amphigena, Newfield Amer. bor. — Ascoma cum stipite $1/2$ mm. alt., cupula $1/3$ mm. lat. Diagnosis ex exempl. Ellisionis erui.

50. **Dasyscypha perforata** (Saut.) Sacc. *Peziza perforata* Saut. Pilz. 1850

Salzb. pag. 8. — Minima, stipitata, alba, cupula globosa, villosa, disco punctiformi luteo.

Hab. in foliis Alni putrescentibus, prope Stuhlfelden. — Inter *D. bicolorum* et *D. cerinam* media. Vix visibilis, stipite altitudinem ascomatis in longitudine æquante.

51. ***Dasyscypha hystricula*** (E. et E.) Sacc. *Lachnella hystricula* E. 1851 et E. Journ. Myc. 1888, p. 99. — Minuta, alba, sparsa, substipitata, sicca clausa, setis rectis, continuis v. obsolete septatis, rugosis, 50-75 \approx 4-5; ascis oblongo-clavatis, sessilibus, 25 \approx 7-8; paraphysibus non visis; sporidiis superne distichis et verisimiliter subfusoideis, (spurie?) 3-septatis, circiter 12 \approx 2 (immaturis visis).

Hab. in pagina inferiore foliorum Magnoliæ grandifloræ St. Martinsville Amer. bor. (LANGLOIS).

52. ***Dasyscypha albo-pileata*** (Cooke) Sacc. *Peziza albo-pileata* Cooke 1852 Hedw. 1875, p. 82, Grev. III, f. 182. — Epiphylla sparsa v. subgregaria, stipitata, candida dein ochracea, stipite gracili subnudo, cupula mox expansa extus floccis hyalinis, albidis, brevibus ornata, disco obscuriore; ascis cylindraceis; sporidiis linearibus minutis.

Hab. ad folia decidua Magnoliæ, New Jersey Amer. bor. (ELLIS). — Affinis *D. ciliari* sed major.

53. ***Dasyscypha australis*** (Speg.) Sacc. *Lachnella australis* Speg. 1853 F. Fung. n. 320. — Hypophylla, hinc inde (3-8) laxe gregaria, minutissima, uda turbinato-applanata, 400-500 μ . diam., sicca non v. vix contracta, margine involuta, deorsum attenuata ac in pedicellum glabrum, rigidulum concolorem, 500-800 \approx 50-80, desinens, extus sordide alba, densiuscule villosula, pilis breviusculis rectis vel flexuosulis cylindraceis, 60-80 \approx 3, apice non attenuatis obtuseque rotundatis 3-5-septatis saepèque ad septa constrictis dense minutaque papillulosis hyalinis vestita; disco pallescenti-ceraceo; ascis cylindraceo-subclavatis, antice attenuatis, vertice non v. vix subtruncatalis ac minute l-foveolatis, postice longe crasseque attenuato-pedicellatis, 45 \approx 2,5-3, octosporis, pseudoparaphysibus longioribus, 65-75 \approx 2,5, utrinque attenuatis, apice acutissimis, parce obvallatis; sporidiis acicularibus vel subclavulatis, utrinque attenuato-acutatis, 10 \approx 0,7-0,8, continuis, hyalinis.

Hab. ad folia dejicta putrescentia Berberidis ilicifoliæ in silvis in Staten island Fuegiæ.

54. ***Dasyscypha tautilla*** (Ph. et H.) Sacc. *Lachnella tautilla* Phill. 1854 et Harkn. Calif. p. 1. — Sparsa, stipitata, minuta, alba, cupuli-

formis, basi pilis gracilibus tecta; ascis late clavatis, octosporis; sporidiis ovoideis, $4 \approx 1$; paraphysibus gracilibus filiformibus.

Hab. in pagina inferiore foliorum *Garryæ ellipticæ* in California.

55. *Dasyscypha albo-citrina* (Cooke) Sacc. *Peziza albo-citrina* Cooke 1855

Grev. VII, pag. 47. — Stipitata, villosa, alba; cupulis turbinatis, demum expansis, $\frac{3}{4}$ mm., stipite brevi; disco citrino; ascis cylindraceis; sporidiis linearibus, rectis, 10μ . long.; paraphysibus acuminatis.

Hab. in foliis *Vaccinii*, Darien Georgia Amer. bor. — Per affinis *D. virginellæ*.

IV. In Monocotyledoneis.

56. *Dasyscypha imbecillis* (Karst.) Sacc. *Lachnum imbecille* Karst. 1856

M. F. I, p. 172. — Sparsa, stipitata, albo-tomentosa, cupula concava, disco luteo, 0,3-0,4 mm. lat., stipite latitudine cupulæ longiori, subæquali tenui; ascis cylindraceo-subclavatis; sporidiis distichis, fusoideo-bacillaribus, guttulatis, $10-16 \approx 1,5-2$; paraphysibus $5-6 \mu$. cr., apice acutis.

Hab. ad culmos et folia *Airæ cæspitosæ* et *Eriophori vaginati* putrescentes prope Mustiala Fenniæ.

57. *Dasyscypha elatior* (Karst.) Sacc. *Lachnum elatius* Karst. M. 1857

F. I, p. 171. — *D. virginea* sat similis sed major, cupula 1 mm. lat., disco dilutissime luteo stipiteque 1,5 mm. alt.; ascis 5μ . cr.; sporidiis $8-14 \approx 2-2,5$; paraphysibus 4μ . cr.

Hab. ad culmos graminum, terra tectos, in Mustiala Fenniæ.

58. *Dasyscypha caduca* (Rehm) Sacc. *Trichopeziza caduca* Rehm 1858

Ascom. n. 160, an *Lachnum acutipilum* Karsten myc. fenn. I, p. 173 cfr.: Cooke in Grevillea III, p. 128, tab. 41, f. 189. — Plerumque sparsa, stipitata, turbinata, albido-flavescens, sicca plerumque collapsa et clausa, humida rigida, stipite brevi, crasso, disco dilatato concavo, flavidо-rosello, piloso-hirta, 0,5 mm. diam.; pilis longis, acutis, vix scabris, simplicibus, hyalinis, c. $3-4 \mu$. crassis obsessa; ascis clavatis, 8-sporis, $50-60 \approx 6-8$, apice jodo cœrulecente; sporidiis fusoideo-ellipticis, rectis vel subcurvatis, interdum uno apice obtusis, 1-cellularibus, sœpe 2 nucleolis præditis, hyalinis, distichis, $10-15 \approx 2-2,5$; paraphysibus lanceolatis, ascis superantibus, basi 4μ . cr.

Hab. ad calamos sicclos Nardi strictæ in alpibus editioribus tyrolensisibus, juxta glaciem æternam.

59. **Dasyscypha carneola** Sacc. *Hyalopeziza* Sacc. Mich I, p. 253, 1859

F. ital. f. 1371. — Gregaria, brevissime stipitata, pallida, disco concaviusculo, carneo v. subaurantiaco, usque $\frac{1}{2}$ mm. diam., extus, præcipue margine, velutina, pilis apice subincrassatis, subhyalinis; ascis cylindraceis, 40 \approx 5, brevissime stipitatis, 8-sporis; paraphysibus (?) cylindraceis, sursum cuspidatis, 70-80 \approx 6 obvallatis; sporidiis minutissimis, cylindraceis v. clavulatis, rectis curvulisve, 5-6 \approx 1, hyalinis.

Hab. in foliis putrescentibus Moliniæ cœruleæ, Conegliano Ital. bor. (SPEGAZZINI). — Cum *D. deminuta*, *D. patenti*, *D. Carricis*, *D. rosea* pluribus notis convenit, tamen diversa videtur. —

***H rhodoleuca** Sacc. Mich. II, p. 79: ascomatibus subsessilibus concaviusculis, $\frac{1}{6}$ mm. d., extus pilis non septatis capitatis hyalinis, 50 \approx 5, capitulo 10 d., vestitis; disco amœne roseo; ascis cylindraceis subsessilibus, apice rotundatis 40-45 \approx 6; paraphysibus longioribus, utrinque cuspidatis, roseo-guttulatis, obvallatis, octosporis; sporidiis fusoideis, 10-11 \approx 1- $\frac{1}{2}$, hyalinis. In culmis graminum in Gallia (BRUNAUD). *Peziza graminum* B. et Br. forte hue spectat, sed, ignota fructificatione, dubia. A typo differt sporidiis duplo longioribus ceterisque.

60. **Dasyscypha acutipila** (Karst.) Sacc. *Lachnum acutipilum* Karst. 1860

M. F. I, p. 173. — Gregaria vel sparsa, stipitata, alba vel albida, villosa; cupula planiuscula vel convexiuscula, sicca concava, margine piloso-fimbriato, circiter 0,8 mm. lat.; stipite brevissimo, circiter 0,2 mm. alt.; ascis cylindraceo-clavatis, 40-48 \approx 4-5; sporidiis fusoideo-filiformibus, vulgo rectis, ut plurimum guttulatis, 8-16 \approx 1-1,5; paraphysibus 4-5 μ . cr., apice acutis.

Hab. ad caules graminum, præprimis Phragmitis communis, putrescentes in Fennia australi et media præsertim Wasa. — Pili cupulæ articulati, apice longe cuspidati, (basin versus) 4 μ . cr., discus interdum lutescens; cupula primitus sphæroidea, clausa.

61. **Dasyscypha controversa** (Cooke) Rehm Ascom. n. 114 Phill. Exs. 1861

n. 70, *Peziza* Cooke in Grev. IV, p. 41, t. 51, f. 264. — Breve stipitata, gregaria, cupula subhemisphærica; pilis filiformibus, obtusis, hyalinis, 4-5 μ . cr. obsessa; disco rubescente; ascis clavatis, 36 \approx 5, octosporis; sporidiis ellipsoideis, rectis, unicellularibus, 6 \approx 1,5; paraphysibus lanceolato-acuminatis, hyalinis.

Hab. ad culmos Phragmitis in Germania et Britannia. — Ascoma $\frac{1}{4}$ - $\frac{1}{3}$ mm. lat. Desideratur in Phillipsii Man. Brit. Disc.

— Var. **albescens** Rehm Ascom. n. 907: ascomatibus breve vel vix stipitatis albido-roseis, pilis flavidulis, septatis asperulis, $60 \approx 4-5$ obsessis; ascis $40-45 \approx 5$; sporidiis $6-9 \approx 1,5-2$. In culmis Phragmitis pr. Berlin (SYDOW).

62. **Dasyscypha Arundinariæ** (Berk.) Sacc. *Peziza Arundinariæ* B. 1862
North Am. Fungi n. 727, *Pez. cannea* Cooke Disc. U. S. I, p. 293.
— Parva, cupula cyathiformi extus albo-tomentosa; stipite demum glabratu discoque concavo pallide umbrinæ.

Hab. in culmis emortuis Arundinariæ in Carolina infer. Amer. bor.

63. **Dasyscypha pudica** (B. et R.) Sacc. *Peziza pudica* Bomm. et 1863
Rouss. Fl. Myc. Brux. pag. 126. — Ascomatibus stipitatis, albis, extus villosis, disco concavo albo, stipite villoso compactiusculo; ascis cylindraceis; paraphysisibus linearibus; sporidiis fusiformibus.

Hab. in foliis et radicibus Holci mollis in Belgio. — Fungillus fractus leviter cruentatus.

64. **Dasyscypha plagopus** (Wormsk.) Sacc. *Pez. plagopus* Wormsk. 1864
fl. Camtsch. ined. et in Fr. Syst. Myc. II, p. 93. — Pusilla, breviter stipitata, turbinato-infundibuliformis, extus villo sericeo adpresso, circa marginem piloso, disco luteo.

Hab. in foliis gramineis Kamtschatkæ. — Proximis minor, $\frac{1}{2}$ mm. alta et lata. Stipes brevis, cum cupula turbinata omnino confluens. Villus stramineus adpresso, sed ad marginem regularem subinflexum pili rigidi subflexuosi vel crispati, longius extantes. $\beta.$ stipite evidentiore, villo albo discum occultante. In ramulis dejectis Rubi triflori ibidem.

65. **Dasyscypha rufula** Quél. X Suppl. p. 12, tab. IX, fig. 4. — Ascomatibus cyathiformibus, 1 mm. diam., tenuibus, tenacibus, diaphanis, lanosis, carneis, ciliis serrulatis, albis præditis; stipite brevissimo, $100 \mu.$ longo; hymenio concavo, albo; sporidiis subcylindraceo-fusiformibus, $12 \mu.$ longis, 1-2-guttulatis, hyalinis.

Hab. in cæspitibus Graminacearum exsiccatis, Jura.

66. **Dasyscypha Eriophori** (Quél.) Sacc. *Erinella* Quél. X Suppl. 1866
(1880), p. 12, t. XI, f. 1. — Ascomatibus urceolatis, dein apertis, $500-700 \mu.$ diam., villosis, subtiliter denticulato-ciliatis, albis dein roseis; stipite brevi, basi attenuato, concolori, hymenio plano, pallide roseo-ochraceo; sporidiis fusoideis, acicularibus, $16 \mu.$ longis.

Hab. ad basim culmorum exsiccatorum Eriophori angustifolii, Normandia (A. LE BRETON). — Sec. Le Breton (Excurs. Crypt. p. 15), asci cylindracei, sessiles, apice rotundati jodoque

cœrularescentes, 50-70 \approx 6,5-8; sporidia 13-20 \approx 1,5-3; paraphyses ascos superantes lanceolato-auctæ.

67. **Dasyscypha hyalina** (Phill.) Sacc. *Peziza hyalina* Phill. Grev. 1867 IV, p. 51, f. 266. — Gregaria vel sparsa, breviter stipitata, minuta, pubescens, disco pallide lutescente, plano, marginato; ascis clavatis, octosporis; sporidiis fusiformibus, rectis vel vix curvis, 12-20 \approx 5-6; paraphysibus linearibus.

Hab. ad basim foliorum emortuorum Luzulæ silvaticæ, Conway, North Wales Britanniæ.

68. **Dasyscypha rosea** Rehm Ascom. n. 157. cfr. Cooke in Grevillea III, p. 128, tab. 41, f. 192. — Plerumque sparsa, interdum binata, brevissime stipitata, cyathoidea, pilosa, sicca plerumque complicata vel urceolata, albido-rosea, humida concava, urceolata, disco roseo, margine albido-fimbriato, c. 1 mm. diam., prosenchymerico-contexta, c. 0,5 mm. diam.; pilis brevibus, subscabris, septatis, apice subincrassatis et sœpe crystallis conglomeratis coronatis, hyalinis, 5-6 μ . cr. obsessa; ascis clavatis, 8-sporis, 45-50 \approx 6, apice jodo cœrularescente; sporidiis oblongo-ellipticis, plerumque nucleis 2 parvulis præditis, rectis obtusis, hyalinis, 1-cellularibus, distichis 20-24 \approx 1,5; paraphysibus lanceolatis, acutis, ascos superantibus, hyalinis, basi c. 4 μ .

Hab. ad calamos sicclos in alpibus tyrolensisibus juxta glaciem æternam.

69. **Dasyscypha luteo-disca** (Peck) Sacc. *Lachnella luteo-disca* Peck 1869 33 Rep. p. 31. — Ascomate minuto, 350-700 μ . diam., expansa, plana vel vix concava, marginata, extus furfuraceo-pubescens, alba, vel disco lutescente, siccо subaurantiaco; stipite brevi vel obsoleto; ascis subcylindraceis; sporidiis stipatis vel distichis, fusoides, 10-12 μ . longis.

Hab. in calamis emortuis Scirpi validi, Manlius Amer. bor.

70. **Dasyscypha deminuta** (Rob. et Desm.) Sacc. *Peziza deminuta* 1870 Rob. et Desm. 14 Not. p. 185, *Lachnella* Phill. Disc. pag. 253, *Lachnum* Rehm Asc. n. 707. — Minutissima, sparsa vel conferta, breviter stipitata $\frac{1}{2}$ mm. diam., extus albo-tomentosa, primo globosa, dein aperta hemisphærica; disco concavo, luteo, aureo vel aurantiaco; ascis tereti-clavatis; sporidiis oblongis 12 \approx 2 (Phill.); paraphysibus filiformibus.

Hab. in calamis siccis Junci in Gallia, Belgio, Germania, Britannia. — Ascomatis pili exteriores longi et tenues. Sec. Rehm loc.

cit. ascii 36-5; sporidia 5-6 \approx 1,5; paraphyses fusoideo-acutæ; hinc diversa species videtur.

71. **Dasyscypha curvipila** (Karst.) Sacc. *Helotium curvipilum* Karst. 1871

M. F. I, p. 162. — E sclerotio sicco rufo, foliis innato, gregatim enata, breviter stipitata, albida, cupula e sphæroideo hemisphærica, sicco sphæroideo-contracta, pilis flexuoso-curvatis, albidis dense vestita, 0,2 mm. lat., stipite brevi vel brevissimo, in cupulam abeunte; ascis cylindraceo-clavatis 32-36 \approx 4-5; sporidiis distichis, fusoideo-elongatis, vulgo rectis, guttulatis 7-8 \approx 5; paraphysisibus filiformibus.

Hab. ad folia *Caricis vesicariæ* emortua locis uidis prope Mustiala Fenniæ.

72. **Dasyscypha crinella** (E. et E.) Sacc. *Lachnella crinella* Ell. et 1872

Everh. in Bull. Torr. Cl. 1883, p. 76. — Sparsa, stipitata, pruinosa, alba, tenuis, delicata, disco 250-375 μ . diam., margine pilis brevibus ciliato; stipite longitudine circiter diametrum disci æquante; ascis oblongo-cylindraceis, 38-40 \approx 4-5, sessilibus; paraphysisibus non visis; sporidiis distichis, clavato-fusoideis, 9-12 \approx 2, hyalinis.

Hab. in foliis emortuis *Caricis crinitæ*, West Chester Am. bor. — Differt a *P. caricinella* ascis acutioribus, sporidiis simpli-cibus minoribus clavatis, et a *P. Caricis* sporidiis oblongo-cylindraceis.

73. **Dasyscypha Caricis** (Desm.) Sacc. *Peziza* Desm. in Ann. S. N. 1873

1840, p. 12. — Per exigua, sicca globulosa v. sabglobulosa, uda cupulata vel hemisphærica extus pilis albido-griseis minutis tomentosa, $\frac{1}{2}$ mm. diam.; disco planiusculo flavo-aureo; stipite gracili, pilosulo; ascis.....

Hab. in utraque pagina foliorum aridorum *Caricum* in Gallia.

74. **Dasyscypha caricicola** Sacc. *Hyalopeziza Caricis* (Desm.?) Sacc. 1874

F. Ven. Ser. IV, n. 82, F. ital. fig. 112. — Albo-lutescens, minuta, gregaria, mediocriter stipitata, undique pilosa margineque longius ciliata, pilis pallidis subcapitatis, disco concaviusculo (in sicco) nigricante-olivaceo; ascis cylindraceo-subclavatis basi vix attenuatis, 38 \approx 4-4 $\frac{1}{2}$; paraphysisibus nullis obvallatis, 8-sporis; sporidiis subdistichis, cylindraceis, 6 \approx 1, rectis curvulisve, hyalinis.

Hab. in pagina inferiore foliorum emortuorum *Caricis* silvaticæ in silva Montello, Treviso Ital. bor. — *Peziza Caricis* Desm. eadem ac mea esse posset, sed ejus fructificatio mihi ignota et cupulæ discus subaurantiacus.

75. **Dasyscypha callimorpha** (Karst.) Sacc. *Lachnum callimorphum* 1875

Karst. M. F. I, p. 173. — Gregaria, sessilis vel breviter stipitata, tomentosa; cupula planiuscula, sicca sphæroideo- vel hemisphærico-contracta, disco luteo vel aurantiaco-luteo, 0,3-0,5 mm. lat.; ascis subclavato-cylindraceis 35-45 \approx 5; sporidiis fusoideo-bacillaribus, guttulis 6-8 præditis vel spurie pluriseptatis, rectis, 17-20 \approx 1,5-2; paraphysibus subulatis, 3 μ . cr.

Hab. ad folia Caricis ericotorum exsiccata in Fennia saltem australi et ad folia Eriophori angustifolii (spor. 14-19 \approx 2-4) pr. Aberdeen Britanniæ (TRAIL). — Pili breviusculi, inarticulati, apice attenuati vel acuti, 3-4 μ . cr.

76. **Dasyscypha glabrescens** (C. et Ph.) Sacc. *Peziza glabrescens* 1876

C. et Phil. Grev. VIII, p. 62. — Sparsa, stipitata, candida; cupulis 0,5-1 mm. cyathiformibus, primitus subvillosis, demum nudis, glabris; ascis clavatis; sporidiis lanceolatis, utrinque rotundatis, guttulatis, 15-19 \approx 3-4; paraphysibus filiformibus.

Hab. in Rhipogono (?) Melbourne Australiæ.

77. **Dasyscypha pudicella** (Quél.) Sacc. *Erinella pudicella* Quél. XIII 1877

Suppl. p. 9, t. VIII, f. 25. — Ascomatibus cyathiformibus, velutino-gossypinis, pilis granulosis præditis, albo-niveis, dein roseo-maculatis 300-500 μ . diam.; stipite tenui, 300 μ . alto, hymenio albo, dein reseo; sporidiis acicularibus, 10-11 μ . longis.

Hab. gregaria in graminaceis minoribus Jura Galliæ.

V. In Acotyledoneis.

78. **Dasyscypha aspidiicola** (B. et Br.) Sacc. *Peziza* B. et Br. Ann. 1878

N. H. n. 771, *Helotium aspidiicolum* Rehm Asc. n. 560. — Ascomatibus minimis, cir. 0,2 mm. diam. et alt., cyathoideis vel globulo-so-stipitatis, gregariis vel sparsis, subscabris, albis, siccis alutaceis, extus floccoso-furfuraceis subhemisphæricis; ascis brevibus, gracilibus, clavatis, 20 \approx 6, octosporis; sporidiis elongatis, subclavatis obtusis, rectis vel curvulis, continuis, hyalinis 6 \approx 2; paraphysibus filiformibus.

Hab. in rhizomate Aspidii Filicis-maris in Britannia, Germania, Austria et Belgio.

79. **Dasyscypha filicea** (C. et Ph.) Sacc. *Peziza filicea* C. et Phil. 1879

Discom. p. 254. — Nivea, stipitata, sparsa; cupulis 0,1-0,2 mm. lat., parvulis, villosis, turbinatis, in stipitem brevem productis; ascis clavato-cylindricis; sporidiis fusiformibus, biseriatis, guttulatis,

denum pseudo-septatis 15-20 = 2-3; paraphysibus elongato-fusiformibus, acutis.

Hab. in *Filicibus*, Dunedin et Chedder Britanniæ. — Pili septati, leves, apice granuloso-capitati.

80. **Dasyscypha fimbriifera** (B. et C.) Sacc. *Pez. fimbriifera* B. et 1880
C. Cub. Fung. n. 681. — Nivea, minuta; stipite brevi decolorante
pruinoso; cupulis extus farinoso-tomentosis marginem versus floc-
cis longis rectis ornatis.

Hab. in stipitibus *Filicium* in insula Cuba.

81. **Dasyscypha Carestiana** (Rabenh.) Sacc. *Pez. Carestiana* Rabenh. 1881
F. Eur. n. 913, Hedw. V, p. 189, *Helotium* Karst. M. F. I, p. 161.
— Sparsa vel subgregaria, stipitata, alba, sicca pallida vel luteola
cupula concava, sicca subsphæroideo-contracta, pilis flexuoso-cur-
vatis, remote articulatis, subæqualibus, albidis dense vestita, 0,1
mm. lat., stipite brevissimo in cupulam abeunte; ascis cylindraceo-
clavatis, 32 = 4,5; sporidiis distichis, aciculari-elongatis vel acicu-
lari-oblängatis, guttulatis, 6-8 = 1,5-2; paraphysibus filiformibus,
sursum leviter incrassatulis, 1 μ . cr.

Hab. in *Asplenio Filice-femina*, in Italia bor. et Fennia.

82. **Dasyscypha Filicis-maris** (Saut.) Sacc. *Pez. Filicis-maris* Saut. 1882
Pilz. Salzb. p. 9, Wint. Hedw. 1881, p. 132. — Pusilla, ceracea, albi-
da, turbinato-cupularis, disco punctiformi, stipite brevi æquali;
ascis anguste clavatis, 30-35 = 3,5-5; sporidiis 8, distichis, oblongo-
clavatis, continuis, hyalinis, circ. 5 = 1.

Hab. in stipite vetusto *Aspidii Filicis-maris*, Mittersill
Salzburg. — Cupula pilis dense tecta, 1 mm. lat.; stipes vix visibilis.

83. **Dasyscypha Ulei** (Winter) Sacc. *Peziza Ulei* Winter in Hedw. 1883
1885, p. 258. — Hypophylla, sparsa, primitus globosa, clausa, dein
aperta, cupulæformis, margine in sicco semper connivente, hume-
ctata magis erecta, intense rubicunda, 260-340 μ . lata, pilis niveis,
longis asperulis densissime obsita, stipite brevi pallido prædita;
ascis cylindrico-subclavatis vertice rotundatis, deorsum parum at-
tenuatis, sessilibus, 8-sporis, 47-65 = 5-7; sporidiis subdistichis, fu-
siformibus utrinque acutis, parum inæquilateralibus, hyalinis, con-
tinuis, 15-18 = 2,5; paraphysibus sublinearibus lanceolatis i. e. e basi
cylindrica sursum parum incrassatis, apicem versus iterum at-
tenuatis et subacutatis, ascos superantibus plerumque parce ra-
mosis.

Hab. ad folia viva *Gleichenia dichotoma* prope S. Fran-
cisco in Brasilia (E. ULE).

84. **Dasyscypha ericina** (Quél.) Sacc. *Erinella ericina* Quél. XIV 1884

Suppl. p. 9, t. XII, f. 22. — Ascomatibus cyathiformibus, subtiliter velutinis, niveo-albis, 2-3 mm. diam., tenuibus; stipite filiformi, flexuoso, 2-6 mm. longo; hymenio luteo; sporidiis lanceolatis, 11 μ . longis, 2-3-guttulatis.

Hab. in muscis et in residuis vegetabilium in ericetis Alsatiæ.

85. **Dasyscypha erythrostigma** (B. et Br.) Sacc. *Peziza* B. et Br. 1885

Ann. N. H. 1866, n. 1168, t. IV, f. 31, nec Mont. — Minuta, stipitata, punctiformis, pallide rufa, pilis delicatis vestita; disco demum convexo; stipite curvato, distincte celluloso; ascis clavatis; sporidiis minutis, subglobosis vel ellipsoideis.

Hab. parasitica in *Chætosphæria phæostroma* in Britannia. — Stipes curvulus, distincte cellulosus. Quoad genus dubia.

86. **Dasyscypha Rhytismatis** (Phill.) Sacc. *Lachnella* Phill. Disc. 1886

p. 250. — Minuta, gregaria, stipitata, alba; ascomate primo globoso dein expanso, pilis sursum capitellatis tecto, stipite longiusculo, basi villoso; ascis cylindraceis, octosporis; sporidiis oblongo-fusiformibus 3-5 \times 1; paraphysibus asco longioribus, acute terminatis.

Hab. in Rhytismate acerino, Forres in Scotia (REV. JAMES KEITH). — Habitatione, sporidiis, pilis capitatis mox dignoscenda. Ascoma 300 μ . lat., stipes 400 μ . long.

B. Pili ascomatis colorati.

I. Trunci-lignicolæ.

87. **Dasyscypha cerina** (Pers.) Fuck. Symb. p. 305, Sacc. F. ital. 1887

fig. 1316, *Pez. cerina* Pers. Syn. pag. 651, Fr. Syst. Myc. II, p. 92, Nees Syst. f. 283, *Pez. marginata* Holmsk. Ot. II, p. 39, t. 20, ses. spec., *Pez. bicolor* Schum. Saell. p. 423, *P. lutescens* Pers. in Roem. Bot. Mag. p. 113, *P. pulchellæ* var. Pers. Myc. Eur. p. 261, *P. biformis* Fl. Dan. t. 1620, fig. inf., *Lachnella* Phill. Disc. p. 233, t. VII, f. 44, Pat. Tab. 170. — Hemisphærica, surfuraceo-villosa, lutescenti-olivacea, disco concavo luteo; pilis simplicibus, obtusis, septatis, asperulis, brunneis, 4-5 μ . cr.; ascis clavatis, 40 \times 5; sporidiis ellipsoideis, continuis, hyalinis, 5-8 \times 2,5-3.

Hab. ad ligna putrida frequens in Suecia, Fennia, Britannia, Italia, Gallia, Belgio, Amer. bor. — Stipes nunc distinctus, brevis 1 mm. fere long., crassus, sursum fere attenuatus, glaber, ater; nunc omnino nullus; cupula sœpe connivens, colore expallente, etiam in

disco, qui junior obscurus, demum subochraceus; sapor amarus.

— ***D. grisea** Pers. Myc. I, p. 264: minuta, sessilis, lutescenti-grisea, cupulis globosis duriusculis pubescentibus, ore minuto aperto. Ad ligna. Levis varietas. — ***D. cerinella** Spieg. F. Arg. II, p. 91 (*Helotium*): laxe gregaria, sessilis, villosula, hemisphærica, sicca globoso-contracta, minuta $\frac{1}{3}$ - $\frac{1}{2}$ mm., griseo-fulvescens, disco cereo, roseo-alutaceo; ascis cylindraceo-subclavatis, 40-45 \approx 3-5, paraphysibus filiformibus vix longioribus obvallatis, obturaculo iodi ope dilute cœrulecente; sporidiis distichis, elliptico-elongatis, utrinque acute rotundatis, 7-8 \approx 3-3½, primo continuis, dein medio (spuria?) uniseptatis, hyalinis. In fragmentis ligneis saliciniis putrescentibus en la Boca del Riachuelo, Argentina.

88. **Dasyscypha calyculæformis** (Schum.) Rehm Ascom. n. 111, *Peziza* 1888 Schum. Saell. p. 425 et in Fl. Dan. t. 2032, f. 2, *Lachnum* Karst. M. F. I, p. 178. — Sparsa, stipitata, villosa, subinfundibuliformis, umbrino-fusca, disco lutescenti-fusco vel sordide lutescenti-fusco vel sordide lutescente, 1,5 mm. lat., stipite crassiusculo, cupula breviore; ascis cylindraceo-subclavatis, 60-60 \approx 4-6; sporidiis distichis, eguttulatis, vulgo rectis, 10-12 \approx 2 (Karst.), 9 \approx 2 (Rehm); paraphysibus apice acutis, ascos superantibus.

Hab. ad lignum vetustum salicinum etc. in Dania, Fennia, Britannia, Germania, Africa austr. — Stipes in specim. britannicus atro-brunneus, glaber. — Var. **latebricola** Rehm Ascom. n. 111, cfr. Cooke in Grevillea III, p. 128, t. 41, f. 180: differt a forma normali: sporidiis ellipticis, haud acuminatis, 1-cellularibus, hyalinis, 6-7 \approx 2,5, distichis in ascis clavatis 40-45 \approx 6, 8-sporis; paraphysibus lanceolatis, apice acuminatis et ascos superantibus 4 μ . cr.; pilis ascomatis filiformibus, obtusis, scabris, fuscis, 6 μ . cr. Ad ramulos putridos imprimis Rhododendri ferruginei in alpibus editis tyrolensisibus.

89. **Dasyscypha Stevensoni** (B. et Br.) Sacc. *Peziza Stevensoni* B. 1889 et Br. Ann. N. H. n. 1485. — Minuta subglobosa, colore rubente, extus minute granuloso-puberula; stipite brevissimo, superne incrassato; sporidiis minutis, subcymbiformibus 12-13 μ . longis.

Hab. in ligno decorticato, Glamis Britanniæ. — Simil adsunt hyphæ repentes albæ, ramos erectos exserentes, in quorum apice conidia generantur obovata, 25 μ . long. Sec. Phill. Disc. p. 235; ascoma 200 μ . lat.; stipes longiusculus vel subnullus; asci clavati; sporidia oblongo-fusoidea, 8 \approx 2; paraphyses filiformes; pili delicati, hyalini.

90. **Dasyscypha crassipes** (Wallr.) Sacc. *Pez. crassipes* Wallr. Fl. 1890
crypt. n. 2572. — Ascomatibus obconico-piriformibus fusco-rufe-
scentibus, vertice primum in osculum subrotundum, dein in discum
concavum rufum reclusis, extus fibrillis parcis brevibus pallide-
griseis obsitis, intus stipiteque curto crasso rufescentibus.
Hab. ad caudices Carpin i cæsos in Thuringia (WALLR.).
91. **Dasyscypha atro-olivacea** Fuck. Symb. mycol. App. I, p. 337. 1891
— Cupulis gregariis, 1 mm. diam., in stipitem deorsum attenuatis,
subturbanatis, quandoque obliquis, pilis densis, olivaceis, articulatis
obsitis, antice subglobosis, ore connivente, umbilicato, albo-mar-
ginato, disco minuto, concavo, fusco; ascis stipitatis, cylindraceis
8-sporis, $48 \approx 8$; sporidiis oblique monostichis, ovatis, utrinque
subapiculatis, continuis, hyalinis $6 \approx 3$; paraphysibus filiformibus.
Hab. in caudicibus putridis Aceris Pseudoplatani pr.
Reichartshausen Rhenogoviæ. — *D. cerinæ* affinis.
92. **Dasyscypha ochracea** (Schw.) Sacc. *Peziza ochracea* Schw. Syn. 1892
p. 172. — Major quam *D. virginea*, eleganter ochracea, extus vil-
losa villis concoloribus, stipite mediocri; cupula dilatata, primum
contractim subclausa, disco vivide ochraceo.
Hab. in ligno pineo Kaign's Point, Philadelphia Amer. bor.
— Gregibus latis lignum non corruptum occupat.
93. **Dasyscypha arida** (Phill.) Sacc. *Peziza arida* Phill. Grev. V, 1893
p. 117, t. 89, fig. 13. — Stipitata, sparsa, globosa, dein expansa,
firma, extus setis tuberculatis, brunneis, septatis vestita; stipite
brevi, gracili, brunneo; ascis cylindraceis, octosporis; sporidiis
ovoideis, $7 \approx 4$; paraphysibus filiformibus.
Hab. in cortice Pin i, Blue Canon, Sierra Nevada Amer. bor.
— Ascomata 8 mm. diam., in sicco *Hysterium* simulantia.
94. **Dasyscypha variegata** Fuck. Symb. Myc. App. II, p. 61 et App. 1894
III, p. 31. — Cupulis gregariis, plus minusve dense stipatis, car-
noso-coriaceis, primo globosis, clausis, demum disco concavo, sor-
dido vel dilutissime rubello-fusco apertis, breviter sed distinrete
stipitatis, 1-2 mm. lat., orbicularibus seu pressione irregularibus,
marginatis, siccis margine involuto clausis, extus stipiteque pilis
crispulo-subtomentosis, primo fuscis, demum, præcipue marginem
versus, albo-variegatis obsitis; ascis oblongo-cylindraceis, stipitatis,
8-sporis, $80 \approx 8$; sporidiis primo distichis, demum in asci superiore
parte oblique monostichis, fusiformibus, parum inæquilateralibus,
continuis, hyalinis, $12 \approx 4$; paraphysibus simplicibus, utrinque at-
tenuatis.

Hab. in truncis putridis Fraxini Reichartshausen Rhenogoviæ. — Status conidicus, teste Fuck., sistit *Sarcopodii* speciem acervulis plerumque effusis confluentibusque, tomento saturatiore rubello-fusco, margine albo, conidiis 8-1 in sporophororum simplicium verticalium apicibus orientibus.

95. **Dasyscypha vix-visibilis** (Schw.) Sacc. *Pez. vix-visibilis* Schw. 1895

Syn. p. 175. — Minutissima subfasciculatim proveniens, fusca, subfibrillosa, obconica, abiens in stipitem brevem, margine albo pruinoso involuto, disco cinereo-fusco ascigero, ore semper contracto.

Hab. in fibris interioribus corticis Castaneæ, Bethlehem Amer. bor. — Vix nisi per lentem detegenda. Provenit circum *Sphæronæma parabolicum*.

96. **Dasyscypha turbinulata** (Schw.) Sacc. *Peziza turbinulata* Schw. 1896

Syn. p. 173. — Stipitata, gracilis, sparsa, turbinata, extus eleganter sulphurea pilosa, stipite versus basim subnudo subsulcato, disco vivide aurantio-rubro.

Hab. in rimis corticalibus Castaneæ in Amer. bor. — Vix 1 mm. alta.

97. **Dasyscypha uncinata** (Phill.) Sacc. *Peziza uncinata* Phill. Grev. 1897

V. pag. 117. — Minuta, stipitata, sparsa, pilis brevibus cinereis, sursum incurvis tecta; disco cœruleo-griseo; ascis cylindraceis v. subclavatis, 30-4; sporidiis fusiformibus rectis v. vix curvis, 6-1.

Hab. in ligno quercino et in acubus Pini, Blue Canon, Sierra Nevada, in montuosis Amer. bor.

98. **Dasyscypha prolificans** (Schw.) Sacc. *Peziza prolificans* Schw. 1898

Syn. p. 172. — Minuta, stipite tenui crassiusculo, subfasciculata, sicca clausa et sphæriomorpha, extus æruginoso-farinacea, pubescens, uda aperta, disco atro-viridi, saepe prolificante (i. e. cupula in cupula crescenti), limbo plerumque undulato, margine tumido.

Hab. in disco cæso ramorum et truncorum, Salem Amer. bor.

99. **Dasyscypha sericea** (A. et S.) Sacc. *Pez. sericea* A. et S. Lus. 1899

p. 325, t. III, f. 7, Fr. Syst. Myc. II, p. 93. — Cupularis, breviter stipitata, extus villo viridi circa marginem albicante sericeonitente adpresso tecta, intus ochraceo-ferrugineo.

Hab. in lignis nudis duris virore late tinetis, locis silvaticis umbrosis madidis in Lusatia. — Color initio totus pulchre viridis, extus profundior subfuscescens; sed in adultis in ochraceo-ferrugineum, nonnumquam sordide olivaceo-sordide olivaceo-intus mutatur, villorumque simul apices una cum cupulæ margine introflexo albescunt. Stipes firmus, crassus. Cupula 2-6 mm. lata, demum tota hians.

100. **Dasyscypha membranata** (Schw.) Sacc. *Peziza membranata* 1900
Schw. Syn. p. 175. — Submembranacea, extus rugulosa, fibris ad-
pressis, tota nigra, statu sicciori e stipite longiusculo abiens in
cupulam tubæformem, irregulariter limbo lobatam, intus cinere-
scens, statu madido tota cinerescens.
Hab. rara ad ligna dejecta, Bethlehem Amer. bor.
101. **Dasyscypha annulata** (Holms.) Sacc. *L. annulata* Holms. Ot. II, 1901
p. 30, t. 13, Fr. Syst. Myc. II, p. 106. — Stipitata, conferta, cu-
pulis subglobosis furfuraceis obscure cinereis, stipite annulato, di-
sco depresso nigro.
Hab. ad ligna putrida autumno in Dania. — Species dubii loci.
Stipes 5 mm. longus, annulo furfuraceo cinctus. Cupula globosa,
subclausa, cinereo-nigrescens, margine subconnivente.
102. **Dasyscypha armeniaca** (Pers.) Sacc. *P. armeniaca* Pers. Myc. 1902
Eur. p. 268, Fr. S. M. II, p. 95. — Gregaria, substipitata; asco-
matibus globoso-turbanatis, pubescenti-farinaceis, subcinnamomeis.
Hab. rarissime ad ligna exsiccata in Gallia. — Junior subglo-
bosa, villosa albicans, adulta magis pulveracea, subfulva.
103. **Dasyscypha borealis** (E. et H.) Sacc. *Lachnella borealis* Ell. 1903
et Holw. in Arthur Bull. III, Minn. p. 35. — Breviter stipitata,
circiter 1 mm. diam., globosa et vix aperta, dense tomentosa, sicca
atque pilis griseis (marginalibus 80-10 \approx 2½) tecta; disco pallido;
ascis cylindraceis, sessilibus, 75-80 \approx 7-8; paraphysibus validis, 2,5
 μ . crass., apice acutis, asco parum longioribus; sporidiis clavato-
fusoideis, nucleolatis, hyalinis, rectis vel leniter curvis, distichis,
20-22 \approx 3-4.
Hab. in ligno putri in America boreali. — Sicca *Cyphellam*
Ravenelii in mentem revocat.
104. **Dasyscypha luteo-alba** (Schw.) Sacc. *Peziza luteo-alba* Schw. 1904
Syn. p. 173. — Minuta, stipite gracili, in cupulam latiusculam
expanso, extus cum cupula luteo-villosa, disco aperto candido.
Hab. ad cortices Bethlehem Amer. bor.
105. **Dasyscypha illota** (B. et C.) Sacc. *Peziza illota* B. et C. Cub. 1905
Fungi n. 682. — Stipite brevi solido; cupula crateriformi, tomen-
tosa. cervina; hymenio rufo.
Hab. ad cortices in insula Cuba. — Ascoma 12 mm. diam. de-
scribitur; an revera *Dasyscypha*?

II. Ramicolæ

106. **Dasyscypha clandestina** (Bull.) Fuck. Symb. mycol. pag. 305, 1906

Peziza clandestina Bull. Champ. p. 251, Fr. Syst. Myc. II, p. 94, *Lachnum* Karst. M. F. I, pag. 178, Pat. Tab. 283. — Gregaria, stipitata, cervina v. griseo-lurida, furfuraceo-villosa; cupula turbinato-cyathoidea, disco pallido, 0,5-1 mm. lat.; stipite subæquali breviusculo circiter 0,4-0,5 mm. alt., firmulo; ascis cylindraceo-clavatis, 42-45 = 4-5; sporidiis fusoideo-oblongatis v. fusoideo-elongatis, rectis, 5-8 = 1,5; paraphysibus grandiusculis, apice acutis, 5-6 μ . cr.

Hab. ad caules et ramos dejectos putrescentes, præsertim Rubi idæi, fruticosi, Sambuci, Solani, Quercus in Suecia, Fennia, Germania, Gallia, Belgio, Amer. bor. — Pili articulati, fusi, 4,5-5 mm. crass., apice subclavato.

107. **Dasyscypha albo-furfuracea** (Saut.) Sacc. *Pez. albo-furfuracea* Saut. Pilzfl. Salzb. p. 8. — Subhemisphærica, furfuraceo-villosa, albida, margine inflexo, disco planiusculo, albo, stipite linearis.

Hab. in sarmenis Rubi idæi pr. Mittersill Salzburg. — $1\frac{1}{2}$ -1 mm. crassa, *P. cerinæ* propinqua, teste Saut.; sed sec. Wint. Hedw. 1881, p. 130 vix diversa a *D. clandestina*

108. **Dasyscypha scabro-villosa** (Phill.) Sacc. *Peziza* Phill. Grev. VII, 1908 pag. 22. — Gregaria, stipitata; disco albo-lutescente; ascomatibus extus pilis longis, rugosis, saepe capitato-granulosis tectis; ascis cylindraceis, octosporis; sporidiis fusiformibus, 15-21 = 2-3; paraphysibus latis, sursum acutis, asco longioribus.

Hab. in sarmenis Rubi nutkani in California.

109. **Dasyscypha Lentaginis** (Schw.) Sacc. *Peziza Lentaginis* Schw. 1909 Syn. p. 115. — Sparsim gregaria, breve stipitata, stipite tenuisculo, concolore, cupula dilatata, extus tenuissime fibrosa, badia, subpulverulenta, disco rotundo aut flexuoso marginato, pallide fusco-badio; tota 2-4 mm. lat.

Hab. in ramis Viburni Lentaginis, Hope Jersey Amer. bor. vulgata.

110. **Dasyscypha epixantha** (Cooke) Sacc. *Peziza epixantha* Cooke 1910 Grev. VII, p. 3. — Stipitata, brunnea, extus floccosa; cupula cyathiformi, 1 mm., disco pallidiore; ascis cylindraceis; sporidiis linearibus, 0,1 mm.; paraphysibus fusiformibus, utrinque acutis.

Hab. in ramis quercinæ in California Am. bor. (HARKNESS). — Pili brunnei septati sursum pallidiores, capitulo subgloboso ruguloso hyalino deciduo ornati.

111. **Dasyscypha succinea** (Phill.) Sacc. *Peziza succinea* Phill. Grev. VI, p. 116, t. 89, f. 12. — Stipitata, sparsa vel aggregata, glo-

bosa, dein exspansa, concava, firma, extus floccis lutescentibus tecta; stipite brevi, rufo-brunneo, disco carneo; ascis cylindraceis, octosporis; sporidiis ovatis, $10-14 \times 3-5$, paraphysibus linearibus, septatis.

Hab. in ramis emortuis quercinis, Blue Canon, Sierra Nevada in montuosis Amer. bor. — Ascomata 5 mm. diam.

112. **Dasyscypha sulphurella** (Peck) Sacc. *Peziza sulphurella* Peck 1912

30 Rep. p. 59. — Pallide lutea; ascomatibus minutis, numerosis, stipitatis, siccis clausis, pilosis, pilis septatis, rugosis, capitatis; ascis cylindraceis; sporidiis oblongis vel subcylindraceis, $7,5 \mu$. longis; paraphysibus asco longioribus sursum acutis.

Hab. in ramis Myricæ Gales, Adirondack Amer. bor. — Differt a *D. brunneola* colore atque stipite piloso. Totus fungus quam *D. myricacea* valde minor.

113. **Dasyscypha Ellisiana** (Rehm) Sacc. *Peziza Ellisiana* Rehm in 1913

Grev. IV, p. 169, *P. calycina* Schwz. in Herb. Soc. Nat. Philadelphia sec. Ellis Myc. univ. n. 716. — Sparsa, primitus hemisphærica, dein breviter stipitata atque dilatata, 1-2 mm. lata, luteo-villosa, disco aurantiaco-concaviusculo; sporidiis fusiformibus, utrinque acuminatis, simplicibus, hyalinis, $18 \times 2,5$, biseriatis; ascis clavatis, sessilibus, $45-60 \times 6-7$; paraphysibus filiformibus, ascos superantibus, septatis, 2μ . cr.

Hab. in ramulis aridis Pini rigidæ Newfield N. J. Amer. bor. (J. B. ELLIS). — Pili ascomatis di lute viridi-lutescentes, obtusi, simplices, scabri, 6 mm. crass. Apex ascorum jodii ope cœrulescit.

114. **Dasyscypha fusco-sanguinea** Rehm Ascom. n. 112, Cooke Grev. 1914

III, p. 122, t. 41, f. 179, *Lachnella* Karst. Symb. XXIV, p. 16. — Sparsa, primitus globosa, hirta, dein breviter stipitata; stipite circ. 1 mm. longo, patellæformis, extus cum stipite parum crasso fusco-badia, villosa, sicca vario modo complicata, humectata explanata, disco plano, recenti sanguineo-roseo, exsiccatu roseo-flavido, 3 mm. diam., orbicularis, prosenchy matico-contexta, pilis brevibus, scabris, obtusis, simplicibus, fuscis, septatis, $0,5 \mu$. crassis obsessa, margine flavidæ-ciliata; ascis clavatis, 8-sporis 70×10 ; sporidiis obtuse ellipticis vel ovato-elongatis, interdum subbiscoctiformibus, 1-cellularibus, hyalinis, distichis, 15×5 , rectis; paraphysibus rigidis, filiformibus, haud clavatis, fuscidulis, 2μ . cr.

Hab. ad ramos siccis Pini Cembræ et pumilionis in alpibus editis tyrolensisibus et Pini silvestris in Fennia. — Sec.

Karst. ascoma albido-tomentellum; asci 90-100 \approx 9; sporidia 15-20 \approx 6. An species diversa? Teste Rehm *D. arida* Phill. est hujus varietas.

115. *Dasyscypha flavo-virens* Bres. Trid. p. 92, t. 104, f. 1, Rehm 1915

Ascom. n. 762. — Stipitata ex hemisphærico plano-concava, 2-4 mm. lata, extus brunneo-olivacea, hirsuta, disco aureo-vitellino; stipite fusco-olivaceo, villoso, 1- $1\frac{1}{2}$ mm. longo, $\frac{1}{2}$ mm. crasso, substantia ceracea, albida; ascis cylindraceis basi parum constrictis jodo haud tinctis 70-80 \approx 7-8, (50 \approx 7 sec. Rehm); paraphysibus filiformibus, septatis, ramosis, apice subincrassatis, 2 μ . latis; sporidiis late ovatis, hyalinis, 7-8 \approx 4-5, pili ascomatibus sub micr. fuscidulis, septatis, granulis minutis concoloribus conspersis, 120 \approx 4 mm.

Hab. ad ramulos Laricis laxe gregaria in silvis alpinis tridentinis et germanicis. — A concolore *D. cerina*, et ab affinibus *D. fusco-sanguinea*, *D. calyculiformi* sporidiis præsertim distinguitur. Excipulum grosse parenchymaticum, brunneum; pili asperuli, septati, obtusuli, brunnei, 120 \approx 6.

III. Foliicolæ.

116. *Dasyscypha brunneola* (Desm.) Sacc. *Peziza* Desm. Crypt. Fr. I, n. 1156, Grev. III, p. 125, t. 42, f. 2, *Lachnum* Karst. M. F. I, p. 180, *P. dryophila* Pers., Pat. Tab. fig. 284. — Sparsa, stipitata, cupula juniore subsphæroidea, dein (humida) plana, marginata, brunneo-villosa, disco albido vel albido-pallescente, circ. 0,7 mm. lat.; stipite glabro, eburneo, brevi vel brevissimo; ascis cylindraceo-clavatis, 38-42 \approx 4,5-5,5 jodo non cœrulescentibus; sporidiis fusoideo-elongatis vel fusoideo-oblongatis, vulgo rectis et eguttatis, 6-11 \approx 2-2,5; paraphysibus apice acutis, circ. 4 μ . cr.

Hab. supra folia exsiccata Quercus et Æsculi in Gallia, Italia, Britannia, Fennia, Amer. bor. — Var. **fagicola** Phill. Disc. p. 238: multo obscurior, pilis obtusioribus intusque granulosis; ascis tereti-clavatis; spor. 7-10 \approx 2. In foliis Fagi prope Bristol Britanniae.

117. *Dasyscypha parvula* (Bomm. et Rouss.) Sacc. *Lachnella parvula* Bomm. et Rouss. Fl. mycol. p. 127. — Ascomatibus minutis, sessilibus vel breviter stipitatis, amœne citrinis, extus pilis brevibus concoloribus tectis; ascis clavatis, 33 \approx 3; sporidiis linearibus, minutis.

Hab. in foliis emortuis quercinis, Watermael in Belgio.

118. **Dasyscypha fuscescens** (Pers.) Rehm Ascom. n. 457 (nec 458), 1918

Peziza fuscescens Pers. Syn. p. 654, Myc. Eur. p. 265, Fr. Syst. Myc. II, p. 95, Grev. III, t. 40, f. 154. — Stipitata, fuscescens, cupula hemisphærico-globosa, extus pilis badiis obsita; pilis marginalibus simplicibus, septatis brunneis, apice capitulo hyalino crystallophoro, 7-8 µ. cr. terminatis; ascis clavatis $42 \approx 6$; sporidiis tereti-oblongis, obtusis rectiusculis, hyalinis, $10 \approx 1$ (Phill.) v. $8 \approx 2$ (Fuck.); paraphysibus filiformibus.

Hab. ad folia dejecta Fagi in umbrosis Sueciæ, Germaniæ, Britanniæ, Amer. bor. — Gregaria, pusilla, humida semipatula, sicca globosa clausa. Pili præcipue circa marginem regularem ciliati. Sec. Phill. a *D. brunneola* differt præsertim paraphysibus filiformibus nec lanceolatis. Teste Fuck. st. conidioph. est *Sphaeridium vitellinum* Fres.

119. **Dasyscypha phæotricha** Dur. et Lév. Expl. Sc. Alg. t. 28, f. 5 1919

(Pez.) [absque diagnosi]. — Gregaria, cupulata, stipitata, villosula lateritio-fusca, margine incurvo, disco nigricante; pilis clavulatis septatis, fuscis; ascis fusoideis..... immaturis.

Hab. in foliis emortuis, dejectis Quercus Ilicis, in Algeria. — Ascoma 1 mm. lat., stipes subæquilongus, tenuis, badio-fuscus.

120. **Dasyscypha capillaris** (Fr.) Sacc. *Peziza capillaris* Fr. Syst. 1920

Myc. II, p. 89. — Tenera, fusco-pallida, cupula subhemisphærica, extus stipiteque longo capillari deorsum nigricante villosa.

Hab. inter folia dejecta Fagi locis irriguis in Suecia. — Stipes gracilis, longitudine varius, 3-4 mm. longus et ultra, flexusus. Cupula primo globoso-clavata, dein hemisphærico-concava, demum planiuscula, subpellucida, 2-3 mm. lata. Color virens.

121. **Dasyscypha puberula** (B. et C.) Sacc. *Peziza puberula* B. et C. 1921

North Am. Fungi n. 729. — Cupulis globosis, cervinis, furfuraceo-tomentosis; pedicello brevi pallido; disco concolore, sero expanso, lætiore.

Hab. in areis decoloratis foliorum dejectorum Fraxini in Carolina inf. (RAVENEL).

122. **Dasyscypha Vincæ** (Lib.) Sacc. *Peziza* Lib. exsicc. n. 325, *He-* 1922

lotium Fuck. Symb. p. 316 et Lambotte Fl. Belg. II, p. 547. — Ascomatibus sparsis, minutis, planis, orbicularibus, carnosoceraceis, rufo-brunneis, extus griseolis, pilosis, stipite brevissimo, papilliformi, nigro; ascis elongatis, stipitatis, octosporis, $54 \approx 6$; sporidiis oblique distichis, ovato-oblongis, rectis curvatisve, $10 \approx 4$.

Hab. in foliis *Vincæ minoris* in Arduennis, Belgio et Germania.

123. **Dasyscypha callochætes** (E. et E.) Sacc. *Lachnella callochætes* E. et E. Journ. Myc. 1888, pag. 99. — Stipitata, 1 mm. alta, $\frac{1}{2}$ - $\frac{3}{4}$ mm. lat. extus stipiteque pilis rectis, nigris, fragilibus, obscure septatis, 150-200 \approx 6-7, sursum pallidis, basi vix incrassatis tecta; ascis clavato-cylindraceis, 60-70 \approx 7-8, paraphysibus copiosis filiformibus cinctis; sporidiis distichis, oblongo-fusoideis, hyalinis, plasmate imperfecte 1-2-partito, 12-14 \approx 2,5-3, inæquilateris.

Hab. in foliis delapsis *Myricæ ceriferæ*, Newfield in America boreali.

124. **Dasyscypha Eucalypti** (Berk.) Sacc. *Peziza Eucalypti* Berk. Fl. Tasm. pag. 274, t. CLXXXIII, f. 13. — Pallide olivacea, cupula plana, margine pilis rigidis, atro-purpureis ciliato, stipite cylindrico; sporidiis 10-11 μ . longis, oblongo-cymbiformibus.

Hab. in foliis *Eucalypti* in Tasmania (ARCHER). — Cum *P. misella* Desm. comparanda, etsi certe diversa.

125. **Dasyscypha fuscidula** (Cooke) Sacc. *Peziza fuscidula* Cooke Disc. U. S. II, p. 22. — Sparsa vel subgregaria, brunnea, stipitata, pilis septatis brunneolis margine fimbriata, disco pallido; ascis cylindraceis minutis; sporidiis spermatioideis.

Hab. in foliis *Andromedæ* New Jersey Amer. bor. — Dif- fert a *Trich. marginata* Cooke statura majore, stipite distincto, pilis diversis, paraphysibus acutis, disco pallidiore. Stipes vero variat subinde brevior.

126. **Dasyscypha subhirta** (Schw.) Sacc. *Peziza subhirta* Schw. Syn. p. 173. — Minutissima, substipitata, sparsa, lutea, extus minutim hirsuta, ascomate dilatato.

Hab. epiphylla, Bethlehem Amer. bor.

127. **Dasyscypha trichodea** (Ph. et Plowr.) Sacc. *Peziza Ph. et Pl.* Grev. III, pag. 125, t. 42, f. 4, *Lachnella Phill. Disc.* p. 234. — Gregaria vel sparsa, stipite brevi; ascomate primo globoso, dein cupulato, pallide brunneo, pilis rigidis, longis fulvis extus tecto, disco pallide albo; ascis tereti-clavatis 30 \approx 5; sporidiis distichis, oblongis, 7 \approx 1,5 binucleatis.

Hab. in foliis emortuis subputridis *Pini silvestris*, Trefriw, North Wales Britanniæ. — Ascoma 200-300 μ . lat., pili basi incrassati, continui.

128. **Dasyscypha pulverulenta** (Lib.) Sacc. *Peziza pulverulenta* Lib. 1928

Plant. Arduenn. n. 125, *Lachnum* Karst. M. F. I, p. 175, *Trichopeziza* Fuck. Symb. p. 297. — Gregaria aut subconferta, breviter stipitata vel subsessilis, alba vel albida, villosa; cupula primum subsphæroidea, dein planiuscula, sicca subsphæroideo-contracta, villo sulphureo-albo, aureo-pulverulento obsessa, 0,7-1 mm. lat.; stipite brevissimo vel fere nullo; ascis cylindraceo-subclavatis, 30-34 \approx 3-4; sporidiis fusoideo-oblongatis vel fusoideo-elongatis, 6-8 \approx 1,5-2; paraphysibus apice acutis.

Hab. ad folia *Pini silvestris* putrescentia in Gallia, Germania, Belgio, ad folia *Pini rigidæ* N. Y. Amer. bor. — Pili ascomatum 3 mm. crass., apice subclavato-obtuso.

129. **Dasyscypha solfatara** (C. et Ell.) Sacc. *Peziza solfatara* C. et E. Grev. VII, p. 7, Phill. Disc. p. 246. — Sparsa; cupulis hemisphæricis, citrinis, subtus papillato-affixis v. breve stipitatis, tomentosis, pilis asperulis, granulato-capitatis; ascis cylindraceis; sporidiis linearibus, 8 \approx 1; paraphysibus filiformibus.

Hab. in foliis pineis, New Jersey Amer. bor. et pr. Bristol et Carlisle Britanniae.

IV. Caulicolæ.

130. **Dasyscypha caulincola** (Fr.) Sacc. *Peziza caulincola* Fr. Syst. Myc. II, p. 94, *Helotium caulincola* (Fr.) Karst. M. F. I, p. 134, *P. albomarginata* Schum. Saell. p. 436, *P. Serratulæ* Pers. Myc. Eur. I, p. 263. — Lutescenti-fusca, pallida, cupula turbinato-globosa farinoso-pubescente, stipite brevi firmo glabro; ascis cylindraceo-clavatis; sporidiis oblongo-fusoideis, 7 \approx 1-1,5, hyalinis; paraphysibus fusoideo-acutis, ascos excedentibus.

Hab. ad caules plantarum frequens in Suecia, Fennia, Britannia, Germania, Belgio, Italia, in Arundinaria Amer. bor., in foliis Palmarum Cuba. — Minuta, gregaria, firmula persistens, demum glabrescens; tum ægre dignoscitur a *Phialea cyathoidea*, cum qua convenit loci, substantiæ, formæ et coloris mutationibus. Color pallescens, sordide et opace lutescens, pallide umbrinus etc. Discus ob marginem inflexum suboccultatus.

131. **Dasyscypha longipila** (Peck) Sacc. *Peziza longipila* Peck Bull. N. Y. Mus. 1887, p. 20, t. II, f. 15-19. — Ascomate minuto, 350-500 μ . diam., inferne in stipitem brevem attenuato, pilis rigidis, erectis, septatis, alutaceo-brunneis, longis (superioribus 250-350 \approx 3), dense tecto; disco albescente, setis marginalibus in sicco abs-

condito; ascis cylindraceis, 65-75 \approx 6,5-7,5; sporidiis oblongis vel subfusiformibus, rectis vel vix curvis, 7,5-10 \approx 2-3, hyalinis.

Hab. in caulis Eupatorii maculati, Adirondack Amer. bor. — A *Tr. relicina* ascomate stipitato non castaneo differt.

132. **Dasyscypha fuscata** (Rehm) Sacc. *Helotium fuscatum* Rehm 1932
Ascom. Alp. n. 5. — Sparsa v. gregaria, crasse stipitata, c. 1 mm. alt., primitus globosa, dein turbinata, demum cyathiformis, extus flavidio-puberula, margine subinvoluto, striatulo, humida explanata, urceolata; disco rubro-fusco, c. 1-2 mm. diam., parenchymatico-fuscidulo-contexta, fibris marginalibus brevibus, obtusis, subscabris, flavidulis, distantibus, c. 3 μ . cr.; ascis clavatis, 8-sporis, 50 \approx 5, apice jodo dilute cœrulecente; sporidiis elongatis, obtusis, rectis, 1-cellularibus, hyalinis, distichis, 6 \approx 2; paraphysibus hyalinis filiformibus.

Hab. ad caules putridos Aconiti Napelli prope glaciem æternam, Taschach-Gletscher in valle Piz alpium tyrolensium 2000 m. alt. (DR. REHM).

133. **Dasyscypha cypheloides** (Rehm) Sacc. *Trichopeziza cypheloides* 1933
Rehm Ascom. Alp. n. 12. — Gregaria, interdum sparsa, lentiformiter-globosa, breviter stipitata, dein fere sessilis, nigro-purpurea, hirta, parenchymatico-fusco-contexta, pilis longis, fusco-purpureis, simplicibus, subscabris, obtusis, c. 5 μ . cr., obsessa, primitus subclausa, humida urceolato-aperta, disco pallido, 1 mm. diam.; ascis clavatis, apice incrassatis, 8-sporis, 45 \approx 6, apice jodo cœrulecente; sporidiis fusiformibus, rectis vel subcurvatis, subobtusis, distichis, 1-cellularibus, interdum 2 nucleis instructis, hyalinis, 10 \approx 5; paraphysibus ascos superantibus, filiformibus, circ. 2 μ . diam.

Hab. ad caules putridos Aconiti Napelli juxta glaciem æternam alpium tyrolens. Taschach-Gletscher in valle Piz, alt. 2000 m. (DR. REHM), Kartel-Gletscher in valle Moos (BRITZELMAYR). — Color ascomatis in aqua solubilis.

134. **Dasyscypha vestita** (Quél.) Sacc. *Helotium vestitum* Quél. 11, 1934
Suppl. p. 22, t. XII, fig. 13. — Tubiformis, ceracea, citrina, velo surfuraceo, brunneo vestita; stipite basi attenuato, flexuoso, 4-5 mm. longo, hymenio plano, 3 mm. lato, citrino, fimbriis veli marginato; sporidiis cuneatis, elongatis, 22-24 μ . longis, guttulato-granulosis.

Hab. fasciculatim in caulis plantarum exsiccatis in Jura.

135. **Dasyscypha Cyathula** (Pers.) Pez. *Cyathula* Pers. Mycol. Eu- 1935

rop. I, p. 265, Weinm. 444, *P. caulincola* β. *Cyathula* Fr. Syst. Myc. 2, p. 95. — Gregaria sparsaque, firmula, ascomate subferrugineo, infundibuliformi, extus tomentoso; stipite sursum incrassato, demum glabro.

Hab. ad stipites dejectos plantarum in Rossia minore: legit Dr. BONGARD (Herb. Ac. Sc. Petrop.). — Species pulchella! Ascoma primo clausum, dein satis late apertum, intus parum dilutius, margine subintumescente subinvoluto. Stipes initio villosus, concolor, dein glaber, basi nigrescens.

136. **Dasyscypha stipiticola** (Schw.) Sacc. *Peziza stipiticola* Schw. 1936

Syn. p. 172. — Minuta, subsparsa, substipitata; cupulis globoso-turbinatis in stipitem crassum abeuntibus, primum clausis, extus farinaceo-pubescentibus, margine albescientibus; pilis ceterum cervinis; disco pallido.

Hab. in stipitibus dejectis, Bethlehem Amer. bor.

V. In Monocotyledoneis.

137. **Dasyscypha groenlandica** (Rost.) Sacc. *Lachnella groenlandica* 1937

Rostr. Fungi Groenl. p. 537. — Gregaria, brevi stipitata, badiostrigosa, turbinato-cyathoidea, 0,5-1 mm. alt.; ascis anguste cylindraceo-clavatis, circiter 64 ± 5; sporidiis fusoideo-elongatis, 9-10 ± 1,2, hyalinis, continuis; paraphysibus filiformibus.

Hab. ad caules exsiccatos Platantheræ hyperboreæ, Kob. befjord Groenlandiæ.

138. **Dasyscypha lanariceps** (B. et Ph.) Sacc. *Peziza lanariceps* C. 1938 et Phill. Grev. VIII, p. 62. — Sparsa, stipitata, ochraceo-fusca; cupulis turbinatis, demum apertis, 0,2 mm. diam., villosis, granulis purpureis conspersis; ascis clavatis; sporidiis cylindricis, utrinque attenuatis, 15-20 ± 3; paraphysibus elongato-fusiformibus, acutis.

Hab. in Rhipogono (?) Melbourne Australiæ. — Pili extiores 80-90 μ. long., rugulosi, granulis purpureis mixti.

139. **Dasyscypha Oncospermatis** (B. et Br.) Sacc. *Pez. Oncosperma-* 1939 *tis* B. et Br. Fung. of Ceyl. n. 942. — Cupulis breviter stipitatis cyathiformibus, tomentosis, luteis; sporidiis fusiformibus angustis, hinc acutis illinc obtusiusculis 20-25 μ. long.

Hab. ad *Oncosperma fasciculatum*, Habgalla Ceylon. — Minuta, pallide flava, sparsa, 300 μ. alta.

140. **Dasyscypha Staritzii** (Rehm) Sacc. *Lachnella Staritzii* Rehm 1940

Ascom. Ill. et Exs. n. 709. — Brevissime stipitata, patellaris, primus subglobosa, dein urceolata, disco aurantiaco-rubro, extus ro-

seo hirtello, margine imprimis pallidiore, subpruinoso, gregaria, flavidо-rubra, prosenchymerico-contexta, c. 0,5-8 mm. diam., pilis crebris, simplicibus, obtusis, subscabris, hyalinis v. rubro-fuscidulis, 30-4-5 obsessa; ascis clavatis, 45-6, 8-sporis; sporidiis fusiformibus v. bacillaribus, plerumque rectis, 1-cellularibus, hyalinis, 10-12=1,5; paraphysibus lanciformiter acutatis, ascos longe superantibus, basi 5-6 μ . cr.

Hab. ad calamos Junci sicclos prope Königstein a. d. Elbe Saxoniæ.

141. **Dasyscypha Rehmii** (Star.) Sacc. *Trichopeziza Rehmii* Staritz 1941 in Rehm Ascom. n. 658. — Sparsa, sublentiformis, calyciformiter tenuiter stipitata, circ. 1 mm. alta, sicca ore involuto, pilis fuscis obpresso subclausa, humida usque ad 1-1 $\frac{1}{2}$ mm. diam. dilatata, disco pallide fuscidulo, concavo, extus nigro-fusca, hirtella, contextu prosenchymerico chalybeo, pilis rectis longiusculis, simplicibus, obtusis, vix septatis, scabris, fuscis, basi saepe chalybeis, c. 4-5 μ . cr. circa osculum obsessa; ascis cylindraceis, sessilibus, 8-sporis, 45-8; sporidiis distichis, ellipticis, rectis, 1-cellularibus, plerumque binucleatis, hyalinis, 12-2-2,5; paraphysibus lanciformibus, acutatis, hyalinis, basi 5-6 μ . cr. ascosque longe superantibus.

Hab. ad calamos sicclos Junci conglomerati, Keulenberg prope Königsbrück Saxoniæ (STARITZ). — Ascomatis contextu chalybeo pilis brunneis mox dignoscenda species.

142. **Dasyscypha patens** (Fr.) Rehm Ascom. n. 156, *Peziza clandestina* β . *patens* Fr. Syst. Myc. II, p. 94, *Lachnum patens* Karst. M. T. I, p. 179, *Trichopeziza* Sacc. Mich. II, p. 330, *Pez. palearum* Desm. Ann. S. N. 1846, p. 365, *Lachnella* Phill. Disc. p. 239. — Sparsa vel subgregaria, stipitata, lutescenti-pallescens vel albido-fulva, furfuraceo-tomentosa, cupula turbinato-cyathoidea v. acetabuliformis, disco pallescente, circiter 0,5 mm. lat., stipite subæquali aut sursum paullulum incrassato, 0,4-0,6 mm. alt.; ascis cylindraceo-subclavatis, 35-46=4-5; sporidiis fusoideo-elongatis, rectis, plerumque guttulatis, 10-12=1,5-2; paraphysibus grandiusculis, apice acutis, 4-5 μ . cr.

Hab. in culmis Airæ, Psammæ, Tritici in Gallia, Germania, Fennia, Britannia, Belgio.

143. **Dasyscypha spherocephala** (Wallr.) Rehm Ascom. n. 263 (ut 1943 var.), *Peziza* Wallr. Fl. crypt. p. 456, *D. patens* var. *spherocephala* Rehm, *Lachnum patens* β . *spherocephalum* Karst. M. F. I, p. 180, *D. cephaloidea* Fuck. Symb. myc. pag. 306. — Sparsa,

stipitata, 2 mm. alta; stipitibus crassiusculis, parum curvatis, adpresso-pilosis, griseis; cupulis globosis, seminis Papaveris magnitudine, undique pilis patulis, brevibus, lutescentibus, siccis pallidis clavatis, septatis obsitis, semper clausis v. humectatis ore regulari, pusillo apertis, disco pallido; ascis linearibus, acuminatis, 8-sporis, 45-48 = 4-5; sporidiis fusiformibus, hyalinis, 6-8 = 2; paraphysibus lanceolato-acutis, ascos superantibus, 5-6 μ . cr.

Hab. in culmis siccis adhuc stantibus Tritici repentis, Secalis, Calamagrostidis in Thuringia pr. Östrich ad Rhenum et in Fennia. — *D. clandestinæ* affinis. Pili saepe apice crystallifero, 6 = 5.

144. ***Dasyscypha calycioides*** (Rehm) Sacc. *Peziza calycioides* Rehm 1944

Ascom. n. 159. — Sparsa, distincte plus minusve stipitata, stipite filiformi, plerumque curvato, nigro-fuscidulo, 0,3-8 mm. long. cyathoidea, badio-strigosa, c. 0,3 mm. diam.; disco flavidulo, prosenchymatico-flavidulo contexta; pilis fuscis, distincte articulatis, obtusis, apice subscabris, c. 6 μ . cr., longis sessa; ascis clavatis, 8-sporis, 36-45 = 6, apice jodo cœrulecente; sporidiis ellipticis plerumque rectis, 1-cellularibus, hyalinis, distichis, 9-12 = 2-2,5; paraphysibus lanceolatis, ascos longe superantibus, hyalinis, basi c. 5 μ . cr.

Hab. ad calamos sicclos Nardi strictæ in alpibus editioribus tyrolensisibus juxta glaciem æternam. — Var. ***microspora*** Rehm Ascom. n. 567: sporidiis 6 = 1,5. In culmis ibidem. Fugax, ægre observanda.

VI. In Acotyledoneis et ad terram.

145. ***Dasyscypha grisella*** (Rehm) Sacc. *Helotium grisellum* Rehm 1945

Ascom. n. 766. — Sparsa, imprimis in pagina inferiore frondium, turbinata; stipite brevi, fuscidulo; cupula grisella, subfurfuracea, circ. 0,3-4 mm. alt. et 0,2-3 mm. diam., subclausa, fuscidula sive subhyalina, prosenchymatico-contexta, pilis fuscidulis v. subhyalina, undulatis, simplicibus, 60 = 3, sessa; ascis clavatis, apice subacutatis, lata basi sessilibus, saepe subcurvatis, 8-sporis, 30 = 6-7; sporidiis bacilliformibus, oblongis, rectis v. subcurvatis, 1-cellularibus, hyalinis 6-8 = 1-1,5, 2-3-stichis; paraphysibus filiformibus, teneris, hyalinis, haud clavatis.

Hab. ad frondes emortuas, putrescentes Pteridis aquilinæ Königstein a. d. Elbe Saxoniæ (KRIEGER).

146. ***Dasyscypha seminis*** (C. et Ph.) Sacc. *Peziza seminis* Cooke 1946

et Phil. Grev. IX, p. 104. — Ochracea, stipitata; cupulis clavatis, demum concavis $\frac{1}{2}$ -1 mm. diam., extus tomentosis; stipite crasso, vix deorsum attenuato 1 mm. long.; ascis clavatis; sporidiis arcte ellipticis, minutis 5-2,5.

Hab. in Sclerotio semine in Arduennis (LIBERT).

147. **Dasyphypha terrestris** (B. et Br.) Sacc. *Helotium terrestre* B. 1947

et Br. Fung. Brisb. p. 69, t. XV, f. 15-17. — Ascomate stipitato parvo, planiusculo, cornicolori, lurido, extus villoso; ascis elongatis; sporidiis monostichis, ellipticis, utrinque breviter appendiculatis uniguttulatisque, 10 μ . longis; paraphysibus filiformi-clavulatis.

Hab. ad terram udam, Brisbane Australiæ. — Ascoma 4-5 mm. lat., stipes 1 mm. longus.

COMESIA Sacc. Conspl. gen. Discom. p. 6 (1884) [Etym. a cl. botanico et mycologo neapolit. HOR. COMES], *Helotii* spec. Auct. — Ascomata obconica, sessilia vel brevissime stipitata, disco plano, glabra v. minute ciliata, carnosoceracea. Ascii crassi, polyspori, paraphysati. Sporidia continua, hyalina, oblonga v. (in subg. *Comesiella*) subglobosa. *? Roma?*

I. *Eu-Comesia*. Sporidia tereti-oblonga.

1. **Comesia Felicitatis** (Crouan) Sacc. Conspl. gen. Discom. p. 6, *Helotium Felicitatis* Cr. Finist. p. 48, t. ult. fig. 20. — Sessilis, initio hemisphærica dein subcylindracea, carnea, disco plano; ascis fusoideis, polysporis; sporidiis cylindraceis utrinque truncatis, 4-5-guttulatis; paraphysibus capillaribus sursum lenissime incrassatis; contextu excipuli minute celluloso.

Hab. ad terram inter muscos et Glœocapsas, Finistère Galloë (Domina FELICITAS CROUAN). — Ascoma 1 mm. lat.

2. **Comesia cunicularia** (Boud.) Sacc. *Peziza cunicularia* Boud. 1949 Ascob. p. 68. — Minutissima, tota alba, sparsa, substipitata v. piriformis, extus glabra, margine minute ciliata, disco plano; ascis crassis subcurvatis, fissura bilabiata dehiscentibus; sporidiis numerosis (48-64), oblongo-fusoideis, hyalinis; paraphysibus septatis ascos subæquantibus.

Hab. ad sterlus Cuniculorum locis argillosis pr. Montmorency Galliæ. — Ascoma $\frac{1}{4}$ - $\frac{1}{2}$ mm. lat. Cfr. *Ryparobius argenteus*.

3. **Comesia Leveillei** (Crouan) Sacc. *Ascobolus Leveillei* Cr. Finist. 1950 p. 57, tab. ult. f. I, Boud. Ascob. p. 68. — Ascomate niveo, gla-

bro hemisphærico, stipite brevissimo cylindraceo, tomentoso; ascis clavatis, curvis; sporidiis 48 subcylindraceo-oblongis, utrinque rotundatis, intus tenuiter granulosis et plasmate fere bipartito; paraphysibus filiformibus erectis ascos superantibus, septatis.

Hab. in fimo Muli inter folia dejecta, Finistère Galliæ. — Ascoma $\frac{1}{2}$ mm. lat. An *Ryparobius*?

II. *Comesiella* Sacc. Sporidia globulosa.

4. **Comesia fusca** (Crouan) Sacc. *Helotium fuscum* Cr. Finist. p. 47. 1951

— Sparsa; ascomata 1 mm. diam. metiente, brunneo, sessili, subhemisphærico, convexo; ascis oblongo-clavatis, 16-sporis; sporidiis rotundatis, granulosis; paraphysibus filiformibus, apice brunneis.

Hab. ad terram nudam, Brest Galliæ.

TROMERA Mass. in Arnold Flora 1858, p. 507, Körb. Par. p. 453 (Etym. *tromeros* tremulus). *Pezizæ* sp. Fr., *Peziculæ* et *Retinocyli* sp. Fuck. *Biatorella* Mudd. — Ascomata superficialia sessilia v. subsessilia, obconico-patellata, submarginata, excipulo discoque concaviusculo subceraceis, hypothecio subcarnoso. Ascii oblongi, polyspori, paraphysati. Sporidia continua, hyalina globulosa v. oblonga. Pycnidia nectrioidea, sphærospora, teste Fuck. — Genus anceps, hic Lichenibus hinc Dermateis accedens.

1. **Tromera resinæ** (Fr.) Körb. Par. p. 453, Karst. Rev. p. 153, *Lecidea* Fr. Obs. I, p. 180, *Peziza* Fr. S. M. II, p. 149, *Pez. myriospora* Hepp. *Tromera myr.* Anzi, *Tr. xanthostigma* Mass. *Biatorella resinæ* Mudd., *Retinocyclus flavus* Fuck. Symb. App. I, p. 332. — Sessilis, primo urceolato-concava et margine dilutiore crasso cincta, dein planiuscula, margine evanescente, crassiuscula, dura, firma, rufo-aurantiaca, minuta, ascis clavatis seu elongato-ventricosis, 90 ± 18 jodo intense cærulescentibus; sporidiis numerosissimis, sphæroideis, continua, flavescentibus vel subhyalinis, 3μ . diam.; paraphysibus filiformibus, clavulatis, articulatis.

Hab. in resina et in cortice Abietis excelsæ, Pini silvestris etc. in Suecia, Fennia, Britannia, Germania, Gallia, Italia et America boreali. — Teste Fuckel, *Nectria Resinæ* Fr. (sporulis globosis flavidis 3μ . diam.) sistit st. pycnidicum. — Var. **stipitata** Karst. l. c. provenit in Fennia.

2. **Tromera difformis** (Fr.) Rehm Asc. n. 577, Karst. Rev. p. 154, 1953 *Peziza* Fr. S. M. II, pag. 151, *Lecidea* Nyl., *Tromera sarcogynoides* Mass., *Biatorella difformis* Wainio, *Tromera myriospora*

var. *sarcogynoides* Anzi. — Sessilis, firma, plana vel convexiuscula, marginata vel immarginata, irregularis, nuda, opaca, nigra, 0,5-0,8 mm. lata, disco fuscescente; ascis subclavatis, 50 \approx 12; sporidiis numerosis (50-80 circ.) sphæroideis, hyalinis, 2-2,5 μ . diam.; paraphysibus filiformibus, apice subramosis, incrassatis fuscidulisque.

Hab. in cortice et ligno resinoso Abietum in Italia, Suecia, Fennia, Germania, Austria, Helvetia. — Gelatina hymenialis jodo intense cœruleascit. — A *Tr. resinæ* ascomatibus junioribus atro-cœruleis mox dignoscenda.

3. **Tromera lignaria** Karst. Symb. Myc. Fenn. XXV, p. 23. — Superficialis, solitaria, raro in cæspitulos sociata (plerumque 3 insimul), sessilis, rarius basis constricta, aperta, concaviuscula, demum plana vel convexiuscula, margine obtuso, demum subimmarginata, nuda, atra, rugulosa, 0,3 mm. lat.; ascis clavatis obtusis, jodo dilute cœrulecentibus, 80-100 \approx 12-14, myriosporis; sporidiis ellipsoideo-sphæroideis, hyalinis, 1,5-2 \approx 1,5 vel 1,5 μ . diam.; paraphysibus filiformibus 1 μ . cr.

Hab. ad lignum abietinum vetustum prope Mustiala Fenniæ. — An *Tympanidi* adscribenda?

4. **Tromera microtheca** Karst. Symb. Myc. Fenn. XXV, p. 22. — Superficialis, sessilis, gregaria, solitaria, rarissime in cæspitulos sociata (2-3 insimul), nuda, nigra, opaca, subrugulosa, aperta, plano-orbicularis vel ovalis, tenuiter marginata, vix 0,3 mm. lata; ascis clavatis, obtusissimis, jodo dilute cœrulecentibus, 36-40 \approx 10-12, myriosporis; sporidiis oblongatis vel elongatis, rectis, hyalinis, 2-3,5 \approx 1,5; paraphysibus haud bene discretis; hymenio jodo fulvescente.

Hab. in ligno vetusto abietino ad Mustiala, Fenniæ. A præcedente, cui proxime accedit, præcipue ascis minoribus, sporidiis longioribus cupulisque minoribus et tenuius marginatis recedit.

5. **Tromera olivacea** (Fuck.) Sacc. *R. olivaceus* Fuck. Symb. mycol. 1956 App. I, p. 332. — Cupulis resina et plerumque villo olivaceo insidentibus, ceterum glabris, subgregariis, carnosis, subsessilibus, usque ad 2 mm. latis, primo clausis, margine distincto, atro, olivaceo, involuto, demum planis, disco carnoso, olivaceo, sicco atro; ascis elongatis, sessilibus, polysporis, 120 \approx 14; sporidiis minutissimis, perfecte globosis, simplicibus, hyalinis seu dilute flavescentibus, 3 μ . diam.

Hab. in ligno vetusto molli Laricis pr. Aepfelbach et Hall-

garten Rhenogoviae — Sec. Rehm Ascom. n. 577 ad *Tympnidem* species hæc pertineret.

Sect. 2. Phæosporæ Sacc.

PHÆOPEZIA Sacc. Mich. I, p. 71 (1877) Consp. gen. Disc. p. 6 (1884) [Etym. *phaios* obscurus et *Peziza*). — Ascomata carnosa v. subceracea, sessilia v. subsessilia, cupulata v. scutellata, fusca, rarius læte colorata, glabra v. pilosula. Asci elongati, paraphysati, 6-8-spori. Sporidia continua, globosa, ellipsoidea v. oblonga, fusca. — Genus hucusque paucas species continens, quæ vero series plures, hyalosporarum parallelas, constituunt; si species numerosæ innotuerint, dividendum.

I. *Eu-Phæopezia*. Sporidia globosa.

1. **Phæopezia murina** (Fuck.) Sacc. Consp. gen. Disc. pag. 6, March. 1957 Copr. p. 38, *Plicaria* Fuck. Symb. pag. 326, *Peziza* Fuck. F. R. n. 1597, Cooke Myc. f. 76. — Cupulis valde aggregatis, carnosis, pallide rufis, planis, in stipitem brevem, crassum attenuatis, margine crenulato, vix elevato, extus rugulosis, basi fibrillis repentibus, fuscis cinctis; ascis longissimis, cylindraceis, 8-sporis, longitudinaliter plicatis et demum quasi laceratis, 92 \times 12 (parte sporifer.); sporidiis in ascis superiori parte monostichis, perfecte globosis, 12 μ . diam. episporio crasso, granuloso-tuberculato, fusco.

Hab. in stercore putri murino et vulpino culmisque proximis in Rhenogovia et Belgio. — Cellulæ excipuli exacte hexagonæ. Ascomata, teste Marchel, sessilia v. subsessilia, 1-1,5 mm. lat., initio alba dein violacea; paraphyses asco longiores, 3,5-5 μ . cr., septatæ, hyalinæ, sursum subinde divisæ; ascis 150-180 \approx 16-17.

2. **Phæopezia melanosperma** Cr. Fin. pag. 50. — Sessilis, brunneo-1958 atra, subhemisphærica, fragilis, 1-2 cm. lat., margine lobato, disco brunneo, initio concaviusculo, dein convexo; ascis cylindricis, octosporis; sporidiis globosis, nigricantibus, granulosis; paraphysibus filiformibus, sursum furcatis, apice filiformi, brunneo.

Hab. ad terram adustam, Morlaix Galliæ. — Cellulæ excipuli globosæ, majusculæ, in filamenta septata, nodulosa abeuntes.

3. **Phæopezia Radula** B. et Br. Ann. N. H. XVIII, pag. 77, Cooke 1959 Mycogr. fig. 293. — Magna, sessilis, cupula hemisphærica, demum depressa, extus verrucosa, atra, intus atro-vinosa; ascis cylindraceis; sporidiis globosis, episporio tuberculatis, brunneis, 22-25 μ . diam.; paraphysibus septatis, sursum clavatis.

Hab. ad terram in silvis prope Bristol Britanniae. — Ascoma

2-3 cm. diam., extus verrucosum; verrucæ subconoideæ secedentes.

4. **Phæopezia atro-spora** (Fuck.) Sacc. *Pez. atro-spora* Fuck. F. rhen. 1960 n. 1224, Cooke Myc. f. 294, *Ph. ferruginea* Fuck. Symb. mycol. p. 326. — Ascomatibus sessilibus, primo globosis, clausis, dein hemisphæricis, margine involuto, subapplanatis, usque ad 2,5 cm. latis, extus verrucosis, ferrugineis, disco levi, obscuriori; ascis cylindraceis; sporidiis globosis, verrucosis, brunneis, 18-20 μ . diam.; paraphysibus filiformibus.

Hab. ad terram udam in silvis pr. Rabenkopf Germaniae.

5. **Phæopezia scabrosa** (Cooke) Sacc. *Peziza scabrosa* Cooke Mycogr. p. 170, f. 295. — Sessilis; cupula concava, demum expansa, atro-brunnea, extus verrucosa; margine primo subincurvo, regulari; ascis cylindraceis; sporidiis globosis, 15 μ . diam.; episporio tuberculatis, brunneis; paraphysibus supra clavatis.

Hab. ad terram pr. Portland Maine Amer. bor. (BOLLAS). — Ascoma 12-15 mm. lat.

6. **Phæopezia albo-fusca** (Cr.) Sacc. *Peziza albofusca* Cr. Fin. p. 54. 1962 — Ascomate 1 mm. diam., albo, basi filamentis mycelicis cincto, sessili, hemisphærico, dein concavo, margine tenui, translucido, erecto; hymenio brunneo; ascis cylindraceis, octosporis; sporidiis sphæroideis, granulosis, primo hyalinis dein brunneis; paraphysibus filiformibus.

Hab. in fimo muli ad Sphagna, Finistère Galliae.

II. *Aleurina* Sacc. Sporidia ellipsoidea v. oblonga.

7. **Phæopezia retiderma** (Cooke) Sacc. *Peziza retiderma* Cooke Mycogr. p. 176, f. 306. — Sessilis, badia, cupula hemisphærica, interdum contorta, integra, carnosa, extus reticulato-venosa, circa marginem pallidore; ascis cylindraceis; sporidiis ellipticis, brunneis, verrucosis, 14-7; paraphysibus numerosis, septatis, linearibus, sursum brunneis.

Hab. ad terram, Portland Maine Amer. bor. (FULLER). — Asco-
ma 2-4 cm. lat.

8. **Phæopezia applanata** (Rab. et Gonn.) Sacc. Conspl. gen. Disc. p. 6, 1964 *Peziza* Rabenh. et Gonn. Myc. Eur. t. VI, f. 5 (absque diagnosi) nec Fr. — Ascomatibus sessilibus scutellato-cupulatis, glabris, paleo castaneis, disco atro-castaneo, concavo; ascis cylindraceis, basi nodulosis, octosporis; sporidiis in ascii parte sup. oblique monostichis ellipsoideis, leniter inæquilateralibus, utrinque rotundatis, fu-

ligineis; paraphysibus filiformibus, sæpe e basi furcatis, apice lenissime clavulatis.

Hab. ad terram in Germania (?). — Ascoma ex icone 2 cm. lat.

9. **Phæopezia Puiggarii** Speg. Fung. Arg. IV, n. 220. — Uda applicata-scutellata 3-5 mm. diam., sicca varie contracto-corrugata, tenuiuscule submembranacea, gelatinoso-coraciuscula, glaberrima, extus intusque aterrima; ascis cylindraceis, sursum obtuse rotundatis crasseque tunicatis, deorsum longe attenuato-stipitatis, 80-100 \approx 6-6,5, octosporis, aparaphysatis, cito diffluentibus, muco obvallatis; sporidiis oblique monostichis, ellipticis, sæpe inæquilate-ralibus, utrinque obtusulis, 9-10 \approx 4-5, olivaceis, eguttulatis, continuis, corpusculis (spermatiis?) globosis vel ellipticis 3-3,5 \approx 1,5, hyalinis immixtis.

Hab. ad ramenta putrida coacervata prope Apiahym Brasiliæ meridionalis (Dr. J. PUIGGARI).

10. **Phæopezia apiculata** (Cooke) Sacc. Mich. I, p. 71, *Peziza* Cooke 1966 Mycogr. fig. 79. — Cupulis sessilibus, scutellatis, circ. 2-6 mm. diam., carnosso-tenacellis, atris, levibus, glabris, disco concavo, margine integro; ascis cylindraceis, deorsum crassiuscule attenuatis, apice subtruncatis, 200 \approx 18; paraphysibus filiformibus, septulatis apice fuscidulis obvallatis; sporidiis oblique monostichis, oblongo-ellipsoideis, utrinque breve mucronatis, 20-25 \approx 10-11, atro-fuscis; mucronibus tandem fuscis.

Hab. ad cortices putres et humum, Lucca et Padova Italiæ.

11. **Phæopezia elastica** Pat. et Gaill. in Bull. Soc. Myc. Fr. IV, p. 99. 1967 — Cupulis sessilibus, 1-3 mm. diam., sparsis vel gregariis, extus brunneo-fuscis rugosis, intus atris, matrici per fibrillas brunneas, hyphis septatis conflatas adfixis, margine crasso, excipulo compacto, elastico, bulgarioideo; ascis brunneis, cylindraceis, elongatis, arcte adpressis, octosporis; sporidiis monostichis, 13-15 \approx 6-7, brunneis, ovoideo-obtusis, levibus, biguttulatis; paraphysibus? indistinctis.

Hab. ad lignum emortuum Mapire Amer. austr. (GAILLARD).

— Pars exterior cupulæ fit cellulis magnis angulatis rugosa.

12. **Phæopezia phæospora** (Hazsl.) Sacc. *Pyronema phæosporum* Hazsl. 1968 Eing. Disc. pag. 16. — Carnosa, concava, albo-flavida, mediocris, mycelio albo inserta; disco brunneo-flavo, farinoso; ascis cylindraceis, in stipitem tenuem sensim attenuatis, 80 μ . long. dein clavatis; sporidiis senis (semper?), ovoides, brunneis, 12-16 \approx 6-7.

Hab. in horto botanico Buda-Pest Hungariae (Doct. DIETZ).

13. **Phæopezia crinita** (Bull.) Sacc. *Peziza crinita* Bull. ch. p. 249, 1969
t. 416, f. 2, Pers. Myc. Eur. pag. 255, Fr. Syst. Myc. II, p. 86,
Cooke Mycogr. f. 361. — Exigua, hemisphærica, ciliata, subtus
cinerea, disco rubro-purpureo; ascis cylindraceis; sporidiis ellipsoi-
deis asperulis, dilute fuligineis, 20 ± 12 ; paraphysibus sursum cla-
vulatis, fuscellis.

Hab. ad ligna putrida in Gallia (BULLIARD, LIMMINGHE). — Pla-
ne sessilis, sed non applanata, inferne, sed præcipue ad marginem
pilis confertis nigris septatis, 300μ . longis, hirta, 4-5 mm. lat.
Caro alba.

14. **Phæopezia fuscocarpa** (E. et Hol.) Sacc. *Peziza fuscocarpa* E. et 1970
Hol. in Journ. of Myc. I, pag. 5. — Sessilis orbicularis, 3-4 mm.
diam, extus pruinosa-tomentosa, olivaceo-flava, disco concavo, atro-
olivaceo, margine incurvo; ascis linearibus $65-80 \pm 4,5-5$; para-
physibus filiformibus; sporidiis monostichis, ovato-oblongis, brun-
neis, binucleatis, $7-8 \pm 3-3,5$.

Hab. in lignis vetustis, Jowa Amer. bor.

Sect. 3. Hyalodidymæ. Sacc.

- HELOTIELLA** Sacc. Conspl. gen. Disc. p. 6, 1884 (Etym. demin. ab
Helotio), *Pezizæ, Helotii et Mollisicæ* sp. Auct. — Ascomata mi-
nuta, ceracea, sessilia, plerumque pallida vel lacticoloria, discoidea
scutellata vel urceolata, glabra, scabra vel margine breve ciliata.
Asci typice octospori, paraphysati. Sporidia ovata, oblonga, vel
fusoidea, 1-septata, hyalina.

I. *Eu-Helotiella*. Ascomata margine pilosula vel scabra.

1. **Helotiella Citri** (Penz.) Sacc. Conspl. Disc. p. 6, *Peziza Citri* Penz. 1971
Mich. II, p. 399, F. ital. fig. 1130. — Ascomatibus gregariis, ses-
silibus, convexulis, lentiformibus, immarginatis, extus marginem
versus pilosiusculis, candidis, 0,2-0,4 mm. lat.; ascis cylindraceo-
clavatis, sessilibus, $50-60 \pm 10-20$, paraphysibusque jodi ope cœr-
lescentibus; sporidiis in asco distichis fusoideo-ellipticis, obtusiu-
sculis, plerumque medio septatis ibique haud constrictis, 2-4-gut-
tulatis, hyalinis, $9-12 \pm 4-5,5$.

Hab. in cortice ramorum Citri Limonum in frigidariis, Strà
prope Patavium (O. PENZIG).

2. **Helotiella cornuta** (Ell.) Sacc. *Peziza cornuta* Ellis Torr. Cl. 1882, 1972
p. 73. — Gregaria, minuta, sordide alba, sicca, carnea, sessilis,
tenui-membranacea, subfibrosa, margine pilis 4-5-septatis, albis,

sparse fimbriata; ascis clavato-cylindraceis, 50-60 \approx 8-10, filiformi-paraphysatis; sporidiis distichis, fusoideis, trinucleatis, demum uniseptatis, 12 \approx 3-4, hyalinis, utrinque appendicula brevi auctis.

Hab. in ligno dejecto Castaneæ, Westchester Pennsylvania Amer. bor.

3. **Helotiella carnosula** (Rhem) Sacc. *Peziza carnosula* Rehm Ascom. 1973

Alp. n. 4 (*Helotium*). — Gregaria, sessilis, primitus subglobosa, mox explanata, sicca vario modo complicata et urceolata, carneo-vitellina, extus pallidior, albo-scabra, humida dilatata, plana, circ. 1 mm. diam., hyalino-prosenchymatico-contexta; ascis clavatis, apice incrassatis, 8-sporis, 90 \approx 9; sporidiis oblongis, obtusis, hyalinis, 1, demum distincte 2-cellularibus, rectis, distichis, 10-12 \approx 5; paraphysibus filiformibus, subdichotomis, apice in clavam irregularem, hyalinam, c. 3 μ . crassam exeuntibus.

Hab. ad caules putridos Aconiti Napelli et Cirsii spinosiss. prope glaciem æternam, Alpeiner-Gletscher in valle Stubai et Taschach-Gletscher in valle Piz in alpibus tyrolensibus (DR. REHM).

II. *Helotinia*. Sacc. Ascomata glabra.

4. **Helotiella Vaccinii** (Rehm) Sacc. *Helotium Vaccinii* Rehm Ascom. 1974

Alp. n. 8 — Gregaria orbicularis, primitus subglobosa, concava, extus fusca, humida disco explanato concaviusculo, una cum margine hyalino, circ. 1 mm. diam., sessilis, hyphis longis, fuscis, septatis, circ. 3 μ . crassis, simplicibus basi circumdata, prosenchymatico-contexta; ascis clavatis, apice incrassatis jodoque cœrule-scentibus, 8-sporis, 45 \approx 8; sporidiis oblongo-ellipticis, rectis vel subcurvatis, hyalinis, 1-2-cellularibus, distichis, 14 \approx 3; paraphysi-bus ascos superantibus, filiformibus, apice sensim 4 μ . cr. hyalinis.

Hab. ad ramum siccum Vaccinii uliginosi, Peischlkopf in alpibus tyrolensibus (BRITZELMAYR).

5. **Helotiella citrinella** (Rehm) Sacc. *Niptera citrinella* Rehm Ascom. 1975

n. 262. — Gregaria, in cortice interiore sessilis et per rimas epideridis fissæ protuberans, patellæformis, primitus orbicularis, dein irregularis et confluens, plana, margine tenui concolori cincta, fla-va vel citrinella vel æruginosa, circ. 1 mm. diam., contextu par-enchymatico flavo; ascis clavatis, apice incrassatis, 8-sporis, 90 \approx 9, apice jodo cœrulescente; sporidiis fusoideo-elongatis, acutatis, plerumque subcurvatis, 2-cellularibus, hyalinis, distichis, 21-24 \approx 3;

paraphysibus filiformibus, haud clavatis, ascos superantibus, guttulatis, 3-4 μ . cr. hyalinis.

Hab. ad ramos putridos pendulos *Alni viridis* in alpibus editoribus tyrolensisibus.

6. **Helotiella Russellii** (B. et C.) Sacc. *Peziza Russellii* B. et C. North 1976
Am. Fung. n. 739, Cooke Disc. U. S. I, p. 296. — Erumpens, sessilis, fasciculata, lateritia; margine obtuso; disco leviter concavo; ascis clavatis; sporidiis biseriatis, oblongis, utrinque angustatis, demum uniseptatis, 15 μ . longis.

Hab. in corticibus New England Amer. bor. (RUSSELL). — Primo obtutu *Nectriam* collapsam refert. Hymenium in sicco pulvulentum.

7. **Helotiella Mercurialis** (Boud.) Sacc. *Pseudopeziza Mercurialis* 1977
Boud. Bull. Soc. Myc. IV, pag. 84, tab. 17, f. 6. — Minutissima, erumpens, sparsa aut subcæspitosa, luteo-ochracea, 0,2-0,4 mm. lata, lenticularis aut hemisphærica, margine tenuissimo fulvo, sessilis, intus et extus concolor, regularis, margine fulvescente; paraphysibus gracilibus dichotomo-ramosis, non aut vix septatis, ramulorum apicibus non incrassatis 1,5 μ . rarius 2 μ . cr.; ascis octosporis, clavatis, 60-65 \times 10-11, vacuis foramine vix conspicuo; sporidiis hyalinis, oblongo-fusiformibus subcurvatis aut flexuosis, primo continuis denique medio septatis et subconstrictis, intus paululum granulosis præcipue ad extremitates, 11-15 \times 4-5.

Hab. ad caules exsiccatos *Mercurialis perennis* in umbrosis Montmorency Galliæ.

8. **Helotiella drosodes** (Rehm) Sacc. *Helot. drosodes* Rehm Ascom. 1978
n. 566. — Gregaria, orbicularis, hemisphærica vel subcyathoidea, sicca fuscescens vel flavidula, urceolata, humectata subexplanata, marginata, fere hyalina, 1 (1-2) mm. diam., inter *Torulam* aliquam nidulans, prosenchymatico-flavidulo-contexta; ascis clavatis, apice incrassatis, 8-sporis, 60 \times 8 jodoque violascentibus; sporidiis ellipticis, subacutis, rectis vel navicularibus, hyalinis, 2-cellularibus, 10-13,3; paraphysibus filiformibus, apice subincrassatis.

Hab. ad basim cauli putridorum *Adenostyli* dis alpinæ, Bauernalpkopf alpium Allgoviæ (BRITZELMAYR).

9. **Helotiella erythrostigma** (Rehm) Sacc. *Mollisia erythrostigma* 1979
Rehm in Voss Mat. Krains V, p. 225, fig. 6, a, b. — Sparsa vel gregaria in pagina superiore foliorum dealbatorum, contextu parenchymatico-hyalino vel rubidulo, primitus globosa, clausa, dein patellaris, tenerrime marginata, glabra, humida hyalino-flavidula

vel rubidula, sicca demum cinnabarino-rubra, 0,15-0,35 mm. diam.; ascis clavatis, 90-10-12; sporidiis ellipticis vel subfusoideis, primus 1-cellularibus, nucleolis 2 instructis, dein 2-cellularibus, rectis hyalinis, 12-17-6, distichis; paraphysibus filiformibus 1,5-2 μ. cr. superne dichotomis, hyalinis.

Hab. in foliis languidis anni præteriti Cerastii alpini in ober. Bärentale Carnioliae. — *Calloriæ Primulæ* affinis dicitur.

10. **Helotiella herpotricha** (Berk.) Sacc. *Peziza herpotricha* Berk. 1980 Dec. n. 304, t. I, f. 2. — Cupulis convexis pallide rufis dein obscurioribus, glabris, immarginatis, hyphis ramosis repentibus insidentibus; ascis brevibus clavatis; sporidiis oblongis curvulis 1-septatis saepius biguttulatis, polystichis.

Hab. in foliis viventibus, saepius epiphylla, Caripi in Brasilia. — Ramuli hypharum breves clavati basi uniseptati; asci ex icon. 8-spori.

SOLENOPEZIA Sacc. (Etym. *Solenia* et *Peziza*). — Ascomata sessilia, minuta, urceolata, urceolato-oblonga, rarius hemisphærica, cerea, extus villosa vel pilosa, brunnea vel lacticoloria. Ascii typice octospori, paraphysati. Sporidia ovoidea, oblonga vel subfusoidea, 1-septata, hyalina.

1. **Solenopezia Solenia** (Peck) Sacc. *Peziza Solenia* Peck 25 Rep. 1981 p. 99. — Ascomatibus minutis, fere cylindraceis, pilosis, brunneis, margine contracto apertis; sporidiis oblongis, stipatis vel distichis, uniseptatis, saepe 4-nucleatis, subhyalinis, 12 μ. longis, paraphysibus filiformibus.

Hab. in caulis emortuis Eupatorii ageratoidis, Watkins Glen Amer. bor. — Ascomatibus elongatis *Soleniam* in mentem revocat. Teste auctore, a *P. Eupatorii* Schw., ad quam quidam mycologus retulit, satis diversa.

2. **Solenopezia trabinelloides** (Rehm) Sacc. *Helotium trabinelloides* 1982 Rehm Ascom. n. 853. — Gregaria vel conferta, sessilis, primus globosa, clausa, dein urceolata, rubro-aurantiaca, margine integro cincta, 0,2-1,5 mm. diam., extus puberula hyphis fasciculatis, subobtusis, flavidis, septatis, asperulis, creberrimis, 60-70-5, e pariete externa excipuli parenchymatici abeuntibus; ascis clavatis, subcurvatis, apice obtusis, 8-sporis, 90-7-9; sporidiis oblongis vel oblongo-fusiformibus, obtusis, rectis vel subcurvatis, hyalinis, 2-cellularibus, 10-17-4, distichis; paraphysibus filiformibus,

1 μ . cr., apice rotundatis et sæpe subcurvatis, 2 μ . cr. plasmate oleoso rubro impletis.

Hab. ad truncum decorticatum fagineum prope balnea Herculis Hungariæ (LOJKA). — Jodii ope ascoma paraphysesque, imprimis porus ascorum, cœruleo tinguntur.

3. **Solenopezia vulpina** (Cooke) Sacc. *Pez.* Cooke Hedw. 1875, p. 82. 1983

— Gregaria, sessilis, punctiformis, vulpino-rubra, sicca testacea; cupula subglobosa, plano-depressa, primum leniter tomentosa, demum subglabra 0,5 mm. lata, margine tumido; ascis cylindraceis; sporidiis ellipticis, minutis, binucleatis, deinde uniseptatis, hyalinis.

Hab. ad lignum cariosum, New Jersey (ELLIS), Buffalo (CLINTON).

4. **Solenopezia Britzelmayriana** (Rehm) Sacc. *Trichopeziza Britzel-* 1984
mayriana Rehm Ascom. Alp. n. 10. — Gregaria sessilis, globo-

sa, primitus subclausa, dein urceolata, extus cervino-fuscidula, disco pallido, demum subdilatato, 1 mm. diam., strigosa, prosenchy-matico-contexta; pilis simplicibus, longissimis, septatis, rectis, p. p. spiraliter contortis, obtusis, dilute fuscis, circ. 3–5 μ . crassis; ascis clavatis, 8-sporis, 90 \approx 6, apice jodo cœrulecente; sporidiis cylindricis, obtusis, rectis, hyalinis, 2-cellularibus, distichis, 10–11 \approx 2; paraphysibus ascos superantibus, hyalinis, basi circ. 5 μ . diam., apice acutatis.

Hab. ad crates prope ripam fluvii juxta Augustam-Vindelico-
rum seu Augsburg (BRITZELMAYR).

5. **Solenopezia Dorcas** (B. et Br.) Sacc. *Peziza Dorcas* B. et Br. 1985

Fung. of Ceyl. n. 944. — Sessilis, sparsa, globosa, aperta, extus cervina tomentosa; hymenio fusco; sporidiis subellipticis, medio constrictis, biseriatis uniseptatis binucleatis, 12 μ . longis.

Hab. in plantis quibusdam monocotyledoneis in ins. Ceylon. — Primitus Sphæriaceam villosam imitatur.

6. **Solenopezia consimilis** (Oud. et Rehm) Sacc. *Lachnum consimile* 1986

Oud. et Rehm in Oud. Contr. mycol. XII, p. 25. — Ascomatibus sparsis vel approximatis, sessilibus, $2/3$ mm. diam., hemisphaericis, solidis, undique pilis articulatis vel plus minusve sinuositis, basi brunneis, sursum pallidioribus et apice hyalinis ornatis; hymenio leniter brunneolo; ascis cylindraceis vel clavatis, 25 \approx 4; sporidiis monostichis, cylindraceis, curvis, utrinque obtusis, 1-septatis, loculis inæqualibus, 12–15 \approx 2; paraphysibus asco longioribus, fu-

siformibus.

Hab. in ligno putri in Nederlandia (DESTREE).

7. **Solenopezia columbina** (Kalch. et C.) Sacc. *Peziza columbina* K. 1987

et C. Grevillea X, p. 144. — Sessilis, pallide griseo-fusca; cupulis hemisphæricis, pilis acutis, hyalinis, mollibus, dense obsitis; disco obscuriore, $\frac{1}{2}$ mm. diam., margine incurvo; ascis clavato-cylindricis; sporidiis lanceolatis, distincte uniseptatis, hyalinis, $14 \times 3,5$; paraphysibus filiformibus.

Hab. ad cortices in Africa australi (MAC-OWAN).

BISPORELLA Sacc. Consp. gen. Disc. pag. 6 (1884) *Bispora* Fuck. Symb. p. 310 nec Corda (Etym. a *Bispora*, cuius videtur status ascophorus). — Ascomata initio subclavata dein cupulata, stipitata, ceracea, glabra, subiculo conidiophoro *Bisporæ monilioidis* insidentia. Ascii octospori, paraphysati. Sporidia oblonga, 1-septata, hyalina.

1. **Bispora monilifera** (Fuck.) Sacc. Consp. gen. Disc. p. 6, *Bispora* 1988 Fuck. Symb. pag. 310, t. IV, fig. 54, *Hymenoscypha* Phill. Disc. p. 130. — Cupulis 1-3 mm. altis, 1-2 mm. lat. in fungi conidiophori mycelio ortis, primo stipitiformibus, subclavatis, plerumque curvatis, demum antice magis magisque dilatatis, disco concavo, maturis patellæformibus v. planis, marginatis, ceraceis, longe firmeque stipitatis, glabris, pallide fuscis, disco parum obscuriori, sparsis subcæspitosisque, quandoque basi connatis; ascis oblongis elongatisve, 8-sporis, $64-68 \times 8$; sporidiis primo distichis, demum in asco valde elongato submonostichis, oblongis, utrinque parum attenuatis, sed obtusis, distincte uniseptatis, 12×4 , loculis uniguttulatis, hyalinis; paraphysibus filiformibus, simplicibus subclavatis.

Hab. in ligno putrescente Fagi et Carpi in Germania et Britannia. — Fungus conidiophorus sistit *Bisporam monilioidem* Corda: mycelium primo in ligno mucido, maculas griseas sublimitatas formans, ramosissimum. Conidia demum in stipitibus hyalinis orta, catenulata, singula oblongo-lanceolata, utrinque obtusa, uniseptata, primo hyalina, demum fusca opaca.

LANZIA Sacc. Consp. gen. Disc. p. 6 [1884] (Etym. ab egregio mycologo et diatomologo doct. M. LANZI, romano). — Ascomata stipitata, patellata, ceracea, disco plano v. concavo, stipite plerumque longitrorsum plicatulo. Ascii elongati, octospori. Sporidia oblonga, demum 1-septata, hyalina.

1. **Lanzia flavo-rufa** (Sacc.) Consp. gen. Disc. pag. 6, *Helotium* 1989 *flavo-rufum* Sacc. Fung. Ven. ser. IV, n. 99, F. ital. f. 116. — Cupulis ceraceis, gregariis, discoideo-patellatis, $\frac{1}{2}-\frac{3}{4}$ mm. diam..

subtus convexulis et in stipitem subæquilongum productis, glabris, intus albis, disco jugiter plano purpureo-ferrugineo, margine initio luteolo non elevato, extus stipiteque longitudinaliter (præcipue senio) plicatis, flavo-cerini; excipuli contextu molli tenuiter areolato, flavescente, cellulis marginalibus elongato-clavatis; ascis cylindraceo-clavatis basi breve attenuatis, apice subtruncatis, 60-65 \approx 5-6, pseudoparaphysibus crassitudine variis obvallatis, 8-sporis; sporidiis oblongis, 8-12 \approx 3 $\frac{1}{2}$ -5, utrinque obtusis, medio tenuissime constrictis ibique (spurie?) 1-septatis, hyalinis.

Hab. in truncis putrescentibus uidis Robiniae Pseudaciæ Patavii Ital. bor. (G. BIZZOZERO). — Cupulæ insident tapeto velutino atro-fuligineo conidiophoro cum *Lanzia* forte metagenetice connexo et *Chalaran Ampullulam* Sacc. Syll. IV, pag. 334, sistente.

2. ***Lanzia rugipes*** (Peck) Sacc. *Helotium rugipes* Peck 26 Rep. p. 82. 1990
— Ascomatibus concavis, dein expansis, saepe minute umbilicatis, extus brunneo-flavidis, disco viridulo-brunneo vel flavidio, stipite brevi, deorsum attenuato, usque ad cupulam plicato-lacunoso; ascis cylindraceis, tenuibus; sporidiis monostichis, uniseptatis, ellipticis, 7-8 \approx 3-4.

Hab. in ligno putri, Worcester Americæ borealis. — Ascoma 4-8 mm. lat.

3. ***Lanzia crassipes*** (Pat.) Sacc. *Peziza crassipes* Pat. Tab. An. Fung. 1991 p. 124, f. 279. — Tenuis, concava, dein plana, 1-2 mm. lat. intus brunneo-rufescens, extus albo-rubida, margine recto, stipite basi incrassato, albo vix rubescente, 4-6 mm. long.; ascis brevibus, deorsum abrupte attenuatis, octosporis; sporidiis distichis, ellipsoideo-fusoideis, 12-14 μ . long., hyalinis, inæqualiter bilocularibus; paraphysibus acicularibus, rubello-guttulatis.

Hab. ad terram in quercetis, Polygny Jura.

NIPTERA Fr. Summa Veg. pag. 359 (1849) em. De Not. Prof. Disc. p. 375, Sacc. Consp. gen. Disc. pag. 6, nec Fuck. (Etym. *nipter* malluvium, vas). — Ascomata cupulato-patellata, sessilia v. subsessilia, superficialia, minuta, ceracea, excipulo cellulis arctis fuscis contexto. Ascii elongati, octospori, paraphysati. Sporidia oblonga v. fusoidea, 1-septata, hyalina-luteola. — Est *Mollisia didymospora*.

1. ***Niptera Raineri*** (De Not) Comm. Critt. I, pag. 375, *Cenangium* 1992 *Raineri* De Not. Micr. Dec. I, n. 4, f. 4. — Nitens, seriatim gregaria, obconico-subsessilis, ex urceolato cupulata, sicca vix rugulosa, fu-

ligineo-nigra, arescendo subclausa vel rimose dehiscens, $\frac{1}{2}$ mm. diam., disco pallide cinereo; ascis tereti-clavatis, subsessilibus, apice rotundatis, octosporis; sporidiis distichis, fusoideis, utrinque acutis, constricto-1-septatis, ex hyalino fuscellis, 11-12 \approx 3-4; paraphysibus bacillari-clavulatis.

Hab. ad ramulos sicclos Cytisi radiati in montibus Lariensis Ital. bor. (DE RAINER).

2. **Niptera Vossii** (Rehm) Voss Mater. IV, pag. 13, f. 6, a, b, *Mollisia Vossii* Rehm Ascom. Ill. et Exs. n. 710. — Gregaria, emergens, dein lata basi sessilis, orbicularis, patellariformis, plana, c. 1 mm. diam., sicca margine subplicata, extus nigro-fusca, disco livido, fusco-parenchymatica; cellulis marginalibus interdum subelongatis; ascis clavatis, apice obtuse acutatis, 8-sporis, 75 \approx 12; sporidiis clavatis, 2-cellularibus, cellula superiore latiore, medio haud constrictis, rectis, interdum subcurvatis, hyalinis, 2-stichis, 21-25 \approx 6; paraphysibus filiformibus, hyalinis, versus apice sensim incrassatis.

Hab. ad ramulos emortuos Cytisi radiati, Oberkauth juxta Niedertiefenbach prope Gottschee Carnioliae (Prof. Voss).

3. **Niptera Saliceti** (Rehm) Sacc. *Mollisia Saliceti* Rehm Ascom. Alp. 1994 n. 20. — Gregaria, in rimis epidermidis cortici inferiori fuscatae insidens, sicca fusca complicata, humida explanata, patellæformis, concava, demum plana, vix marginata, extus fuscidula, disco cinereo-albido, c. 0,8-1 mm. diam.; ascis clavatis, subcurvatis, 8-sporis, 50 \approx 8; sporidiis obtusis, oblongis, 2-cellularibus, rectis v. subcurvatis, hyalinis, distichis, 10-12 \approx 3-5; paraphysibus filiformibus, apice sensim 3 μ . incrassatis, hyalinis.

Hab. ad ramum siccum Salicis cujusdam apud glaciem æternam. Taschach-Gletscher in valle Piz alpium tyrolens. c. 2000 m. alt. (DR. REHM). — *Habitus Pyrenop. ligni.*

4. **Niptera laricina** (Rehm) Sacc. *Pyrenopeziza laricina* Rehm Asc. 1995 Ill. n. 511. — Sessilis, gregaria, primitus subclausa, dein patellaris, sicca atra, extus rugulosa, varie plicata, saepe hysteriformis, humida extus atra, explanata, disco pallido, parenchymatico-fusco-contexta, circ. 0,5 mm. diam.; ascis clavatis, 8-sporis, 40-50 \approx 9; sporidiis obovatis vel biscoctiformibus, rectis vel subcurvatis, 1-2-cellularibus, distichis, hyalinis, 10-12 \approx 5; paraphysibus filiformibus, superne vix incrassatis.

Hab. ad ramulos sicclos Laricis europææ in monte Arlberg

tyrolensi, c. 1300 m. (BRITZELMAYR), ad ramulos emortuos Abietis in silva Cansilio Ital. bor. (SPEGAZZINI).

5. **Niptera exsiliens** Speg. Mich. I, p. 472. — Sessilis v. brevissime 1996
pedicellata, solitaria v. gregaria, hinc inde per epidermidem ramu-
lorum erumpens, extus atro-cærulea, disco pallidiore griseo, linea
candida marginato, magnitudine valde ludens, $\frac{1}{4}$ -1 mm. diam., con-
textu parenchymatico, marginem versus prosenchymatico, densis-
simo, membranaceo, ascis cylindraceo-subclavatis, apice obtusiu-
sculis, in pedicellum crassum desinentibus, 55-65 \approx 5-7, 8-sporis;
paraphysibus guttulatis, crassiusculis; sporidiis clavulatis medio
uniseptatis, ad septum non vel vix constrictis, hyalinis, 10-15 \approx 2-3.

Hab. in ramulis emortuis dejectis Quercuum, quandoque pa-
rasitica in Colpomate quercino, Conegliano Ital. bor.

6. **Niptera Rhododendri** (Ces.) De Not. Comm. Critt. I, pag. 375. — 1997
Sparsa, sessilis, cupulata, arescendo hysterioideo-subclausa, coria-
ceo-ceracea, fuligineo-atra, $\frac{3}{4}$ -1 mm. lat., glabra; ascis clavatis,
75 \approx 12; sporidiis oblongis, obtusis, rarius ovoideis, rectis curvu-
lisve medio constricto-1-septatis, 16-18 \approx 6, distichis, hyalinis;
paraphysibus filiformibus, apice clava ovoidea brunnea, 6 μ . crass.
terminatis.

Hab. in capsulis siccis (et in ramulis?) Rhododendri fer-
ruginei in Italia boreali et Tyroli.

7. **Niptera Rollandii** Boud. Bull. Soc. Myc. IV, p. 83, t. 17, f. 5. — 1998
Minutissima, sessilis, hemisphærica, alba, 0,2-0,4 mm. lata, crassa
levis v. vix pruinosa, alba, hymenio concolore plano dein convexo,
marginе crasso vix superante cincto; paraphysibus gracilibus, sim-
plicibus aut ad basim tantum divisis, continuis, sed ad apicem
clava oblonga, tri-septata, 15 \approx 4-5 terminatis; ascis elongato-su-
soideis, octosporis, 90-100 \approx 8-9; sporidiis oblongo-fusiformibus,
hyalinis, quadri-guttulatis, guttulis oleosis, sæpe divisis et tunc
plus minusve granulosis, primo continuis, denique medio septatis
et paullulum constrictis, 13-15 \approx 4.

Hab. ad cortices Alni glutinosæ prope Isle-Adam Galliæ
(ROLLAND). — An *Helotiella*?

8. **Niptera subbiatorina** Rehm Ascom. n. 858. — Sparsa, sessilis, pri- 1999
mitus globosa, dein patellaris, ceracea, extus flavidula, glabra, con-
textu parenchymatico, flavidulo, versus marginem album hyalina,
disco hyalino-flavidulo, subplano, 0,1-0,3 mm. diam.; ascis cylin-
draceis, apice rotundatis, 8-sporis, 60-66 \approx 5-6; sporidiis oblongis
vel oblongo-ellipticis, obtusis, rectis, hyalinis, 2-cellularibus, 7-9

≈ 3, distichis; paraphysibus vix 1 μ. cr., apice subramosis, dilute flavidulis et conglutinatis.

Hab. ad truncum decorticatum *Castaneæ vescæ*, Gratsch prope Meran Tyrol (Dr. EGGERTH). — *Biatorinæ* subaffinis sed gonidia nulla.

9. **Niptera castanea** (Sacc.) *Pyrenopeziza? castanea* Sacc. Mich. I, 2000 p. 422, F. ital. f. 1400. — Ascomatibus gregariis superficialibus, sessilibus, $\frac{1}{3}$ - $\frac{1}{2}$ mm. diam., concavis, obtuse marginatis siccis non contractis, hyphis repentibus basi cinctis, excipuli contextu parenchymatico ad marginem prosenchymatico; ascis cylindraceo-clavatis, breve stipitatis, 70-80 ≈ 9-10, apice rotundatis; paraphysibus filiformibus obvallatis, octosporis; sporidiis distichis fusoideis rectis curvulisve, 15 ≈ 4 (spurie?) 1-septatis, non constrictis, hyalinis.

Hab. in ligno decorticato putrescente *Castaneæ vescæ* in Euganeis ad Bressèo Ital. bor.

10. **Niptera hypogæa** (Bres.) Sacc. *Mollisia hypogæa* Bres. Trid. 2001 p. 69 et 105, t. 75, f. 1. — Sessilis, carnosò-ceracea, explanato-concava, extus nigra, margine albo, puberulo-tomentoso, ut plurimum undulato-repando, disco e fuscidulo vel cinereo subolivaceo albido, 1- $\frac{1}{2}$ mm. lata; substantia subconcolore, inodora; ascis fusoideis, jodo haud tinctis, 70-100 ≈ 10-14; paraphysibus numerosis, asco longioribus, apice subincrassatis v. subacutatis, ramosis, 2 μ. latis; sporidiis hyalinis fusoideis, pluriguttulatis, demum 1-septatis, 15-25 ≈ 3-4, distichis v. congregatis.

Hab. subterranea in rhizomatibus adhuc vivis *Adenostyliidis alpinæ* var. *albifrontis*, quæ nigrificat, juxta rivulos subalpinos Valle di Sole, Trento.

11. **Niptera umbrinella** (Desm.) Sacc. *Peziza umbrinella* Desm. 10, 2002 Not. p. 36. — Sessilis, erumpens, ceraceo-mollis, sparsa, minuta, orbicularis, glabra, umbrino-pallens, plano-convexa, sicca concava, margine acuto, brunneo, integerrimo, vix prominente; ascis clavatis; sporidiis hyalinis, uniseptatis, oblongis, subfusiformibus, 12-14 μ. long.; paraphysibus capitellatis.

Hab. ad caules *Solidaginis* in Gallia bor. — Ascoma $\frac{2}{3}$ mm. lat.

12. **Niptera leucostoma** (Karst.) Sacc. *Mollisia leucostoma* Karst. M. 2003 F. I, p. 203. — Conferta, innato-erumpens, hemisphærica v. multa pressione difformis, cinereo-pallida, margine fimbriato, albicante vel subalbo, sicca subsphæroideo-depressa, nigricans v. fuliginosa, ore contracto sæpeque angulato flexo, 0,3-0,5 mm. lat.; ascis fu-

soideo-cylindraceis, 40-50 \approx 5,5-6, 4-sporis, obturaculo minutissimo vix vel obsoletissime cœrulescente; sporidiis aciculari- raro fusoideo-bacillaribus, rectis vel subrectis, vulgo uniseptatis, 12-27 \approx 2-3; paraphysibus gracilescentibus apicem versus incrassatis.

Hab. ad caules Artemisiæ vulgaris exsiccatos in Fennia australi.

13. **Niptera Coriariæ** (Pass.) Sacc. *Peziza coriariæ* Pass. Diagn. 2004
nuov. funghi n. 16. — Sparsa v. gregaria, sessilis, firma, madore discoideo-applanata, disco subconcavo, cinereo-violacea, in statu sicco contracta, difformis, fusco-nigrescens; ascis breviusculis clavatis, stipitatis, 8-sporis; sporidiis distichis, tereti-oblongis, tenuibus, hyalinis, endoplasmate granuloso, medio tandem tenuiter diviso; paraphysibus crassiusculis, subclavatis, intus granulosis, ascos æquantibus.

Hab. in caulis siccis decorticatis Coriariæ myrtifoliæ, Rocca Prebalza pr. Berceto in Appennino parmensi. — An *Patellea*?

14. **Niptera lacustris** (Fr.) Summa p. 359, *Pez. lacustris* Fr. Syst. 2005
Myc. II, p. 143. — Sessilis, applanata, orbicularis, ceraceo-mollis, plano-convexa, e fuligineo-nigricans; ascis clavatis; sporidiis oblongo-ellipsoideis, 12-15 μ . leng. (Curr.), v. 22-30 \approx 6-9, 1-3-septatis (Nyl.); paraphysibus filiformi-clavulatis.

Hab. ad calamos emortuos Scirpi lacustris (et Alismatis?) in Suecia, Britannia, Germania, Belgio. — Junior adnata, plano-convexa, $\frac{1}{2}$ -1 mm. lata, levigata, glabra, atra, dein dilatata, applanata, tamen subtus libera et unico puncto centrali tantum fixa, extus fuligineo-nigricans, sicca tota atra. Margo primo concavus, dein apertus, inflexus, non omnino integer; demum vero omnino regularis, integer, parum prominens. Discus turgidus, mollis, humidus fuligineo-pallidus, siccus nigricans. «An *Belonii* species?».

15. **Niptera caricinella** Peck 30 Rep. p. 61, t. I, fig. 5-8. — Ascomate minuto, 0,5-0,7 mm. lat., sessili, rubescente vel ochraceo-brunneo (udo), nigro v. nigricante (sicco), disco piano v. subconcavo; ascis clavatis; sporidiis oblongis, confertis, obtusis, uniseptatis, hyalinis, 20-25 μ . longis.

Hab. in foliis Caricis utriculatae, Adirondack Amer. bor. 2007

16. **Niptera phæa** (Rehm) Sacc. *Mollisia phæa* Rehm Ascom. Ill. et Exs. n. 713. — Subgregaria, sessilis, hemisphærico-urceolata vel patellaris, vix v. tenuiter obscurius marginata, plana, extus fusca,

disco pallidiore, c. 0,3-5 mm. diam., contextu parenchymatico fuscidulo; ascis clavatis, subcurvatis, apice obtuse acutatis, 8-sporis, apice jodo cœrulecente, 70-80 = 10-12; sporidiis oblongo-ellipticis, obtusis, hyalinis, primitus 2-cellularibus, medio haud constrictis, 15-18 = 5-6; paraphysibus teneris, dichotomis, apice 5 μ . cr., clavatis et dilute flavidis, itaque ascos superantibus et discum (ope kalii caust. adhibiti solutum et decoloratum) formantibus.

Hab. ad calamos Junci siccos in silva prope Königstein an der Elbe Saxoniæ.

17. **Niptera melatephroides** (Rehm) Sacc. *Mollisia melatephroides* 2008 Rehm Ascom. n. 655. — Gregaria, sessilis, patellariformis, sicca suburceolata, humida plana, irregulariter marginata, scabra, extus nigro-fusca, disco pallide luteolo, c. 1 mm. diam., contextu parenchymatico, fusco; ascis clavatis, apice obtuse acutatis, subcurvatis, 8-sporis, 120 = 18, apice jodo dilute cœrulecente; sporidiis ellipticis, subobtusis, rectis, 2-cellularibus v. bipartitis, quaque cellula binucleata, hyalinis, 18-21 = 4, episporio gelatinoso c. 6 μ . crasso circumdatis; paraphysibus dichotomis, apice clava plus minusve rotunda, c. 6 μ . crassa, hyalina instructis.

Hab. ad culmos siccos Moliniæ cœruleæ in silvis prope Pulsnitz Saxoniæ (STARITZ), — Differt a *Mollisia melatephra* in primis sporidiis 1-septatis, episporio mucoso crasso, paraphysibus dichotomis, apice clavatis.

18. **Niptera Punctum** (Rehm) Sacc. *Micropeziza Punctum* Rehm 2009 Ascom. n. 261. — Minutissima, primo punctiformis, dein orbiculariter basi late adnata, sessilis, subhemisphærica, concaviuscula, margine elevato, disco humido fuscidulo, sicca atra, contextu parenchymatico, fusco, margine prosenchymatico viridi-fusco, sparsa, c. 0,3 mm. diam.; ascis clavatis, subcurvatis, 8-sporis, 55 = 12; sporidiis oblongis, obtusis, interdum elongato-dacryoideis, rectis, rarius subcurvatis, hyalinis, 2-cellularibus, utraqua cellula binucleata, distichis, 15-17 = 4,5; paraphysibus filiformibus, apice subincrassatis et fuscidulis, saepe subincurvatis.

Hab. ad folia sicca Nardi strictæ pr. Teplitz in Bohemia (THÜMEN).

19. **Niptera antarctica** Speg. Fung. Fueg. n. 342. — Cupulis sparsis 2010 v. hinc inde laxe gregariis, primo epidermide tectis, dein, ea sedente, denudato-superficialibus, hemisphærico-cupulatis, 200-250 μ . diam., atris, glabris; contextu margine prosenchymatico (an fimbriatulo?) ceterum parenchymatico olivaceo-fusco; ascis cylind-

draceis v. subfusoideis, antice truncato-rotundatis, crassiuscule tunicatis minutissime bifoveolatis, postice brevissime crasseque attenuato-pedicellatis, 65-80 \pm 5, octosporis; paraphysibus paucis guttulatis obvallatis; jodi ope obturaculo ascorum intense cœruleo-scente; sporidiis recte distichis, cylindraceo-fusoideis, utrinque obtusiusculis, 17-18 \pm 1,5-2, medio 1-septatis non constrictis, hyalinis.

Hab. ad culmos emortuos Rostkoviæ grandifloræ in pratis uliginosis in Staten island Fuegiæ. — Species vere pusilla ascomatibus quandoque matrice fere immersis nonnihil ad genus *Fabræa* vergens.

20. **Niptera lichenicola** (Speg.) Sacc. *Pyrenopeziza lichenicola* Speg. 2011 Mich. I, p. 472. — Ascomatibus laxe gregariis, minutissimis, vix $\frac{1}{10}$ - $\frac{1}{8}$ mm. diam., atris, disco dilutiore, siccis subsphæroideis, humidis patellaribus, contextu excipuli densissime parenchymatico atro; ascis clavato-ellipticis, crasse tunicatis, a paraphysatis (?), octosporis, jodi ope apice intense cœruleoscentibus, 30-35 \pm 12-15; sporidiis didymis, primo muco obvolutis, dein liberis, loculo superiore crassiore, hyalinis, ad septum constrictis, 6-8 \pm 3-4.

Hab. in thallo emortuo *Parmeliæ caperatae* in Monte Cerva prope Belluno Ital. bor.

SCUTULA Tul. Mem. Lich. p. 118 (1844) nec Lour. (Etym. *scutum*).

— Ascomata plano-scutellata, sessilia, ceracea, excipulo distincto prædicta, tandem immarginata, deformia. Ascii octospori, paraphysibus subconglutinatis obvallati. Sporidia ovato-oblonga, 1-septata, hyalina. — Genus hinc Lichenibus, illinc Dermateis affine, habitatione in Lichenibus parasitica peculiare.

1. **Scutula Stereocaulorum** (Anzi) Körb. Parerg. p. 455, Karst. Revis. 2012 p. 151, *Lecidea* Anzi Menip. p. 26, *Biatorina* Th. Fr. Lich. Arct. p. 188. — Congesta, adpressa, primitus concava et marginata, dein planiuscula, immarginata, regularis, nigricans vel obscure carneola vel pallens intus obscura, 0,3-0,4 mm. lata; ascis clavatis, crasse tunicatis, 50 \pm 10, jodo intense cœruleoscentibus, octosporis; sporidiis distichis, ovoideo-oblongis, uniseptatis, biguttatis, hyalinis, 13-19 \pm 4-6; paraphysibus filiformibus, apice clava luteo-fusca 6 μ . crass. terminatis.

Hab. in *Stereocauli* speciebus in Italia boreali, Helvetia, Fennia, Germania. — Hypothecium flavidum.

2. **Scutula miliaris** (Wallr.) Karst. Rev. mon. p. 149, *Peziza* Wallr. 2013

Crypt. p. 499, *Scutula Wallrothii* Tul. Mem. Lich. p. 119, t. 14, f. 14, *Biatorina miliaris* Th. Fr. *Lecidea epigena* Nyl. in Flora 1865, p. 5, *Calloria Peltigeræ* Fuck. Symb. p. 283. — Conferta, plana vel concaviuscula et marginata, dein convexa immarginata, primo pallida v. testaceo-pallida, mox denigrata, 0,3-0,4 mm. diam.; ascis clavatis, octosporis; sporidiis distichis, ovoideis vel ellipsoideis, uniseptatis, hyalinis, 11-15 = 5-7; paraphysibus coalescentibus.

Hab. in thallo Peltigeræ caninæ et Peltigeræ rufescens in Lapponia, Germania, Helvetia, Italia, Gallia. — Gelatina hymenea jodo dilute cœrulescens, dein vinose rubens. —

***S. Heerii** (Hepp.) Karst. l. c. *Lecidea Heerii* Hepp. in Nyl. Lich. Lapp. 152, *Scutula epigena* Rehm Ascom. n. 20: sporidiis minoribus, nempe 8-12 = 4-5. In thallo Peltigeræ rufescens, P. aphthosæ, P. rufescens, Nephromii tomentosi in Lapponia, Fennia, Germania, Helvetia.

3. **Scutula solorinaria** (Nyl.) Karst. Rev. Mon. p. 150, *Lecidea* Nyl. 2014 Lich. Delph. p. 402. — Affinis *Scutulae oxysporæ* sed ascomatibus convexis, sporidiis minoribus, 13-14 = 3,5-4,5, gelatina hymeniali jodo vinoso-rubente.

Hab. in thallo Solorinæ bisporæ in Fennia. — Var. **solorinicola** (Weinm.) Karst. l. c., *Lecidea* Wainio Add. Lich. Lapp. II, p. 30: ascis clavatis 14 μ. crassis, sporidiis 9-12 = 3-3,5, gelatina jodo vinoso-rubente vel sublutescente. In thallo Solorinæ saccatae pr. Kongäs ad mare glaciale.

4. **Scutula Krempelhuberi** Körb. Parerg. p. 455. — Ascomatibus sub. 2015 minutis, sessilibus, patellaribus, jam primitus atris, disco plano dein convexiusculo marginem tenuem concolorem aequante, tandem superante; ascis anguste clavatis, octosporis; sporidiis minutis ellipsoideis e monoblasto typice biblastis, diametro $2\frac{1}{2}$ -3-plo longioribus, hyalinis.

Hab. in thallo Solorinæ saccatae in Berechtsgaden (KREMPELHUBER). — Hymenium sub microscopio cerasino-brunneolum.

5. **Scutula oxyspora** (Tul.) Karst. Revis. Mon. p. 150, *Abrothallus* 2016 Tul. Mem. Lich. pag. 116, t. 16, f. 27, *Lecidea* Nyl. *Nesolechia* Mass. — Plana v. convexiuscula, nigra v. fusco-nigra, intus fuscescens v. obscura, 0,2 mm. diam.; sporidiis ellipsoideo-fusiformibus, uniseptatis, hyalinis, 14-20 = 5-7; paraphysibus haud discretis; gelatina hymeniali, præsertim ascorum, jodo cœruleacente.

Hab. in thallo Platysmatis, Parmeliæ, Cetrariæ, Imbricariæ in Gallia, Italia, Fennia, Britannia, Suecia.

Sect. 4. Phragmosporæ Sacc.

MASSEEA Sacc. (Etym. a cl. mycologo britannico G. MASSEE). — Ascomata subcarnosa, scutellaria, sessilia, quasi *Humariæ*. Ascii tereti-clavati, paraphysati, octospori. Sporidia fusoidea, 3-4-septata, hyalina.

1. **Masseea quisquiliarum** (B. et C.) Sacc. *P. quisquiliarum* B. et C. 2017 Cub. Fungi n. 670, Cooke Mycogr. f. 37. — Lutea, crateriformis, sessilis, margine inflexo; ascis tereti-clavatis, filiformi-paraphysatis; sporidiis fusiformibus curvatis 3-4-septatis, 25 \approx 6-7, hyalinis, levibus.

Hab. ad ramos emortuos in ins. Cuba. — Ascoma 2-5 mm. lat.

VELUTARIA Fuck. Symb. mycol. p. 300 [1869] (Etym. *vellus* v. *velum*). — Ascomata media magnitudine, ceracea, sessilia, aperta, extus pulveraceo-tomentosa, disco colorato, concavo. Ascii stipitati, elongato-oblongi, 8-spori. Sporidia ovata oblongave, 2-pluriseptata, hyalina. Paraphyses filiformes.

1. **Velutaria griseo-vitellina** Fuck. Symb. mycol. p. 300. — Cupulis 2018 gregariis sparsisve, sessilibus, carnosoceraceis, difformibus, usque ad 2 mm. latis, extus margineque elevato, repando, pulveraceo-tomentosis, griseis, disco plano concavove, vitellino, marginem versus dilute æruginoso; ascis stipitatis, oblongis, 8-sporis, 100 \approx 10; sporidiis distichis, oblongis, utrinque parum attenuatis, rectis curvatisve, triseptatis, hyalinis, 24 \approx 5.

Hab. in sarmenis humidis, putridis Rubi fruticosi prope Estrich Rhenogoviæ.

2. **Velutaria atro-citrina** (B. et Br.) Sacc. *Pez. atro-citrina* B. et 2019 Br. Fung. of Ceyl. n. 945. — Sessilis, subhemisphærica, extus floculis citrinis granulatis villosa; hymenio atro; sporidiis clavatis septatis, 45-50 μ . long.

Hab. in ligno putri in insula Ceylon.

3. **Velutaria nephrodigena** (Phill.) Sacc. *Peziza nephrodigena* Phill. 2020 in Grev. XV, p. 17. — Gregaria v. sparsa, sessilis, globosa, dein hemisphærica, primo tenuiter pubescens, dein glabra, alba, ceracea, firma, 200-400 μ . diam.; disco concavo, castaneo-brunneo, margine inflexo; ascis clavatis; sporidiis oblongo-fusiformibus, utrinque rotundatis triseptatis ad septa constrictis, 12-15 \approx 4-5; paraphysibus non visis.

Hab. in frondibus subemortuis *Nephrodi i hispidi*, New Zealand (KIRK).

DERMATELLA Karst. M. F. I, p. 209 [1871] (Etym. a *Dermatea*), *Dermateæ* et *Peziculæ* spec. Auct. — Ascomata ceraceo-duriuscula, patellata, erumpenti-superficialia, glabra vel puberula. Ascii elongati, paraphysati, 4-8-spori. Sporidia majuscula ovato-oblonga, 2-pluriseptata ex hyalino flava. — Est *Pezicula* sporidiis septatis, sed species adsunt spurie septatae, hinc ambiguæ *). In *Dermateam* et *Scleroderrin* Dermatearum sensim transit.

I. *Eu-Dermatella*. Sporidia transverse tantum septata.

1. **Dermatella Frangulæ** (Fr.) Karst. M. F. I, pag. 209, *Tympinis* 2021
Frangulæ Fr. Syst. Myc. II, p. 174, *Pezicula* Fuck. Symb. p. 279, t. IV, f. 46, *Dermatea* Tul. Carp. III, p. 161. — Sessilis turbinato-truncata, rotundata, opaca, nigrescens; disco atro, margine obsoleto; ascis abrupte longe stipitatis, oblongis, 4-sporis, $66 \div 11$ (parte sporifer.); sporidiis monostichis, ovatis, utrinque, obtusis, 3-4-septatis loculis uniguttulatis, $16-20 \div 7-8$; paraphysibus filiformi-clavulatis.

*) *Peziculis* genuinis supra, p. 310, descriptis addendæ sunt sequentes:

1. **Pezicula amœna** Tul. Carp. III, p. 184, t. XXI, f. 1-9, *Dermatea* Tul. 2022
Bot. Zeit. 1853, pag. 54. — Erumpens, cæspitosa gregariaque, subiculo oblongo, initio crassiusculo et quidem intus locellato, postea maxime attenuato; cupulis obconicis, disco immarginato-plano, dilute aurantiacis; ascis amplis, $128 \div 13$, octosporis; sporidiis ovato-oblongis, continuis, pellucidis, curvulis, $16-19 \div 6,5$; paraphysibus.....

Hab. in ramis quercinis corticatis circa Parisios Galliæ et in Britannia.

2. **Pezicula Coryli** Tul. Carp. III, p. 183, *Dermatea Coryli* Tul. in Bot. 2023
Zeit. 1853, pag. 54. — Erumpens, cæspitosa, turbinata, exigua, straminea, disco tandem plano vel convexiusculo, immarginato; ascis oblongis, octosporis, $100 \div 16$; sporidiis ovato-cylindraceis, curvulis, continuis, utrinque obtusis, $23-27 \div 8-10$; paraphysibus filiformibus, apice clavula obovata, $16-19 \div 5-6$, terminatis.

Hab. in ramis cæsis Coryli Avellanæ circa Parisios Galliæ. — Ad-sunt stylosporæ obovato-oblongæ ocellatæ et spermatia recta, exilissima.

3. **Pezicula conigena** (Phill.) Sacc. *Dermatea* Phill. Rev. mycol. 1883, n. X, 2024 p. 23. — Sparsa, coriacea vel subcornea, atro-carnea, discoideo-convexa, 0,25-1 mm. diam.; sporidiis ellipsoideis, 2-guttatis, $18-20 \div 5-8$; paraphysi-bus divisis, filiformibus, hyalinis.

Hab. in strobilis Pini in Arduennis — An diversa a *Pezicula nectrioides* Phill. (n. 1301) ?

Hab. in ramis emortuis Rhamni Frangulæ in Suecia, Germania, Belgio, Italia, Britannia. — Quoad pycnidia et spermogonia cfr. l. c. Sec. Fuck. sporidia subinde muriformia fusca.

2. **Dermatella quercina** (Fuck.) Sacc. *Pezicula quercina* Fuck. Symb. 2025 myc. p. 279. — Cupulis in cæspitulos gregarios erumpentibus, fasciculatis, stipitatis, disco emarginato, convexo, 1 mm. lato, primo fusco, pulverulento, pellucido, demum ochraceo; ascis oblongis, 8-sporis; sporidiis oblongis, plerumque rectis, 3-guttulatis, demum 3-septatis, hyalinis, 24-28 \approx 5-8.

Hab. in ramis corticatis siccis Quercus ad Judensand prope Oestrich Rhenogoviæ. Sec. Rehm a *Pezic. cinnamomea* vix differt. — Var. **Alni** Fuck. Symb. App. II, pag. 56, Rehm Ascom. n. 463: ascis 75 \approx 9; sporidiis 18-26 \approx 6-7, 1-3-septatis, hyalinis; cupulis flavidis prosenchymaticis. Ad ramos Alni in Germania. — Var. **Aceris** Rehm Disc. p. 252: cupulis in cæspitulos seriatos copiosioribus (usque 30); disco flavo-brunneo, margine pallidiore. In trunco Aceris Pseudoplatani, Berlin Germaniæ.

3. **Dermatella Pumilionis** (Rehm) Sacc. *Pezicula Pumilionis* Rehm 2026 Alp. n. 28. — Sessilia, plana, rotunda, dein irregulariter explanata, livido-cinerea, margine haud distincto pallidore, minutissima, perparum conspicua, gregaria, contextu prosenchymatico, flavo; ascis clavatis, superne subacuminatis, 8-sporis, 60 \approx 12; sporidiis vermicularibus, fusoideo-filiformibus, 4-6-cellularibus, curvatis, hyalinis, parallelis, 36 \approx 2,5; paraphysibus filiformibus.

Hab. ad conos putrescentes Pini Pumilionis Haspelmoos prope Augusta Vindelicorum (BRITZELMAYR). — «An potius *Belonidium*?»

4. **Dermatella livida** (B. et Br.) Sacc. *Patellaria livida* B. et Br. 2027 Ann. N. H. n. 775, *Dermatea* (B. et Br.) Phill. Disc. pag. 340, *Pezicula livida* B. et Br. Rehm Ascom. n. 462. — Minuta, gregaria, sæpe conferta, subhemisphærica, sessilis, olivaceo-flavida; disco plano, margine albido, minute squamuloso; ascis subfusoides v. clavatis, 8-sporis; sporidiis oblongo-ellipsoideis, 3-4-guttatis, dein 1-3-pseudoseptatis, 22-25 \approx 5.

Hab. in truncis pineis emortuis, Leicestershire Britanniæ. — Ascoma $\frac{1}{2}$ -1 mm. lat. in sicco griseo-albidum. — ***D. pachyspora**: cfr. Sacc. F. Gall. n. 2223: ascis tereti-clavatis, breve stipitatis, paraphysatis, octosporis, 130 \approx 20; sporidiis oblongis, utrinque obtusiusculis, 30 \approx 10, 4-nucleatis, tandem tenuiter 3-septatis, ad septa non, nisi medium, constrictis. In cortice Pini in Arduennis

(LIBERT). — Var. **tetraspora** Rehm Ascom. n. 462: sporidiis oblongo-ellipticis, subcurvatis, 1-pseudo 2-cellularibus, 2-4 nucleis magnis instructis, hyalinis, 28-30 = 7, quaternis biseriatis in ascis clavatis, apice incrassatis jodoque violascentibus, subcurvatis, 90 = 14; paraphysibus teneris, superne furcatis. Ad ramum putridum, adustum pineum prope Augustam Vindelicorum (BRITZELMAYR).

5. **Dermatella rosella** (Rehm) Sacc. *Dermatea* Rehm Disc. p. 257. 2028

— Gregaria, erumpens, globosa clausa, dein applanata, patellaris, fere immarginata, convexula, disco rosea, extus brunnea, sicca pallidius marginata, 1-2 mm., coriacea; ascis clavatis, supra obtuse acutatis, 75-90 = 7-9, 8-sporis; sporidiis fusoideis, rectis curvulisve, continuis, dein 1-septatis, hyalinis, 18-25 = 2,5-3, distichis; paraphysibus filiformibus 2,5 μ . cr., septatis, sursum 4 μ . crass., flavidis.

Hab. in ramis quercinis putrescentibus prope Sugenheim in Franconia. — Typus aberrans.

6. **Dermatella eucrita** (Karst.) Sacc. *Pezicula eucrita* Karst. M. F. 2029

I, p. 166, *Dermatea abietina* Auersw. Tauschv. 1865. — Solitaria vel saepius caespitosa, brevissime obconico-stipitata, pallide ochracea v. sordide ochraceo-pallescens, cupula planiuscula 0,4-1 mm. lat.; ascis oblongato-clavatis, 95 = 15-18, obturaculo apicali jodo violaceo-fuscescente; sporidiis distichis, oblongatis v. fusoideo-oblongatis, curvulis v. rectis, plerumque guttulis, 4-7 majusculis, praeditis v. spurie 3-6-septatis, 28-36 = 7-9; paraphysibus numerosis, simplicibus v. ramosis, 1,5-2 μ . apice clavatis.

Hab. ad corticem Pini silvestris in Mustiala et Germania.

— Sec. Rehm Ascom. n. 865, asci 75 = 12-15; sporidia 27-33 = 6-7, demum 3-5-septata; paraphyses septatae.

7. **Dermatella australis** (Rehm) Sacc. *Dermatea* Rehm Disc. p. 254. 2030

— Sparsa vel gregaria, erumpens, dein sessilis, globosa clausa dein expansa applanata, vix marginata, 0,1-0,8 mm. lat., uda flavo-rubescens, sicca subpulveracea, ceraceo-coriacea; ascis clavatis, 75-80 = 21, 8-sporis; sporidiis elongatis vel fusoideis, initio continuis 1-guttatis demum 3-septatis, 18-24 = 6-7, distichis, hyalinis; paraphysibus tenuibus, divisis, apice incrassatis, flavis.

Hab. in cortice Cytisi nigricantis in Carniola (Voss).

8. **Dermatella dissepta** (Tul.) Sacc. Conspl. gen. Disc. p. 7, *Pezicula* 2031

dissepta Tul. Carp. III, p. 186. — Ascomatibus byssو hypodermeа fuscа subindeque pulvinulo stromatico vix protuberante insertis, caespitosis, erumpentibus, turbinatis, luteolis, tandem late

apertis planisque; ascis oblongis, octosporis; sporidiis ovato-oblongis, curvulis, pallidis, 4-5-locularibus, 20-29 = 7-10; paraphysibus asco longioribus, apice bifurcatis, crassiusculis, subaureis.

Hab. in cortice Castaneæ et Alni circa Parisios Galliæ. — Ex byssso conidia obovato-oblonga continua obtusa stipitata oriuntur. His saepe commixta spermata breve filiformia, 5-7 μ . long.

9. **Dermatella Houghtonii** (Phill.) Sacc. *Dermatea* Phill. Disc. p. 342. 2032

— Solitaria v. saepius cæspitosa, erumpens, cupulis primo globosis, dein turbinatis, stromate communi saepe insertis, sordide flavis, dein brunneis, disco plano vel convexo, immarginato, pruinoso; ascis late clavatis, octosporis; sporidiis oblongo-ellipsoideis, saepe inæquilateralibus, chlorino-hyalinis, triguttatis, dein bi-tri-pseudo-septatis, 27 = 6-9; paraphysibus filiformibus, ramulosis.

Hab. in ramis emortuis Pini lusitanicæ, Lilleshall, Salop Britanniæ (V. HOUGHTON). — Stroma intus flavo-cinnamomeum.

II. *Dermatina*. Sacc. Sporidia submuriformia. Cfr. *D. Frangulæ*.

10. **Dermatella Fagi** (Phill.) Sacc. *Dermatea* Phill. Disc. p. 344. — 2033

Erumpens, cæspitosa, 3-20 mm. lata, cupulis planis vel leniter convexis, plerumque immarginatis, udis aurantiaco-flavis, siccis ferrugineo-flavis, pruinosis, in stromate evidenti dense confertis, stipite (ubi adest) valido, cum stromate continuo; ascis late clavatis, octosporis; sporidiis ellipticis v. oblongo-ellipticis, plasmate grosse granuloso-farctis, subinde muriformibus, 18-23 = 9-12; paraphysibus subtiliter filiformibus, copiosissimis.

Hab. ia ramulis Fagi silvaticæ ad Kingcausie pr. Aberdeen Scotiæ. — Cupulæ 0,5-1 mm. latæ; conidia oblongo-elliptica v. elliptica, 10-20 = 7-9, sporophoris clavatis suffulta et in superficie stromatis inter cupulas stratum pallidum efficientia. — **D. curbitaria** Cooke in Grev. (in Moro in Amer. bor.), sporidiis fennestratis, brunneis, 30-35 μ . long., huic affinis videtur, sed diagnosis deest.

BELONIUM Sacc. Consp. gen. Disc. p. 7 [1884] (Etym. *belone* acus),

Belonidii spec. Auct. — Ascomata sessilia v. breve stipitata, cereo-coriacea, scutellata, sicca subinde hysterioideo-compressa, atra vel brunnea, saepe margine minuto atro-setulosa, glabra vel pilosula v. byssiseda. Ascii elongati, typice paraphysati, octospori. Sporidia elongata v. teretiuscula, 2-pluriseptata, hyalina. — Ad gen. *Belonidium* et sect. *Belonopsis* Mollisiæ sensim abit.

I. *Eu-Belonium*. Ascomata glabrescentia, byssو nullia.

1. **Belonium graminis** (Desm.) Sacc. *Consp. gen. Disc.* pag. 7, *Peziza* 2034 Desm. VIII, *Not.* p. 5, *Belonidium Hystrix* De Not. Prof. *Disc.* p. 381, *Pyrenopeziza hysterina* Sacc. *Mich. II*, p. 244, *Beloni-dium* Speg. in M. N. n. 920, *Pyrenopeziza gram.* Sacc. *F. ital.* f. 1398 (sporidia juvenilia). — Cupulis erumpentibus, scutellaribus, sessilibus, $\frac{1}{4}$ - $\frac{1}{3}$ mm. diam., glabris, ceraceo-coriaceis, siccis compresso-clausis, udis apertis, nigris, disco cinerescente; excipulo solidiusculo, fuligineo, parenchymatico, ad marginem prosenchymatico setulisque brunneis obtusis, 5 μ . cr., coronato; ascis cylindraceo-clavatis, apice rotundatis, 60-70 \approx 10-12, breve stipitatis; paraphysibus filiformibus articulatis obvallatis, 8-sporis; sporidiis subfusoideis, curvulis, 18 \approx 3,5-4, 4-guttulatis, dein 3-septatis, hyalinis.

Hab. in culmis vaginisque Moliniæ cœruleæ putrescentibus in Ital. bor., Gallia, Germania, Belgio. — Sporidia subinde initio tantum 2-guttata.

2. **Belonium Andropogonis** (B. et C.) Sacc. *Peziza Andropogonis* B. 2035 et C. *North Am. Fungi* n. 743, *Cooke Disc. U. S.* I, p. 297. — Cupulis initio clausis, nigris, demum expansis, placentæformibus, margine undulato, disco flavo-badio; ascis clavatis; sporidiis distichis, oblongis, utrinque angustatis, 3-septatis, 15 μ . longis.

Hab. in caulis *Andropogonis* in Carolina Americæ bor. (CURTIS).

3. **Belonium subgibbosum** (Ellis) Sacc. *Peziza subgibbosa* Ellis *Bull. Torr. Club.* 1876, p. 108. — Cupulis sparsis, stipitatis, minutis, vix 2 mm. alt., aquose albidis, siccis obscurioribus, disco convexo, pallido, margine erecto, angusto cincto; ascis cylindraceis, octosporis; sporidiis oblongis, obtusis, hyalinis, demum triseptatis, 20-25 μ . longis, curvulis.

Hab. in culmis emortuis *Andropogonis*, New Jersey Amer. bor. — *Bel. vexato* peraffine. Excipulum fibrosum, tenuiter striatum, subpubescens.

4. **Belonium aberrans** (Peck) Sacc. *Peziza aberrans* Peck *Geolog. Surv.* p. 14, Febr. 1875. — Minutum cœreum, erumpens, subrotundum v. ellipticum, in sicco hysteroideum, disco pallido v. subaurantio, margine angusto nigricante cincto; ascis subfusoideis, majusculis; sporidiis biseriatis v. stipatis, oblongis v. oblongo-clavatis, obscure 3-septatis, hyalinis, 15-20 μ . long.

Hab. in culmis emortuis *Andropogonis*, Newfield, New Jersey Amer. bor. (ELLIS).

5. **Belonium pallens** Sacc. Mich. II, p. 80, F. ital. f. 1286. — Asco- 2038
matibus plano-patellaribus, concaviusculis, sessilibus, $\frac{1}{4}$ mm. diam.,
glabris, disco pallide cinereis, extus brunneolis; excipuli contextu
parenchymatico, marginem versus prosenchymatico, cinereo-ru-
fescente; ascis cylindraceis, 80-55 = 10-12, apice rotundatis, bre-
vissime stipitatis, aparaphysatis(?), octosporis; sporidiis bacillari-
fusoideis curvulis, deorsum attenuatis, 40-45 = 3-3 $\frac{1}{2}$, 3-septatis,
hyalinis.

Hab. in culmis vaginisque *Brachypodii silvatici* prope
Rouen Galliæ (LETENDRE). — Inter *Mollisiam gramineam* Karst.
et *M. excelsiorem* Karst. fere medium; ob sporidia vero bacilla-
ria et distincte septata inter *Belonia* aptius collocandum.

6. **Belonium filisporum** (Cooke) Phill. Disc. p. 162, *Peziza filispora* 2039
Cooke Grev. III, p. 66. — Cupulis sparsis v. subgregariis molliu-
sculis, hemisphæricis, dein applanatis, extus cornicoloribus, in sicco
brunneis, conniventibus; disco pallido v. sordide albo, leniter con-
cavo; ascis tereti-clavatis; sporidiis filiformibus, rectis curvulisve
3-septatis, 35 = 3; paraphysibus filiformibus, simplicibus.

Hab. in vaginis Graminum in Britannia. — Affine *B. ex-
celsiori*.

7. **Belonium? albescens** (Cr.) Sacc. *Patellaria albescens* Cr. Finist. 2040
p. 46. — Cupulis sessilibus, disciformibus, concaviusculis, 2-4 mm.
lat., extus brunneis, disco griseo-albo; ascis minutis, octosporis;
sporidiis oblongo-cylindraceis, hyalinis, 4-5-nucleatis; paraphysi-
bus simplicibus, apice incrassatis.

Hab. in culmis emortuis *Secalis*, Finistère Galliæ.

8. **Belonium pullum** Phillips et Keith in Grev. VI, p. 75, t. 97, f. 8-11, 2041
Disc. pag. 151. — Sparsum; cupulis sessilibus demum explanatis,
aquoso-cinereis, extus brunneis; ascis cylindraceo-clavatis; spori-
diis fusiformibus, nucleatis, dein 3-septatis, 25-30 = 3-5; paraphy-
sibus linearibus, apice ovatis, dissilientibus.

Hab. ad culmos Poæ et Typhæ, Forres, Bristol Shrewsbury
Britanniæ. — Ascoma 1 mm. lat. excipuli cellulæ exteriores elon-
gatæ setulas adnatas formantes.

9. **Belonium? eustegiæforme** (B. et C.) Sacc. *Peziza eustegiæformis* 2042
B. et C. North Am. Fung. n. 741, *Pez. Arundinariæ* Cooke Disc.
U. S. p. 297. — Applanatum, facile deciduum, piceum, e subiculo
maculæformi orbiculari oriundum, disco pallido.

Hab. in *Arundinaria macrosperrmæ* in Carolina infer.
et Georgia.

10. **Belonium? paludosum** (Cr.) Sacc. *Patellaria paludosa* Cr. Finist. 2043

p. 46. — Cupulis brunneolis, sessilibus, disciformibus, 1 mm. lat.; ascis late clavatis, minutis, octosporis; sporidiis fusoideo-elongatis, 8-nucleatis; paraphysibus sursum bifurcatis et piriformi incrassatis.

Hab. in calamis Junci effusi, Finistère Galliæ. — Cellulæ excipuli sphæricæ.

11. **Belonium Arctii** (Phill.) Sacc. *Mollisia Arctii* Phill. Disc. p. 183. — 2044

Sparsum, sessile, initio subglobosum dein expansum, scabrum, nigricanti-brunneum, margine pallido, serrulato, erecto vel subinvoluto, disco udo pallide aquoso-brunneo; ascis late clavatis; sporidiis linearis-fusoideis, 2-septatis, 30-45 = 5; paraphysibus linearibus, obtusis.

Hab. in caule putri Arctii Lappæ prope Bristol Britanniæ (BUCKNALL). — Ascoma 0,5 mm. lat.

12. **Belonium fuscum** Phill. et Harkn. Calif. p. 3. — 2045

fusco-brunneis, longitudinaliter striatis; stipite brevi, obtuso; disco concolore; ascis late clavatis, apice acutis, octosporis; sporidiis oblongis vel oblongo-fusoideis, rectis curvulivis, 1-3-septatis, 15-20 = 4-6; paraphysibus filiformibus, ramulosis obtusis.

Hab. in caulibus emortuis Saniculæ Menziesii in California.

13. **Belonium raphidosporum** (Rehm) Sacc. *Pyrenopeziza raphido-* 2046

spora Rehm Ascom. n. 298. — Sessile in ligni dealbati superficie, semiglobosum, dimidiatum, primitus poro conspicuo pertusum, dein concavum, urceolatum, 0,5 mm. lat., ore dilatato (ita ut discus subfuscus conspicatur), extus rugulosum et circa ostiolum subalbidum, fusco-nigrum, sparsum, elapsum foveolam nigrocinctam relinquent; hypothecio fuscidulo, gonidiophoro gonidiis viridibus, plus minusve magnis ascoma, hymenio evanido, replentibus; ascis clavatis, 8-sporis (an polysporis?) 50 = 9-12; sporidiis parallele positis, filiformibus, rectis vel subcurvatis, plus minusve acutatis, bicellularibus, utraque cellula 2-guttulata, hyalinis, 21-24 = 1,5; paraphysibus, superne conglomeratis.

Hab. ad truncos putridos Pini Cembræ in alpe tyrolensi Kühtai.

II. *Trichobelonium*. Sacc. Ascomata byssiseda vel pilosula.

14. **Belonium retinolum** (Rab.) Sacc. *Tapesia Karst.* Rev. p. 137, 2047

Helotium retincola Rabenh. in Kalchbr. Szepes. p. 238, t. II, f. 1.

— Ascomatibus 2-3 mm. latis tomento effuso, nigro, subreticulato insidentibus, sparsis, sessilibus primitus planiusculis, dein scutellato-marginatis, margine subflexuoso, extus nigro-tomentosis; disco ceraceo, flavo-ochroleuco; carne alba; ascis clavatis; sporidiis cylindraceis, curvulis, (spurie?) bi-triseptatis.

Hab. in culmis subputridis in Hungaria et ad basin culmorum Phragmitis, Wasa Fenniæ. — Ascoma, sec. Karst., 0,5-1 mm. lat., in sicco valde fragile, initio hemisphæricum, totum atrum.

15. **Belonium Stockii** (Cooke et Phill.) Sacc. *Lachnella Stockii* C. et 2048

Ph. Disc. p. 261. — Gregarium v. congestum sessile demum planum extus rufo-brunneum; pilis brevibus septatis brunneis obsesum, disco concavo pallide rufo; ascis tereti-clavatis; sporidiis elongato-fusoideis, 3-septatis, 20-22 = 2-3.

Hab. in caulis putridis herbarum Bungay Britanniæ (Stock).

— Ascoma 0,5 mm. lat., substantia tenuis firmula.

III. *Scelobelonium*. Sacc. Ascomata stipitata.

16. **Belonium melanosporum** Rehm Ascom. Alp. n. 9 (*Belonidium*). 2049

— Cupulis sparsis solitariis interdum binis, breviter crasse stipitatis, stipite circ. 0,5 mm. long. flavescente vel nigricante, urceo-lato-turbanatis, margine albido involuto, nigrofuscis, siccis extus striatis, juvenilibus margine albido involuto, glabris, circ. 1 mm. alt., 0,5 mm. diam., contextu prosenchymatico, flavidus; ascis clavatis, apice incrassatis, 4-8-sporis, 105 = 15; sporidiis oblongis, obtusis, 4-cellularibus, primitus hyalinis, dein subfuscis, muco lato circumdati, mono- v. subdistichis 20 = 5; paraphysibus hyalinis, filiformibus apice sensim in clavam fuscum, 4-5 μ . cr. abeuntibus.

Hab. ad culmos erectos, siccis Luzulæ glabratæ juxta glaciem æternam Taschach-Gletscher in valle Piz alpium tyrolensium, 2200 m. alt.

BELONIDIUM Mont. et Dur. Fl. Alg. t. XXVIII, f. 8 (1846), De Not.

Prof. Disc. p. 380 (Etym. *bēlone* acus ob formam sporidiorum), *Arachnopeziza* Fuck. — Ascomata subsessilia vel breve stipitata, scutellata vel infundibuliformia, ceracea, læticoloria vel pallida (nec fusco-nigricantia) glabra vel puberula, subinde byssō inserta; ascis elongati, typice paraphysati, octospori. Sporidia elongata vel teretiuscula, pluriseptata. — A sect. *Belonopsis* Mollisiæ (quam cfr.) differt præcipue sporidiis distinctius septatis, a *Belonio* ascomatibus seu excipulis numquam vere atris, nec setulosis.

I. *Eu-Belonidium*. Ascomata glabra, byssso nulla.

1. **Belonidium Moliniæ** De Not. Comm. critt. I, pag. 380. — Cupulis 2050 infundibuliformibus, concavis, pallescentibus; disco marginato, excipulo fibroso-ceraceo; ascis clavatis, grandiusculis; sporidiis tereti-oblongatis, 4-loocularibus, hyalinis.

Hab. in culmis siccis Moliniæ cœruleæ, Valle Intrasca Italiæ borealis.

2. **Belonidium Scolochloæ** (De Not.) Sacc. *Helotium Scolochloæ* De 2051 Not. Comm. Critt. I, p. 379. — Ascomate patellari convexo, lutescente subsessili, excipulo crassiusculo; sporidiis fusoideis vel fusoideo-sublanceolatis, hyalinis, 4-loocularibus.

Hab. in culmis Scolochloæ in tectis prope S. Siro, Genova in Italia boreali.

3. **Belonidium tephromelas** (Pass.) Sacc. F. ital. fig. 1285, *Peziza te- 2052 phromelas* Pass. Erb. cr. ital. Ser. II, n. 397. — Cupulis gregariis subsuperficialibus, plano-scutellatis, breve marginatis, glabris, 1 mm. lat., cinereo-olivaceis, ceraceis; ascis clavatis, breve stipitatis, $80 \times 10-11$, apice rotundatis, 8-sporis; sporidiis distichis fusoideis, curvulis, utrinque obtuse attenuatis, 3-septatis, non constrictis, $20 \times 3,5$, hyalinis.

Hab. in calamis Typhæ et Junci effusi pr. Parma Italiæ.

4. **Belonidium sclerogenum** (B. et C.) Sacc. *Peziza sclerogena* B. et 2053 C. Cub. Fung. n. 695. — Sessile, subhemisphæricum, lutescens, margine infexo, demum applanatum; ascis clavatis; sporidiis bi-serialibus, fusiformibus, curvulis, triseptatis, $30-34 \mu$. longis.

Hab. in petiolis Palmarum in insula Cuba.

5. **Belonidium pruiniferum** Rehm Ascom. n. 304. — Gregarium, sessile, 2054 siccum sphæroideum, vix urceolatum, rosellum, albido-pruinosum, scabrum, humidum dilatato-urceolatum vel subplanum, margine crasso, ruguloso, albido circumdatum, hyalino-flavidulum, plerumque 1-1,3 mm. diam., contextu prosenchymatico flavido, fibris marginalibus distantibus, obtusis, c. 3μ . crassis; ascis clavatis, apice obtuse acuminatis, 8-sporis, 75×9 ; sporidiis oblongis, obtusis, rectis vel subcurvatis, 1-4-cellularibus, 2-4-8 nucleis parvulis instructis, hyalinis distichis, $22-24 \times 3$; paraphysibus filiformibus, superne sensim 3μ . cr., hyalinis.

Hab. ad caules putridos Cirsii spinosissimi aliarumque in alpibus tyrolensisibus juxta glaciem æternam. — Apex ascorum jodo valde cœrulescit sporidiaque exacte 4-cellularia apparent.

6. **Belonidium introspectum** (Cooke) Sacc. *Peziza introspecta* Cooke 2055
Hedw. 1875, p. 84, Disc. U. S. I, p. 295. — Gregarium vel sparsum,
minutum diaphanum, cupula sessili, molli, hemisphærica demum scu-
tellari applanatave aquoso-pallida, $\frac{3}{10}$ - $\frac{4}{10}$ mm.; ascis clavatis,
stipitatis; sporidiis arcte fusiformibus, rectis vel curvulis, hyalinis,
4-5 nucleatis, dein leniter 5-septatis, 40-45 \approx 8; paraphysibus li-
nearibus.

Hab. ad lignum cariosum, New Jersey Amer. bor. (ELLIS).

7. **Belonidium immarginatum** Pat. et Gaill. in Bull. Soc. Myc. Fr. IV, 2056
p. 100 (*Beloniella*). — Cupula 260-400 μ . diam., sessili, glabra,
expansa, immarginata, albido-lactea, opalina, fere ex toto ab hy-
menio efformata; ascis clavatis, octosporis, 80-90 \approx 10-12; spori-
ridiis linearibus, fusiformibus, 4-septatis, hyalinis, 4-guttulatis,
40-43 \approx 4-5; paraphysibus ramosis, apice clavatis, intus granulosis.

Hab. ad ligna emortua Puerto-Zamuro Venezuela Amer. austr.
(GAILLARD). — Jodi ops nulla.

8. **Belonidium glabro-virens** (Boud.) Sacc. *Helotium* Boud. in B. S. 2057
bot. Fr. 1881, p. 95, t. II, f. 6. — Ascomate per exiguo, sessili, gla-
bro, smaragdino, hemisphærico, nitido translucido, margine satu-
ratiore, 0,3-0,5 mm. diam., disco plano vel convexulo; ascis cla-
vatis, octosporis, 80-94 \approx 10-14; sporidiis oblongis, curvulis, 3-septa-
tis, 12-13 \approx 3-4; paraphysibus filiformibus, septatis, ramulosis.

Hab. in ligno putri Fraxini, Montmorency Galliæ. — Adest so-
cia forma conidica stilbiformis virens.

9. **Belonidium album** (Cr.) Sacc. *Lecanidion album* Cr. Finist. p. 45. 2058
— Cupulis albis sessilibus disciformibus, $\frac{1}{5}$ mm. lat.; ascis late
oblongis, octosporis; sporidiis subfusoideis, 6-septatis, hyalinis.

Hab. in Helminthosporio quodam ad ramos Coryli, Fini-
stère Galliæ.

10. **Belonidium beloneum** (Cr.) Sacc. *Helotium beloneum* Cr. Fin. 2059
p. 48, *Peziza belonea* Nyl. in litt. — Ascomate albo, sessili $\frac{1}{5}$
mm. diam., glabro, convexo; ascis clavatis octosporis; sporidiis sub-
cylindricis elongatis, uniserialiter guttulatis.

Hab. ad terram, inter Polytricha, Moulin à Poudre Fini-
stère Galliæ.

11. **Belonidium abscedens** (Cr.) Sacc. *Helotium abscedens* Cr. Fin. 2060
p. 48. — Ascomate albo, sessili, glabro, convexo, 200 μ . lat.; ascis
crasse clavatis, octosporis; sporidiis navicularibus, primo granulo-
sis, dein 5-septatis; paraphysibus filiformibus.

Hab. ad terram in declivitatibus, Finistère Galliæ.

II. *Arachnopeziza* Fuck. Ascomata glabra v. pilosula, byssso inserta.

12. **Belonidium Aurelia** (Pers.) De Not. Prof. Disc. p. 381, Pat. Tab. 2061 anal. f. 285, *Peziza aurelia* Pers. M. E. I, p. 270, *P. Wauchii* Grev. t. 139, *Arachnopeziza aur.* Fuck. Symb. pag. 303, *Polynema* Fuck. Symb. App. I, p. 337, f. 35, *Patellaria bicolor* Curr. Linn. Trans. XXIV, p. 494, t. 51, f. 15-16. — Sessilis, sparsa, subventrica, aureo-rubiginosa, pilis adpressis nunc aureis, nunc rubris fimbriata, subiculo tenui niveo; ascis elongato-clavatis, substipitatis, 8-sporis, 90 ± 10 ; sporidiis oblique monostichis, fusiformibus, plerumque rectis, 3-septatis, utrinque ciliatis, hyalinis, 16 ± 4 ; paraphysibus copiosis, filiformibus, simplicibus.

Hab. ad folia, ramulos ipsaque pericarpia faginea aliasque quisquiliis humi jacentes putres in Germania, Gallia, Britannia, Italia, Belgio, Amer. bor. — Cupulæ majusculæ, submembranaceæ, subcrispatae, 2-3 mm. latæ. Color in fungo recente auroræ. Species vere pulchella.

13. **Belonidium delicatulum** (Fuck.) Sacc. *Arachnop. delicatula* Fuck. 2062 Symb. p. 304. — Cupulis sparsis, in villo effuso, delicatissimo, arachnoideo, laxo, candido insidentibus, primo globosis, demum applanatis concavisve, extus pilis candidis, patentibus villosis, disco concavo, rubro-fusco, usque ad 2 mm. lato, margine vix elevato; ascis oblongis, substipitatis, 8-sporis, 80 ± 8 ; sporidiis farctis, filiformibus, quandoque antice subhamatis, rectis curvatisve, continuis, hyalinis, 40 ± 2 .

Hab. in cortice putri dejecto Quercus ad paginam interiorem pr. Eberbach Rhenogoviæ. — *B. Aureliae* affine.

14. **Belonidium auratum** (Fuck.) Sacc. Mich. I, p. 60, F. ital. fig. 1283, 2063 *Arachnop. aurata* Fuck. Symb. mycol. p. 304. — Cupulis $\frac{1}{2}$ mm. latis, sparsis, villo delicatissimo, arachnoideo, candido, late effuso insidentibus, subtus dense, strigose flavescenti villosis, subglabris, primo globosis obconicisve, clausis, demum apertis, margine prominente, brevissime aureo-ciliato, disco concavo, flavo; ascis oblongis; in stipitem attenuatis, 8-sporis, 96 ± 7 ; sporidiis fasciculatis, filiformibus, curvatis, multiguttulatis, antice obtusis, basi acuminatis, hyalinis, 72 ± 2 .

Hab. in cortice interiore putrido, umido Populi pyramidalis in Rhenogovia et in lignis quercinis, Selva, Bavaria (Treviso) Ital. bor. In specim. italicis asci sunt clavati, deorsum atte-

nuati, 110 \approx 10, paraphysibus filiformibus obvallati, octospori; sporidia bacillaria, curvula, 45-50 \approx 2 $\frac{1}{2}$, pluriguttulata, hyalina.

15. **Belonidium araneosum** (Berk.) Sacc. *Peziza areneosa* Berk. Fl. 2064
Tasm. p. 275, *P. arachnoidea* Cooke Austral. fung. p. 50. — Cupulis primum subglobosis demum ex hemisphærico expansis, sessilibus extus araneosis, floccis subtus super matricem repentibus oriundis, hymenio luteo-aurantiaco; sporidiis curvulis multinucleatis, 50-52 μ . long. linearibus.

Hab. in ligno emortuo in Tasmania (ARCHER). — Minuta, extus byssoides. An aptius *Erinella*?

16. **Belonidium fulgens** (Hazsl.) Sacc. *Tapesia fulgens* Hazsl. Einig. 2065
Disc. p. 13. — A *Belonidio Aurelia* differt mycelio flavo-aureo, pilis superne brunneo-fulgentibus; sporidiis minoribus fusoideis, hyalinis, 3-septatis.

Hab. in ligno quercino prope Karlstadt Croatiæ. — Ex auctore forte sporidia spurie septata.

17. **Belonidium? phlegmaceum** (Ellis) Sacc. *Tapesia phlegmacea* Ellis 2066
Torr. Cl. 1882, p. 19. — Gregarium sessile, orbiculare, planum, tenue, album, molliusculum, subiculo laxe arachnoideo, albo, repente insertum; ascis tereti-clavatis sursum abrupte acutatis, 60 \approx 12; sporidiis tereti-fusoideis, deorsum attenuatis, 30-33 \approx 2,5 (septatis?); paraphysibus supra vix incrassatis.

Hab. in truncis dejectis Magnoliæ, New Jersey Amer. bor.

18. **Belonidium basitrichum** Sacc. Fungi Ven. Ser. IV, n. 87, F. ital. 2067
fig. 113. — Cupulis gregariis, sessilibus, patellaribus, ceraceo-mollibus, ambitu circularibus repandisve, margine obtusiusculo vix prominulo totis, albis, subdiaphanis, dein sordide albidis, in sicco vix contractis, $1\frac{1}{4}$ - $1\frac{1}{2}$ mm. diam., disco concaviusculo vix granuloso, excipuli cellulis prosenchymaticis, stipatis, lutescentibus, saturatibus ibique hyphis conidiophoris (*Helminthosporii*), sepiis instar cinctis; hyphis filiformibus, 100 \approx 5, subsimplicibus, septatis, fuligineis, rigidulis, apice conidia fusoidea, curvula, 35 \approx 8, fuliginea 6-8-septata gerentibus; ascis clavatis 100-120 \approx 14, breve attenuato-stipitatis; paraphysibus filiformibus clavulatis obvallatis, 8-sporis; sporidiis oblique recteve distichis, cylindraceo-fusoideis, curvulis, 36-40 \approx 3 $\frac{3}{4}$ -4, initio 7-8-guttulatis, dein 6-8-septatis, hyalinis.

Hab. in ligno quercino udo putrescente, Selva (Treviso) Ital. bor. — *Bel. minutissimo* certe affine, differt vero cupulis sessilibus, sporidiis 6-8-septatis nec tantum 4-septatis.

19. **Belonidium melaneres** (B. et Br.) Sacc. *Peziza melaneres* B. et 2068
Br. Fung. of Ceyl. n. 950. — Cupulis subglobosis, albidis, subtili-
ter pulverulentis, e strato tenui byssideo oriundis; hymenio con-
cavo cinereo; ascis clavatis; sporidiis breviter fusiformibus uni-
septatis 20 μ . long.

Hab. in ramis in insula Ceylon. — An *Solenopezia*?

20. **Belonidium Ruborum** (C. et Ph.) Sacc. *Peziza Ruborum* Cooke 2069
Phil. Grevill. IX, p. 105. — Subgregarium ceraceo-flavum, subi-
culo tenui, evanido, concolori, nidulans; cupulis concavis, demum
applanatis, extus tomentosis, fuscis $\frac{1}{2}$ -1 mm. lat.; tomento tenui,
sparso; ascis clavatis; sporidiis sublanceolatis, demum triseptatis,
hyalinis, 15 \times 5.

Hab. in sarmenis Rubi in Arduennis (LIBERT).

21. **Belonidium leucorrhodinum** (Mont.) Sacc. *Peziza leucorrhodina* 2070
Mont. Cuba p. 360, t. 13, f. 4, *Scutula leuc.* Speg. Fung. Guar.
II, p. 56. — Sparsum vel aggregatum conglomeratumque; cupulis
amœne roseis vel carneis in sicco concavis, margine crasso obtuso,
humectis subgelatinosis, disco vix impresso, subiculo byssino
niveo radiante orbiculari insidentibus; sporidiis cynbi-fusiformibus
. 3-4-septatis, ascis conformibus inclusis.

Hab. in foliis Malvacearum Cubæ-(RAMON DE LA SAGRA), ad
folia viva arborum in Perisporiaceis parasitans, Guarapí Paraguay
(BALANSA). — Sec. Speg. sporidia 12-14 \times 4, primo 4-blasta, dein
didyma ad septum non vel vix constricta, hyalina; asci 50-60 \times
10-12. Quoad genus ambigua species.

22. **Belonidium Asteroma** (Fuck.) Sacc. Mycol. Ven. Spec. p. 164, 2071
t. XVI, f. 23-28, F. it t. 1284, *Arachnop. Asteroma* Fuck. Symb.
mycol. p. 304. — Cupulis gregariis, sessilibus, basi fibrillis delica-
tis, repentibus, fuscis cinctis, $\frac{1}{2}$ lineam latis, extus furfuraceis,
olivaceis, margine pallidiore, disco olivaceo, demum clauso; ascis
oblongis, stipitatis, 8-sporis, 70 \times 11; sporidiis farctis, vermicula-
ribus, subcylindraceis, utrinque obtusis, primo continuis, demum
4-6-guttulatis, simulate septatis, hyalinis, 36 \times 2-3; paraphysibus
asci longitudine, simplicibus.

Hab. in foliis putridis adhuc stantibus Caricis paludosæ
in Rhenogovia et Italia bor. — Planta italica a descriptione Fu-
ckelii non nihil recedit: cupulæ disco demum late hiante, applanato
nec clauso; ascis cylindraceo-subclavati, breve stipitati, 60 \times 9,
8-spori, paraphysibus filiformibus obsoletis obvallati; sporidia aci-
cularia, 35 \times 3, leniter arcuata, utrinque acutiuscula, hyalina, 4-

502 Discomyceteæ, Pezizeæ, Phragmosporæ, Belonidium.

6-nucleata, subdisticha; cupulæ patelliformes, circulares, sinuosæ, diam. $\frac{1}{2}$ - $\frac{2}{3}$ mm., basi villo fusco parco cinctæ, cærulescenti-olivaceæ.

23. **Belonidium heteromorphum** (E. et E.) Sacc. *Peziza heteromorpha* 2072

E. et E. in Journ. Myc. II, p. 88. — Subiculo brunneo-nigro tomentoso, exteuso, plura cm. longo; cupulis sparsis, initio globosis ore minuto, rotundo apertis, margine albo, disco pallido, demum carneo, expanso, plano vel convexulo margineque undulato, 2-3 mm. diam.; ascis tereti-clavatis 70 ± 6 -7, copiose paraphysatis; sporidiis tereti-fusoideis, curvulis, hyalinis, 3-4-nucleatis et demum pseudo-septatis, $20 \pm 2,5$ -3.

Hab. in basi culmorum *Spartinæ polystachyæ*, Louisiana Amer. bor. (LANGLOIS). — Cupulæ juniores albo-puberulæ.

24. **Belonidium? Aliculariæ** (Oud.) Sacc. *Mollisia Aliculariæ* Oudem. 2073

Bijd. IX, p. 379. — Cupulis $\frac{1}{3}$ - $\frac{1}{4}$ mm. depresso-orbicularibus, sessilibus, basi in plantulæ nutrientis parenchymate absconditis, extus dilute, intus saturatius aurantiacis, ore nitidissime circulari præditis; ascis numerosissimis perfecte cylindraceis versus apicem tantum paullulum incrassatis 70 ± 5 , achromis, membrana tenerrima; paraphysibus achromis, subtilissime filiformibus; sporidiis distichis, 23-30 \pm 2, achromis, bacilliformibus, 5-septatis.

Hab. in *Hepatica Alicularia scalari*, Deurne Hollandiæ.

III. *Lasiobelonium* Sacc. Ascomata subtomentosa, byssso nulla.

25. **Belonidium Tami** (Lamy) De Not. Comm. Critt. I, 381, *Peziza* 2074

Tami Lamy in Desm. Descr. Esp. nouv. 1836, p. 10. — Sparsum sessile minutissimum, globoso-applanatum, tenerrimum, subtomentosum, primo albo-hyalinum, dein fulvum; ascis cylindricis; sporidiis ovoideis (?).

Hab. ad caules putrescentes *Tami communis*, Limoges Galliæ. — Sec. De Not. l. c.: ascomata depresso urceolata, leviuscula, ceracea; excipulum fibrosum; sporidia teretiuscula 4-locularia, hyalina.

26. **Belonidium amœnum** Speg. Fung. Fueg. n. 343. — Cupulis super- 2075

ficialibus, sparsis vel hinc inde 2-3-aggregatis, sessilibus, udis globulosis, disco tumido-hemisphærico 250-400 μ . diam., albo-ceraceo donatis, siccis contracto-concavis, globosis, extus flavescentibus villosulis, pilis longiusculis cylindraceis apice obtusis basi non incrassatis 60-100 \pm 1-2, levibus 4-6-septatis non constrictis rigidulis ex hyalino chlorinis; ascis clavulatis, antice abrupte breveque cunea-

tis apiceque truncatis, deorsum lenissime attenuatis 60-70 \pm 5-6, aparaphysatis, octosporis; sporidiis fasciculatis, bacillaribus, vix subclavulatis, superne obtuse rotundatis, inferne attenuato-acutatis 35-40 \pm 1,5, 3-5-septatis non constrictis, hyalinis.

Hab. ad culmos emortuos putrescentes Rostkoviæ grandifloræ in Staten island Fuegiæ.

27. **Belonidium æruginosum** Dur. et Lév. Expl. Sc. Alg. t. 28, f. 8 2076 (absque diagnosi). — Cupulis sessilibus gregariis, hypophyllis, ex urceolato scutellatis, albo-viridulis, tomentellis, ore coarctato, disco brunneolo; ascis tereti-clavatis, deorsum breve attenuatis, octosporis; sporidiis distichis, cylindraceo-fusoideis, triseptatis, denique constrictis, hyalinis; paraphysibus bacillaribus, asco longioribus.

Hab. in foliis emortuis dejectis Ilicis in Algeria.

IV. *Podobelonium* Sacc. Ascomata stipitata.

28. **Belonidium Campanula** (Nees) De Not. Prof. Disc. p. 380, Karst. 2077 M. F. I, p. 162, *Peziza Campanula* Nees Syst. pag. 71, f. 295, Fr. Syst. Myc. II, p. 123, *P. sulphurea* Batsch Cont. I, fig. 146 (ex unico specimine incompleto). — Tenella, cupula membranacea levi, campanulata, flava vel flavicans, stipite brevi filiformi; ascis elongatis; sporidiis teretiusculis utrinque obtusis, 4-6-locularibus, hyalinis.

Hab. in culmis, stipulis putrescentibus in Germania, Fennia. — Gregaria, pellucida, subdifformis, ultra 2 mm. alta. Stipes cupula brevior, subflexuosus, abiens in cupulam late campanulatam margine subrepandam.

29. **Belonidium capense** (K. et C.) *Helotium capense* Kalch. et Cooke 2078 Grev. IX, p. 25. — Sparsum, aurantiacum; cupulis stipitatis, explanatis vel centro depressis, extus pallidioribus, 3 mm. diam.; stipite deorsum attenuato, pallidiore $2\frac{1}{2}$ mm. long., in cupulam expanso, basi nigricante; ascis clavatis; sporidiis fusiformibus, curvulis, 5-septatis, 50 \pm 6.

Hab. ad ramos, P. Natal Africæ austr. (Wood).

30. **Belonidium vexatum** De Not. Prof. Disc. p. 380, Karst. M. F. I, 2079 p. 163, *Belon. culmicolum* (Desm.?) Phill. Disc. p. 148, t. V, f. 29. — Stipitatum, pallescens vel rosello-albidum; cupula cyathoidea vel subcampanulata, in sicco ore coarctato plicatulo subclausa, margine lacero vel fimbriato, disco saturationi, roseo vel testaceo circ. 0,7 mm. lat.; stipite in cupulam sensim abeunte, circ. 0,5 mm. alt.; ascis cylindraceo-clavatis, 100-150 \pm 10-15 sporidiis cylindra-

ceis, hyalinis, rectis vel leniter curvulis, 3-septatis, utroque apice leviter attenuatis, 28-40 = 3-4; paraphysibus filiformibus, apicem versus inæqualiter incrassatis, interdum furcellatis.

Hab. ad culmos Secales, Baldingeræ arundinaceæ, Airæ cæspitosæ, Calamagrostidum nec non Junci effusi in Fennia, Gallia, Germania, Italia. — *Sec.* Phill. huc spectat *Pez. culmicola* Desm. sed sporidiis sec. diagn. valde recedit. — *Var. β. litoreum* Karst. *Belonidium litoreum* Karst. Symb. pag. 213, *Peziza litorea* Fr. Syst. myc. II, p. 121? Partibus internis satis cum typo convenit, sed cupulæ solitariæ vel cæspitosæ, breviter stipitulatæ virescenti- vel rosello-pallidæ aut pallidæ, siccae pallescentes disco concolori vel subtestaceo-pallido, cupula concaviuscula vel convexa, sicca concava vel planiuscula, margine in sicco tenui, incurvo, sublacero. Ad culmos Phragmitis communis in par. Tammela Fenniæ.

31. ***Belonidium minutissimum*** (Batsch) Phillips Disc. p. 149, *Peziza minutissima* Batsch Elench. fig. 143, B. et Br. Ann. Nat. Hist. n. 171, tab. 15, fig. 21, *Peziza Helminthosporii* Blox. Herb. Kew. — Cupulis initio clavatis vel obovatis, dein cupulatis, albo-pallidis, glabris, stipite brevissimo, basi attenuato; ascis clavato-elongatis, octosporis; sporidiis oblongis vel oblongo-fusoideis, curvulis, constricto 4-septatis, 30 = 6, hyalinis; paraphysibus filiformibus.

Hab. in Helminthosporiis ad ramulos, Twycross et prope Bath Britanniæ.

Sect. 5. **Scolecosporæ** Sacc.

GORGONICEPS Karst. M. F. I, p. 15 (1871) [Etym. *gorgon* et *ceps* pro caput, ob ascoma sporidiis serpentiformibus farctum], *Vibrisseæ* Sect. II, *Sessiles* Phill. gen. *Vibr.* p. 8, *Apostemidium* Karst. M. F. I, p. 15. — Ascomata sessilia vel brevissime stipitata ceracea vel ceraceo-mollia, turbinata vel disciformi-scutellata, glabra, mediocria vel minuta. Ascii cylindracei, paraphysati, typice octospori. Sporidia filiformi-acicularia, saepius septulata, subdiaphana, ascum plerumque æquantia. — Cfr. sect. *Belonopsis* generis *Mollisiæ* nec non gen. *Belonium* et *Belonidium*.

1. ***Gorgoniceps aridula*** Karst. M. F. I, p. 185. — Gregaria vel subsparsa, sessilis vel brevissime stipitata, cupula convexa, immarginata, cæsio-hyalina, sicca irregulari, deorsum fuscescente, disco cæsio-hyalino vel (siccitate) griseo-pallido, 0,3-0,8 mm. lat., sti-

pite brevissimo vel indistincto, fusco-atro; ascis clavatis, 82-130 = 11-14, obturaculo minutissimo jodo leviter cœrulecente; sporidiis conglobatis, bacillaribus, arcuatis v. flexuosis, septatis (vulgo septis 8-20) vel guttulatis, hyalinis, 50-88 = 2-2,5, paraphysibus filiformibus, 1,5-2 μ . cr., sursum 2-3 μ . cr., apice attenuato.

Hab. ad corticem crassiorem et lignum vetustum, Messuby et Mustiala Fenniæ.

2. **Gorgoniceps Guernisaci** (Cr.) Sacc. *Vibrissa Guernisaci* Crouan 2082 Finist. p. 46, Phill. *Vibris.* p. 8, t. II, f. 1-7, Gill. Disc. cum ic. Pat. Tab. an. Fung. fig. 369. — Minuta, lenticularis vel turbinata e carnoso subgelatinosa, hymenio pallide ochraceo vel griseo, convexo; ascis cylindraceis; sporidiis filiformibus, hyalinis, 270 μ . long., pluriguttulato-septulatis; paraphysibus filiformibus apice furcato-clavatis.

Hab. in ramis submersis Salicis in Gallia, Britannia. — Ascoma 1-3 mm. diam.

3. **Gorgoniceps Fiscella** (Karst.) Sacc. *Consp. gen. disc.* p. 7, *Aposte- 2083 midium* Karst. *Mycol. Fenn.* I, p. 186. — Subgregaria, tota cæsia vel livido-pallida, granulis seriatis exiguis adnatis obsessa, disco convexiusculo sicco planiusculo, 0,5-1 mm. diam.; ascis cylindraceis, 200 = 5-6,5, apice jodo non tinctis; sporidiis conglobatis, filiformibus, 120-180 = 1; paraphysibus filiformibus, apice ovoideo.

Hab. ad ramulos alneos et salicinos locis humidis circa Mustiala Fenniæ. — A *Gorg. Guernisaci*, ad quam cl. Auctor in *Rev. mon.* p. 134, refert, videtur differre sporidiis multo brevioribus (120-180 nec 270).

4. **Gorgoniceps vibrisseoides** (Peck) Sacc. *Helotium vibrisseoides* 2084 Peck 32 Rep. 1879, p. 48 et Bull. N. Y. Stat. Mus. 1887, p. 28, t. II, f. 7-9, *Vibrissa turbinata* Phill. Gen. Vibr. p. 8, tab. II, f. 14-18 (1881), *Gorgoniceps turbinata* Sacc. *Consp. Gen. Disc.* p. 7. — Sparsa, turbinata, disco plano vel convexo, ochraceo, flavo, margine tenui cinto, extus cyaneo-grisea, levis; ascis cylindraceis, 8-sporis; sporidiis filiformibus 180 = 2; paraphysibus tenuibus apice ramosis clavulatis, brunneis.

Hab. in ramis Fraxini et lignis aqua dejectis, Shrewsbury Britanniæ et Sandlake et Catskill Mount. Amer. bor. — Ascoma $1\frac{1}{2}$ -1 mm. lat. Species americana est paullo major, 2-4 mm., disco livido vel atro-viridulo.

5. **Gorgoniceps decolorans** (Saut.) Sacc. *Peziza decolorans* Saut. Pilz. 2085 Salzb. p. 9, Wint. Hedw. 1881, p. 131. — Alba, sessilis, concava,

carnoso-ceracea, vetusta nigrescens; ascis anguste cylindraceis, 280-290 = 6-7; sporidiis filiformibus, ascum æquantibus, 1-1,5 μ . crass. hyalinis; paraphysibus filiformibus, septatis, hyalinis, apice clavatis, 5 μ . crass.

Hab. in ligno putri prope Mittersill Salzburg. — 2 mm. lat.

6. **Gorgoniceps leptospora** (B. et Br.) Sacc. *Peziza leptospora* B. et 2086
Br. An. Nat. Hist. n. 1166, t. IV, f. 30, *Vibrissa* Phill. Gen. Vibr.
p. 8, t. II, f. 19-23. — Hemisphærica dein expansa, sessilis; disco pallido vel stramineo, extus pilis adpressis sparsis nigris lurida; margine crenulato; ascis oblongis; sporidiis filiformibus 200-230 = 2, seriatim plurinucleatis, demum septulatis.

Hab. in lignis putrescentibus, Jedburg Britanniæ. — Ascoma 1 mm. diam.

7. **Gorgoniceps obscura** Rehm Ascom. Ill. et Exs. n. 714. — Asco 2087
matibus in subiculo ex hyphis fuscis, subramosis, septatis, longis,
6 μ . crassis, composito sessilibus, sparsis, primitus globosis sub-
clausis, ore hyalino, dein urceolatis denique patellaribus, siccis mar-
gine crasso, demum involuto, humidis explanatis, 2 mm. diam.,
extus fuscis vel fusco-cinerellis, scabris, disco cinereo; contextu
parenchymatico, fusco, versus marginem cellulis prosenchymatico-
elongatis, fuscidulis, septatis, apice 6 μ . crassis et divergentibus;
ascis clavatis apice obtuse acutatis, 75-96 = 9, 8-sporis; sporidiis
filiformibus, rectis vel subcurvatis subacutis, hyalinis, 4-6 demum
8-cellularibus, 2-3-stichis, 30-36 = 1-2.

Hab. in truncis Callunæ vulgaris emortuis et inter mu-
scos abditis pr. Königstein Saxoniæ (KRIEGEL). — An potius *Be-
lonium*?

8. **Gorgoniceps dinemasporoides** (E. et E.) Sacc. *Peziza* E. et E. 2088
Journ. myc. I, p. 42. — Nigra, setosa, minuta, puncto centrali af-
fixa, infra celluloso-fibrosa; margine subfimbriato setisque longis
300-400 μ ., nigris patentibus ciliato; ascis clavulatis, 90-100 =
8-10 in basim sensim attenuatis; sporidiis filiformibus multinu-
cleatis e flavo hyalinis 50-65 μ . longis.

Hab. in vaginis inferioribus culmi Andropogonis, Newfield
N. J. Amer. bor.

9. **Gorgoniceps micrometra** (B. et Br.) Sacc. *Peziza* B. et Br. An. nat. 2089
Hist. n. 773. — Per exigua sessilis subturbinata brunnea, villo
albo affixa ore subcontracto subtiliter striato; ascis clavatis; spo-
ridiis filiformibus.

Hab. in calamis emortuis Junc*i*, Twycross Britanniæ. — Asco-
ma sursum nudum, 100 μ . lat. An *Mollisia*?

ERINELLA Sacc., vix Quél. Ench. p. 301 (cujus species *Dasyphydium* spectant). [Etym. *erion* lana]. — Ascomata urceolata vel scutellata, stipitata vel sessilia, villosa vel pubescentia, ceracea, minuta. Ascii teretes vel clavati, typice paraphysati, octospori. Sporidia filiformia vel acicularia, septulata vel guttulata, ascum subæquanta, subhyalina. — Est *Trichopeziza* vel *Dasyphydium* scolecospora.

I. *Eu-Erinella*. Ascomata stipitata.

1. **Erinella simillima** (B. et Br.) Sacc. *Peziza simillima* B. et Br. 2090
Fung. of Ceyl. n. 941. — Cupulis brevissime stipitatis, primum globosis, demum hemisphæricis, albo-tomentosis; hymenio concavo aurantiaco; ascis clavatis; sporidiis filiformibus 100-120 μ . long.
Hab. in corticibus et ramis in insula Ceylon (THWAITES) et Sarawak (BECCARI).
2. **Erinella calospora** Pat. et Gaill. in Bull. Soc. Phyc. Fr. IV, p. 101. 2091
Cupulis sparsis, 2-3 mm. diam., turbinatis, inferne in stipitem brevissimum angustatis, extus villosis albido-disque, intus flavo-aurantiis; pilis externis cylindraceis, elongatis, mollibus et flexuosis; ascis cylindraceis, substipitatis, 120-150 = 12-15, apice poro minutissimo (jodi ope cœrulecente) instructis; sporidiis longissimis, 90-120 = 2-3, utrinque obtusis, 6-14-septatis, viridulis, lucem refringentibus; paraphysibus simplicibus ramosisve, regularibus, non incrassatis, septatis, hyalinis.

Hab. ad cortices arborum viventium inter Maipures et S. Fernando Amer. austr. (GAILLARD).

3. **Erinella orinocensis** Pat. et Gaill. in Bull. Soc. Myc. Fr. IV, p. 101. 2092
— Gregaria; cupula turbinata, extus alba, villosa, in stipitem brevissimum attenuata, margine pilis rugosis ciliato; disco subaurantiaco; ascis cylindraceis, substipitatis, 80-90 = 10, poro exoperculato (jodi ope cœrulecente) instructis; sporidiis rectis curvulisve linearibus, obtusis, 40-45 = 3, hyalinis, 3-septatis; paraphysibus hyalinis, septatis, apice clavato-incrassatis.

Hab. ad caules putrescentes Cannacearum, Puerto Zamuro, Maipures aliisque locis Amer. austr. (GAILLARD). — Fungus circ. 1 mm. diam., ætate proiecta brunneus.

4. **Erinella longispora** (Karst.) Sacc. *Lachnum longisporum* Karst. 2093
Hedw. 1889, p. 191. — Gregaria vel subsparsa, breviter stipitata vel

subsessilis, albida, villosula, cupula primum subsphæroidea, dein planiuscula, sicca subsphæroideo-contracta, 0,5 mm. lat., disco vitellino, stipite brevi crassiusculo vel fere nullo; ascis cylindraceo-clavatis, obtusis, circiter 60 = 6-7, octosporis; sporidiis distichis, filiformi-fusoideis, obsolete 3-septatis, chlorino-hyalinis, 18-40 = 2; paraphysibus 2 μ . cr., apice acutis.

Hab. ad corticem et ramos arborum frondosarum in Minos, Lafayette Brasiliæ. — Pili ascomatum asperi, apice obtusi, 3 mm. crassi.

5. **Erinella minutissima** (Cr.) Sacc. *Peziza minutissima* Cr. Fin. p. 52. 2094

— Cupulis altis, stipitatis, pilosis $\frac{1}{5}$ mm. lat., margine pilis brevibus apice rotundatis fasciculatis ciliato; disco flavidus; ascis cylindricis 8-sporis; sporidiis filiformibus; paraphysibus fusoideis ascis multo superantibus; excipuli cellulis majusculis angulosis.

Hab. in foliis putrescentibus quercinis, Finistère Galliæ.

6. **Erinella aturensis** Pat. et Gaill. in Bull. Soc. Myc. Fr. IV, p. 100. 2095

— Stipitata, 5-7 mm. alta, gregaria vel sparsa; cupula plana, 0,5-1 mm. diam., orbiculari, tenui, extus pruinaosa et albida, margine pilis albis, brevibus, cylindraceis, rugosis, 30 = 3, ciliata; disco plano, flavo-aureo; stipite gracili, elongato, cylindraceo, albidus, glabro inferne incrassato et rufescente; ascis cylindraceis, apice acuminatis, exoperculatis, 60-70 = 1,5-2; sporidiis filiformibus, eguttulatis, 50-64 = 1,5; paraphysibus linearibus, apice incrassatis, hyalinis, ascis longitudine æquantibus.

Hab. ad folia putrescentia, Atures Amer. austr. (GAILLARD).

7. **Erinella Mapiriana** Pat. et Gaill. in Bull. Soc. Myc. Fr. IV, p. 100. 2096

— Stipitata, gregaria 2 mm. alta; cupula plana, orbiculari, margine pilis hyalinis rugosis 60-65 = 3, cylindraceis, apice obtusis ciliata, extus alba; disco tenuissimo diaphano, 0,5 mm. diam., vix flaveolo; stipite gracili, cylindraceo, albo, glabro; ascis cylindraceis, deorsum attenuatis, apice poro pertusis, 55-60 = 3-4; sporidiis filiformibus, gracilibus, 40 = 1; paraphysibus hyalinis simplicibus, linearibus, superne paullulum inflatis.

Hab. ad folia emortua, Mapire Americæ australis (GAILLARD).

8. **Erinella erratilis** Quél. Champ. Norm. pag. 31, t. III, f. 15, Pat. 2097

Tab. anal. t. 588. — Ascomatibus ovoideis, 300-500 μ . diam., firmis, stipite brevi fultis, villosis, albo-roseis et in sicco granuliformibus; disco ceraceo, plano, incarnato-aurantiaco; sporidiis capillaribus, 40-50 μ . longis.

Hab. in calamis Junci exsiccatis in Normandia (A. LE BRE-

Discomyceteæ, Pezizæ, Scolecosporeæ, Erinella. 509

TON). — An potius *Dasyscypha*? : ex icono Pat. enim sporidia asco multo breviora sunt.

9. **Erinella juncicola** Fuck. Symb. mycol. p. 305, *Peziza virginea* 2098 Lasch in Rab. Fung. eur. p. 517, nec Batsch. — *Dasys. virginea* similis, sed multo minor; cupulis stipiteque cupulam vix æquante subfuscis, pilis brevioribus, inferioribus fuscis, marginis niveis; ascis oblongis, 8-sporis, 67×8 ; sporidiis farctis, anguste filiformibus, 40×2 .

Hab. in calamis putridis Junci effusi, Estrich et Driesen Germaniæ.

10. **Erinella hapala** (B. et Br.) Sacc. *Peziza hapala* B. et Br. Ann. N. H. n. 561. — Minuta, sparsa vel stipata; ascomatibus stipite obconico suffultis, extus furfuraceo-villosis, cervinis, disco plano, obscuriori; ascis clavatis; sporidiis elongatis, filiformibus, flexuosis, plerumque longitudinem asci æquantibus.

Hab. in Caricibus et Juncis emortuis in Britannia et Wai-taki Novæ Zelandiæ.

II. *Erinopsis* Sacc. Ascomata sessilia.

11. **Erinella raphidophora** (B. et C.) Sacc. *Peziza raphidophora* 2100 B. et C. Cub. Fung. n. 683. — Primum globosa, breviter albo-floccosa, plus minus furfuracea, demum expansa, margine rotundato inflexo; ascis elongatis; sporidiis linearibus utrinque acutis, brevioribus 25μ . long., sed sæpius longioribus, utrinque endochromate contracto.

Hab. in lignis putridis in Venezuela et Cuba. — Ascoma 2 mm. diam.

12. **Erinella raphidospora** (Ellis) Sacc. *Peziza* Ellis Torrey Bot. Club. 2101 1876, p. 107. — Gregaria sessilis, minuta, subconfluens, albo-tomentosa, disco plano vel leviter convexo; ascis linear-clavatis, apice acutis vel obtusis; sporidiis filiformibus ascum æquantibus, seriatim multinucleatis.

Hab. in trunco vetusto Pini, Newfield N. J. Amer. bor.

13. **Erinella leucophæa** (B. et C.) Sacc. *Peziza leucophæa* B. et C. 2102 North Pac. Exp. n. 152. — Cupula concava, intus alba, extus floccis brunneis subfasciculatis aspera; sporidiis longis filiformibus.

Hab. in ramulis emortuis in Japonia.

14. **Erinella pseudofarinacea** (Cr.) Sacc. *Peziza pseudofarinacea* Cr. 2103 Fin. p. 52. — Cupulis albis, ovoideis dein turbinatis, 0,5 mm. lat., pilosis, pilis costulas verticales formantibus; margine laciniato;

disco stramineo; ascis clavatis; sporidiis filiformibus hyalinis; paraphysibus linearibus.

Hab. ad ramulos dejectos Pini maritimæ, Morlaix Finistère Galliæ (DE GUERNISAC).

15. **Erinella rhabdocarpa** (Ellis) Sacc. *Peziza rhabdocarpa* Ellis ²¹⁰⁴ Torr. Cl. 1882, p. 19. — Sparsa sessilis, $\frac{1}{2}$ mm. diam., tenuis, margine pilis brevibus fasciculatis obtusis clavato-capitatis fimbriato; excipuli cellulæ vesiculosæ stipatis; ascis oblongo-cylindraceis sessilibus, apice acutiusculis, $65-75 \approx 10$; sporidiis linearibus multinucleatis (an demum multiseptatis?) flavidis ascum subæquantibus, 2-3 μ . crass.

Hab. in ramis emortuis Comptoniæ, New Jersey America boreali.

16. **Erinella miniopsis** (Ell.) Sacc. *Peziza miniopsis* Ellis Bull. Torr. ²¹⁰⁵ 1881, p. 66. — Gregaria, sessilis, minuta, 500-750 μ . diam., pilis tomentosis, albis, crispis tecta; disco cinnabarino, cupuliformi; ascis clavato-cylindraceis, $95 \approx 12$; sporidiis stipitatis, lineari-lanceolatis, 50-65 μ . longis, lutescentibus, nucleatis, demum multiseptatis; paraphysibus filiformibus sursum non incrassatis.

Hab. in cortice exteriori vivo Aceris rubri in Amer. bor.
— Discus udus aurantiacus, siccus miniatus.

17. **Erinella brevipila** (Rip.) Sacc. *Peziza* Rip. Bull. Bot. Franc. 1876, ²¹⁰⁶ p. 220 et Malbr. et Let. Champ. Norm. p. 61, f. 10, Mout. Ascom. Liege p. 139. — Cupulis sessilibus, subsparsis, minutis, scutellatis, pilis brevibus, crassiusculis, rugulosis, brunneo-atris marginatis; ascis cylindraceis, $55-70 \approx 10-16$, octosporis; sporidiis bacillari-fusoideis, 30-40 ≈ 3 , pluri-guttulatis, dein 3-septatis, hyalinis.

Hab. in caulis Centaureæ et Achilleæ in Gallia et Belgio.

18. **Erinella mutabilis** Quél. Champ. Normand. p. 31, nec B. et Br. ²¹⁰⁷ — Sessilis, subcoriacea, tenuiter tomentosa, cervina dea fusca, 1 mm. diam., margine cilio lato albo, disco carneo dein albido; sporidiis acicularibus, 40 μ . longis, seriatim guttulatis, hyalinis.

Hab. in culmis siccis Brachypodii in Normandia (LE BRETTON).

19. **Erinella albo-carnea** (Cr.) Sacc. *Peziza albo-carnea* Cr. Fin. ²¹⁰⁸ p. 52. — Cupulis albis, subsessilibus, hemisphæricis, pilosis; disco incarnato pilis fasciculatis cincto; ascis subcylindricis 8-sporis; sporidiis tereti-filiformibus; paraphysibus fusoideis, excipuli cellulis prosenchymaticis anastomosantibus tenuissimis.

Discomyceteæ, Pezizeæ, Scolecosporæ, Peziza. 511

Hab. ad calamos Junci conglomerati in paludosis Finistère Galliæ.

Species *Pezizearum* mihi ignotæ.

1. **Peziza catulacea** Desm. Cat. pl. omiss. p. 15. 2109
Hab. in agro lillensi Galliæ.
2. **Peziza clavellata** Desm. Crypt. Voges. exs. n. 175. 2110
Hab. in agro lillensi Galliæ.
3. **Peziza Spirææ** Kirchn. in Lotos 1856, p. 246. 2111
Hab. in Bohemia meridionali.
4. **Peziza Armeriæ** Lasch in Klotzsch Herb. myc. n. 1126 et in Bot. Zeit. 1848, p. 292 (nomen). 2112
Hab. in agro driesensi Germaniæ (LASCH).
5. **Peziza Verbenæ** Kirch. in Lotos 1856, p. 246. 2113
Hab. in Bohemia meridionali.
6. **Peziza epidermidis** Opiz Beitr. p. 25. 2114
Hab. in agro pragensi Bohemiæ.
7. **Peziza fungorum** Opiz in Lotos 1856, p. 246. 2115
Hab. in Bohemia meridionali.
8. **Peziza Juniperi** Kirchn. in Lotos 1856, p. 246. 2116
Hab. in Bohemia meridionali.
9. **Peziza Kirchneri** Opiz in Lotos 1857, p. 28. 2117
Hab. in Bohemia meridionali.
10. **Peziza Koerberi** Peyl in Lotos 1858, p. 31, t. 20. 2118
Hab. in Bohemia orientali.
11. **Peziza flava** Sw. Fl. Ind. occid. III, p. 1933. 2119
Hab. in Jamaica.
12. **Peziza phæoloma** Lév. in Demid. Voy. p. 97, t. 4, f. 7. 2120
Hab. in Tauria.
13. **Peziza raudanensis** Lecocq Rech. sur Raudan., Streinz Nom. pag. 433. 2121
Hab. in agro parisiensi Galliæ.
14. **Peziza Artemisiæ** Kirchn. in Lotos 1856, p. 246. 2122
Hab. in Bohemia et Germania.
15. **Peziza limosa** Balb. Fl. lugd. II, p. 311. 2123
Hab. in agro Brixiae Ital. bor.
16. **Peziza aterrima** Lasch in Klotzsch Herb. myc. n. 336. 2124
Hab. in Germania et Suecia.

Fam. 4. **Ascoboleæ** Boud.

Mém. Ascob. p. 20 (1869), Sacc. Consp. gen. Disc. p. 7.

Ascomata carnosa vel carnosò-gelatinosa, sessilia vel brevissime stipitata, plana vel convexa, glabra vel pilosa vel furfuracea. Asci 4-8-polyspori, apice operculatim dehiscentes, paraphysati, a disco mox exsiliens eumque echinatum v. papillatum reddentes. Sporidia globosa, ellipsoidea vel subfusoidea, continua, colorata vel hyalina. — Plerumque fimicola, rarius humi-carbonicola vel phytogenæ.

Clavis analytica generum.

Sect. 1. **Phæosporæ** Sacc. Sporidia colorata.

Boudiera. Sporidia globosa, intra ascum libera.

Ascobolus. Sporidia ellipsoidea vel subfusoidea, intra ascum libera.

Saccobolus. Sporidia ellipsoidea v. subfusoidea, intra ascum sacculo communni inclusa.

Sect. 2. **Hyalosporæ** Sacc. Sporidia hyalina.

Cubonia. Sporidia globosa, octona.

Ascophanus. Sporidia ellipsoidea vel subfusoidea, octona; ascomata nuda.

Lasiobolus. Sporidia ellipsoidea vel subfusoidea, octona; ascomata pilosa.

Ryparobius. Sporidia ellipsoidea vel subfusoidea, numerosa; ascomata perfecta.

Gymnодiscus. Sporidia ellipsoideo-oblonga, numerosa; ascomata margine et excipulo parentia. (Ad *Gymnoscaceas* nutans).

Sect. 1. **Phæosporæ** Sacc.

BOUDIERA Cooke Grev. VI, p. 76 [1877], (Etym. ab E. BOUDIER, celeberrimo monographo Ascobolearum). — Ascomata immarginata, orbicularia discoidea, plana vel convexa, carnosa; hymenio papillato. Asci clavati v. clavato-cylindracei, erumpentes. Sporidia globosa, colorata. Paraphyses distinctæ. — Cfr. *Phœopezia*, sectio sphærosporarum.

1. **Boudiera areolata** C. et Ph. Grev. p. 76, t. 97, f. 12-15, Phill. Disc. p. 285, t. IX, f. 53, *Plicaria tracheja* Rehm Ascom. n. 451, — Atro-brunnea; cupulis orbicularibus convexis; ascis 8-sporis, clavato-cylindraceis, 24 μ . cr.; sporidiis globosis, areolatis, 26-30 μ . diam., hyalinis dein brunneis; paraphysibus robustis, clavatis, septatis, supra purpureo-brunneis, sursum 8-10 μ . cr.

Hab. ad terram udam, N. Wales Britanniæ et in Germania. —

Ascoma 2-2,5 mm. lat. Asci jodo cœrulescunt, paraphyses violascunt.

2. **Boudiera microscopica** (Crouan) Cooke Grev. VI, p. 76, *Ascobolus* 2126
microscopicus Cr. Finist. p. 57 et Ann. Sc. Nat. 1857, t. IV, E,
fig. 20-23. — Sessilis, subhemisphærica, brunneola, glabra, disco
plano; ascis clavatis, octosporis; sporidiis sphæricis, brunneis, echinatis; paraphysibus simplicibus, crassis, sursum latoribus, curvis,
hyalinis.

Hab. in excrementis caninis, Finistère Galliæ. — Contextus
anastomotico-filamentosus.

3. **Boudiera marginata** Phill. et Harkn. Calif. pag. 4. — Gregaria, 2127
initio orbicularis, dein patellata, margine serrulato, sordide ochra-
cea; ascis late clavatis, octosporis; sporidiis sphæricis, mox rufis,
areolatis, 20 μ . lat.; paraphysibus linearibus, obtusulis.

Hab. in fimo cunicolorum in California. — Cellulæ exci-
puli crasse tunicatae.

4. **Boudiera nivea** (Quel.) Sacc. *Ascobolus niveus* Quél. X Suppl. 2128
p. 14, t. IX, f. 18. — Ascomatibus hemisphæricis, dein lenticula-
ribus, 500-800 μ . diam., glabris, hyalinis, diaphanis, margine mem-
branaceo, subtilissimo, undulato; hymenio plano, albo, lilacino-
punctulato; ascis cylindricis; sporidiis sphæroideis, 13-14 μ . diam.,
eleganter granulosis, lilacinis.

Hab. in fimo vulpium, Jura Galliæ. — Cum *Phæopezia ca-*
nina comparanda.

5. **Boudiera hyperborea** (Karst.) Sacc. *Ascobolus hyperboreus* Karst. 2129
M. F. I, p. 80. — Gregaria, sessilis, nuda, planiuscula, pallescens,
minuta; ascis subcylindraceis, 16-18 μ . cr., 8-sporis; sporidiis
monostichis, sphæroideis, papillose asperulis, dilutissime violascenti-
fuscis, 10-14 μ . diam.

Hab. supra fimum Lemmi norvegici in umbrosis humidis
prope oppidum Kola Lapponiæ rossicæ.

6. **Boudiera Brassicæ** (Crouan) Sacc. *Ascobolus Brassicæ* Crouan 2130
Fl. Finist. p. 57 et in Ann. Sc. Nat. 1857, p. 174, tab. IV, B, C,
f. 9-14. — Alba, sessilis, hemisphærica, concava, 2-3 mm. lata,
marginè erecto, denticulato, disco pallide brunneo-violaceo; ascis
tereti-clavatis operculatim dehiscentibus, octosporis; sporidiis sphæ-
ricis, distichis v. monostichis violaceo-brunneis; paraphysibus hya-
linis, apice furcatis et gelatinoso-cohærentibus.

Hab. in trunculis putridis Brassicæ, Finistère Galliæ. — Con-
textus excipuli hexagono-cellulosus. Sporidia ex iconæ, 16-18 μ . lat.

ASCOBOLUS Pers. in Gmel. Syst. p. 1461 ex p. [1791] (Etym. *ascos* ascus et *ballo* ejicio). — Ascomata carnosula, sessilia vel subsessilia scutellata v. plana v. convexula, glabra v. pilosella. Asci cylindraceo-clavati 4-8-spori, operculo apice dehiscens, varie paraphysati, exsilientes. Sporidia ellipsoidea vel breve fusoidea, eximie longitud. rimulosa vel reticulata vel levia, cyanea, violacea vel brunnea.

I. *Eu-Ascobolus*. Ascomata glabra.

a) Species fimicolæ.

1. **Ascobolus æruginosus** Fr. Obs. II, p. 310, Boud. Ascob. pag. 32, 2131 t. VII, fig. 72, *A. marginatus* Schum. Sael. 2, p. 437. — Sessilis gregarius, globosus v. subcylindraceus, planiusculus, immarginatus, glaber, viridis v. flavo-olivaceus; ascis anguste clavatis, 8-sporis; sporidiis ellipsoideis violaceis v. purpureo-violaceis, demum brunneis, anastomotico-rugosis, subinde levibus, 15-17 = 8; paraphysisibus filiformibus, septatis, hyalinis, muco sulphureo-viridi obvolutis.

Hab. in fimo equino, vaccino et vulpino in Suecia, Dania, Britannia, Gallia, Belgio. — Ascoma $\frac{1}{2}$ -1 mm. lat., siccum nigricans. Species Crouani (Finist. p. 57) sporidiis globulosis videtur diversa. Sec. Boud. l. c. hoc forte spectat *A. immarginatus* Becc.

2. **Ascobolus immarginatus** Beccari Erb. Critt. ital. n. 775. — Major 2132 2-10 mm. lat., gregarius, obconicus vel hemisphæricus, concavus, madore ambitu subrecurvus, saepe irregularis, primum luteo-virescens, dein sporidiis emissis, disco fusco-violaceus, glaber, margine nudo, obtuso, levi; ascis oblongis, octosporis; paraphysisibus filiformibus, articulatis, apice dilatatis; sporidiis violaceis, ellipsoideis, obtusis, obscure venulosis, 13-16 = 8-9.

Hab. in fimo in horto botanico, Pisa in Italia media. — Cfr. *Ascob. æruginosus*.

3. **Ascobolus viridulus** Phill. et Plowr. in Grevillea VIII, p. 103. — 2133 Sparsus, sessilis, hemisphæricus, concavus, dein expansus, submarginatus, crassus, pallide flavescens-viridis, grosse granulatus; ascis clavatis, octosporis; sporidiis mono- vel distichis, ellipticis, violaceis, dein brunneis, 13 = 6, rugoso-striatis; paraphysisibus validiusculis, distincte septatis, apice clavatis, muco pallide viridi obvolutis.

Hab. in stercore avis (Columbi?) ad Shrewsbury Britanniæ. — Cupula 0,5-1 mm. diam. Habitus *Asc. furfuracei* juvenilis.

4. **Ascobolus globularis** Roll. Bull. Soc. mycol. IV, pag. 57, t. XV, 2134
fig. I. — Carnosus, luteo-virens, extus furfuraceus, piriformis,
sed basi immersa globulum fingens, $\frac{1}{2}$ -1 mm.; disco plus minus
plano aut convexo; ascis paucis valde prominentibus claviformibus,
maximis, amplis, 280-300 \approx 70-80, breviter pedicellatis, 8-sporis;
sporidiis sphæroideis, maximis, 33 \approx 28, sacco hyalino singulatim
inclusis v. aliquando nudis, intense violaceis, brunnescentibus, le-
vibus vel aliquando-verrucosis; paraphysibus hyalinis, gracilibus,
ad apicem leviter v. vix incrassatis, plerumque simplicibus, junio-
ribus granulosis, dein septatis ascisque gelatina sulphurea immersis.

Hab. ad stercus caprinum in horto botanico Parisiorum
Galliæ.

5. **Ascobolus lapponicus** Karst. M. F. I, p. 80. — Gregarius, sessilis, 2135
glaber, gelatinosus, plano-concavus vel plano-convexus, submargi-
natus, flavidо-pallens, disco virescente-pallido, 0,6 mm. lat.; spori-
diis octonis, oblongato-ellipsoideis, violaceo-fuscis, 18-21 \approx 12-13,5.

Hab. supra fimum Lemmi norvegici prope oppidum Kola
Laponiæ rossicæ.

6. **Ascobolus crenulatus** Karst. M. F. I, p. 77. — Gregarius, sessilis, 2136
primum sphæroideus, dein applanatus, planiusculus, virescenti-lu-
teus, furfuraceus, margine crenulato, circiter 1,5 mm. lat.; ascis
clavatis jodo haud tinctis, 12 μ . cr., octosporis; sporidiis mono-
stichis, ellipsoideis, striatulis, violaceo-fuscescentibus, 12-16 \approx 6-8;
paraphysibus gracilescitibus.

Hab. supra fimum Tetraonum et vaccinum pr. Mustiala
Fenniæ, in Britannia, Oneida et Helderberg Mts. Amer. bor. —
Ab affinis cupulis minoribus, margine crenulato, sporidiis mi-
noribus etc. differt.

7. **Ascobolus stictoideus** Speg. Mich. I, p. 474. — Laxissime gre- 2137
garius v. sparsus, minutus, vix $\frac{1}{3}$ - $\frac{1}{2}$ mm. diam., externe aquose
albidus, disco dilute olivaceo ex ascis exsiliens nigro-punctato,
totaliter in fimo immersus, non v. labiis tantum exsertis, crassiu-
sculis atque laceris; ascis paucis 5-10, magnis, 150-160 \approx 40, cla-
vato-saccatis, breviter et crasse stipitatis, superne crassiuscule tu-
nicatis, octosporis; paraphysibus septulatis, apice curvulis obval-
latis; sporidiis distichis vel irregulariter polystichis, ellipticis, 25
 \approx 14, primo hyalinis, levibus, dein amœne violaceis minute, ac den-
sissime verruculosis, postremo roseo-fuligineis.

Hab. in fimo canino putrescente, Conegliano Ital. bor. —
Tunica ascorum jodi ope obsolete cœrulescit.

8. **Ascobolus cubensis** B. et C. Cub. Fung. n. 707. — Parvus, cra-²¹³⁸
teriformis, viridi-flavus; ascis linearibus; sporidiis ellipticis pallide
fuscis, longitudinaliter rugoso-plicatis, 10-5.

Hab. in fimo suino in insula Cuba.

9. **Ascobolus cervinus** B. et Br. Fung. of Ceyl. n. 1209. — Oliva-²¹³⁹
ceus, extus furfuraceus; margine discreto; ascis rectis, oblongis;
sporidiis minoribus, 15 μ . long.

Hab. in fimo, Peradenya Ceylon. — Affinis *A. æruginosa* sed
asci recti multo minores etc.

10. **Ascobolus Marchalii** Bomm. et Rouss. Fl. Myc. Brux. p. 131. — 2140
Ascomatibus carnosis, pallide flavis, distincte stipitatis vel in sti-
pitem compactum subsulcatum, furfuraceum, 4-5 mm. alt. desi-
nentibus; disco 4-8 mm. lat., margine integro parum inflexo et
compacto, demum laciniato, ob sporidia exslientia brunneo-viola-
scente; ascis claviformibus, 150-180 μ . longis; paraphysibus fili-
formibus, rectis vel curvulis, flavo-viridulis, subinde apice infla-
tis; sporidiis ellipticis, 15-9, primo hyalinis, dein violascentibus,
striatis.

Hab. in fimo vaccino in Belgio. — Affinis *A. lignatili* et *A. Crouani*, a quibus differt stipite saepe distincto v. striatulo, ma-
gnitudine et habitatione.

11. **Ascobolus marginatus** Pat. Tab. an. Fung. pag. 32, fig. 76, nec 2141
Schum. — Carnosulus, plerumque pedicellatus, rarius subsessilis,
3-4 mm. lat., extus flavidus, glaber, margine dentibus furfuraceis
subfimbriato, disco brunneo piano v. concaviusculo, ascis punctato;
ascis cylindraceis octosporis; sporidiis monostichis, ovoideis, 17-
18-9-10, violaceo-purpureis, rimulis ramulosis percursis; para-
physibus filiformibus.

Hab. in fimo aquino, Poligny Galliæ. — *Ascob. lignatili* et
furfuraceo subaffinis.

12. **Ascobolus Daldinianus** De Not. Comm. critt. I, p. 361. — Ha-²¹⁴²
bitu *Ascophani granuliformis*, differt paraphysibus apice haud
incrassatis; sporidiis violaceis cum nucleo subtiliter punctulato.

Hab. in fimo vaccino, Locarno (DALDINI). — Flavo-ochra-
eus, glaber.

13. **Ascobolus furfuraceus** Pers. Obs. I, p. 33, t. IV, f. 3-6, Bern. 2143
Manip. IV, t. VI, f. 5, Boud. Ascob. p. 26, t. VI, f. 6-10, *Elvella*
fimetaria Scop. Fung. Hung. in Ann. Hist. Nat. IV, p. 149, t. 1,
f. 6, *Peziza alra* Huds. Engl. II, p. 637, *Pez. stercorearia* Bull.
Champ. p. 256, t. 376, Sowerb. fung. t. 18, 389, f. 3-6, *Pez. su-*

sca Bolt. fung. t. 109, f. 2. — Sessilis, subconcavus, fuscus virescens, externe furfuraceus; ascis cylindraceo-clavatis, deorsum longe attenuato-stipitatis, apice rotundato-subtruncatis, 140-150 \approx 20-22; paraphysibus filiformibus, septulatis obvallatis, octosporis; sporidiis oblongo-ellipsoideis, 22-30 \approx 10-14, distichis vel oblique monostichis, primo violaceis, dein vinoso-fuligineis, longitudinaliter ramosse striatulis, muco obvolutis.

Hab. in fimo vaccino et equino, rarius felino, cervino et humano in Suecia, Gallia, Germania, Britannia, Belgio, Italia, Austria, Amer. bor. et austr., Africa et Oceania. — Ascoma 2-4 mm. lat. subglobosum dein disco concavo apertum. — Var. **nudus** Kickx Fland. I, p. 479: fulvo-virens, extus nudus. Ibidem in Belgio et Italia.

14. **Ascobolus glaber** Pers. Obs. Myc. I, p. 34, t. 4, f. 5, b, t. 7, a, ²¹⁴⁴ b, c, Syn. p. 677, Nees. Syst. fig. 297, Fr. Syst. Myc. II, p. 164, Boud. Ascob. p. 33, t. VII, f. 13-15. — Sessilis, minutus, glaber, nitens, subconvexus, marginatus, spadiceus, globoso-oblongus, dein piriformis; ascis late tereti-clavatis, 8-sporis; sporidiis ellipsoideis violaceis, longitud. reticulato-rugosis, 25-30 \approx 12-14; paraphysibus filiformibus.

Hab. in fimo vaccino, equino et cuniculorum in Suecia, Germania, Britannia, Gallia, Italia, Fennia. — Species conferta, superficialis, valde varia; quoad diversum maturitatis gradum forma tereti, conica, convexa, depressa etc. gaudet; $1\frac{1}{2}$ -1 mm. lata. Etiam colore varius, primo rubens, dein spadiceus, demum niger, variat aurantius v. albus (*A. albidus* Cr. Finist. p. 57).

15. **Ascobolus rufo-pallidus** Karst. M. F. I, pag. 8. — Subgregarius, ²¹⁴⁵ obconicus, sessilis, glaber, rufescens; disco plano-concavo, papillato, pallidiore, margine tenui fimbriato, 0,5 mm. lat.; sporidiis octonis, ellipsoideis, 16-26 \approx 12-13.

Hab. supra fimum Lemmi norvegici propæ oppid. Kola Lapponiæ rossicæ. — Ipsi auctori dubia species; sporidia primo nigra, denuo examinata hyalina visa.

16. **Ascobolus minutus** Boud. Bul. Soc. Myc. IV, p. XLVIII, t. II, ²¹⁴⁶ f. 1. — Gregarius aut sparsus, minutissimus, 0,5-0,7 mm. æquans, brunneus, extus levis dilutior, immarginatus, obconicus, aut hemisphericus, dein lenticularis, extus glaber fulvus, disco plano aut convexo saturatiore, ascis parum superantibus nigro-papillato; paraphysibus septatis, hyalinis, gelatina lutea saepius conspicua ascisque immersis, ad apicem subclavulatis, 5-7 μ . cr.; ascis minutis,

clavatis, hyalinis, octosporis, 140-170 \approx 13-14; sporidiis minutis, ellipticis, violaceis, densique fuscis, longitudinaliter striatis, striis vix anastomosantibus, 13-15 \approx 7-9.

Hab. in stercore vulpino, Montmorency Galliæ.

17. **Ascobolus vinosus** Berk. Engl. Fl. V, p. 209, Boud. Ascob. p. 31, 2147 tab. VI, fig. 11, Pat. Tab. an. fig. 381, B. et Br. Ann. Nat. Hist. n. 1083, tab. XVI, f. 25. — Sessilis, subglobosus, dein depresso, glaber, vinoso-purpureus, margine laciniato; disco plano v. convexo, livido-purpurascente; ascis late clavatis, octosporis; sporidiis ellipsoideis, hyalinis, mox purpureis, demum brunneis, 23 \approx 13, episporio anastomotico-rugoso; paraphysibus filiformibus, septatis, vix coloratis, muco livido obductis.

Hab. in fimo cuniculorum et vaccino in silvis in Britannia, Gallia et Germania.

18. **Ascobolus amœnus** Oud. Hedw. 1882, n. 11 et Bijd. IX, p. 381. — 2148 Sparsus, minutus, $4/10$ mm. diam., parum prominens, extus pallide ochraceus, glaber, levis, floccorum tenerorum strato insidens, disco ex ascis valde emergentibus nigro-punctato; ascis plurimis in quovis ascomate, amplis clavatis, breve pedunculatis, 230 \approx 35, tunica toto ambitu æquicrassa, 8-sporis; paraphysibus quam plurimis, tenerrimis, septulatis, apice rectis obvallatis; sporidiis distichis, primitus achromis, postea violaceis, denique fuscescentibus, ellipticis, 36 \approx 15, statu colorato tantum densissime subtilissimeque granulatis (verruculosis).

Hab. in fimo Cameli bactriani, Amsterdam Nederlandiæ (VAN LEDDEN).

19. **Ascobolus australis** Berk. Linn. Journ. XVIII, p. 389. — Brunneus; ascis clavatis; sporidiis ellipticis, purpureo-fuscis, levibus, 60-72 μ . longis; paraphysibus linearibus apice latioribus. — 2149

Hab. in fimo, Rockhampton Australiæ (THOZET).

20. **Ascobolus Baileyi** B. et Br. Fung. of Brisb. II, p. 69. — Cupulis 2150 primum ochraceis concavis, demum vinoso-brunneis explanatis, extus leviter granulatis; ascis clavatis prominentibus; sporidiis mono-distichis brunneis, ovatis, levibus, 15-18 μ . longis; paraphysibus linearibus, apice clavulatis.

Hab. in fimo, Brisbane Australiæ (BAILEY).

21. **Ascobolus leiocarpus** B. et Br. Fung. of Ceyl. n. 971. — Cupulis 2151 extus rubiginosis furfuraceis; hymenio fusco; sporidiis ellipticis levibus v. tenuissime lineolatis.

Hab. in fimo Peradenya (THWAIRES), in fimo bubalino Ceylon (BECCARI).

22. **Ascobolus major** B. et C. North Amer. Fungi n. 772. — De- 2152
planatus, margine eximio, prominente, umbrino; sporidiis ellipticis
levibus, $28-30 \mu$. long.

Hab. in fimo in Carolina inf. Amer. bor.

23. **Ascobolus Leveillei** Boud. Ascob. p. 35. t. VII, f. 16, nec Crouan. 2153
— Minutus, congestus, brunneus, extus opacus et minutissime fur-
furaceus, 1 mm. lat., subpiriformis, semimmersus; ascis longissi-
me exsertis, amplis, clavatis, 180 ± 30 , octosporis; sporidiis distichis
obtuse ellipsoideis, hyalinis, dein brunneis, $25-27 \pm 16$, episporio
leviter reticulato; paraphysibus filiformibus, apice vix incrassatis.

Hab. in stercore equino prope Paris Galliæ et in Germania.

— Ascis gelatina sulphurea interposita longe exslientibus facile
distincta species. Asci jodo cœrulescunt, episporium violascit.

24. **Ascobolus sphæricus** Preuss Hoyersw. n. 156. — Immersus, mi- 2154
nutus, sphæricus, rugosus; ascis cylindricis hyalinis albis; spori-
diis octo, ovatis nigrescentibus; episporio glabro; guttulis oleosis
subbinis; paraphysibus clavatis.

Hab. in fimo canino prope Hoyerswerda.

b) Species humi-carbonicolæ.

25. **Ascobolus viridis** Curr. Linn. Trans. XXIV, p. 154, 1863 Phill. 2155
Disc. p. 289; t. IX, f. 54, nec Boud. — Ascomatibus sparsis ses-
silibus v. substipitatis, planiusculis, furfuraceis v. tomentellis ob-
scure viridi-luteis, 1,5-2 mm. lat.; ascis clavatis, jodo non tinctis,
 $18-20 \mu$. lat., octosporis; sporidiis ellipsoideis acuminatis, rugoso-
striatis, violaceo-fuscescentibus, $25-32 \pm 11,5-13$; paraphysibus nu-
merosis, articulatis, 2μ . crassis, muco virente obvolutis.

Hab. ad terram humoso-arenosam locis umbrosis in Britannia,
Fennia, Gallia, Italia, America boreali. — In speciminibus pata-
vinis: ascii cylindraceo-clavati, breve stipitati, apice rotundati, 180 ± 20 , p. sporif. 100 longa; paraphysibus filiformibus, guttulatis ob-
vallati, octospori. Sporidia disticha, ovata, 25 ± 16 , rugoso-stria-
tula, initio 1-guttata, hyalina, tandem pure fuliginea (non viola-
cea). Ascomata olivaceo-virentia. — * **microspermus** Speg. F. Arg.
IV, n. 220: ascomatibus viridibus, breviter turbinatis, disco primo
hemisphærico, dein patellari-applanato, glaberrimo (2-5 mm. diam.);
ascis cylindraceis vel cylindraceo-clavatis, $140-150 \pm 10-20$, apice
obtuse truncato-rotundatis, deorsum attenuato-stipitatis; parathy-

sibus septulatis longioribus apice vix incrassato-nodulosis obval-latis; sporidiis ellipticis, utrinque obtuse rotundatis vel subtrun-cato-rotundatis, 17-18 = 8-9, monostichis vel irregulariter distichis, primo hyalinis, dein dilute roseo-fuligineis, longitudinaliter dense striatulis. Ad terram pingue uliginosam in nemorosis prope la Recoleta Reip. Argentinæ.

26. **Ascobolus carbonarius** Karst. M. F. I, p. 76. — Cupulis gregariis 2156 sessilibus, concavis, demum pressione mutua difformibus, viridi-luteis mox brunneis furfuraceis, hypothecio pallido, 2-4 mm. lat.; ascis cylindraceo-clavatis, jodo non tinctis, 22 μ . cr., 8-sporis; sporidiis ellipsoideis, reticulatim insculptis, monostichis, violaceo-fuscis, 18-26 = 12-16; paraphysibus ramosis, 3 μ . cr.

Hab. ad terram in carbonariis circa Mustiala Fenniæ.

27. **Ascobolus pusillus** Boud. in Bull. Soc. Bot. Fr. 1877, pag. 310, 2157 t. IV, f. 7. — Perexiguus, carnosulus, hemisphæricus, dein plano-convexus, immarginatus, glaber, flavo-purpurascens, extus pallidior; margine saturatiore; ascis amplis, sat brevibus, clavatis, operculatim dehiscentibus; sporidiis distichis, ovoideis, utrinque acutiusculis, 10-11 = 6-8, violaceis, tenuiter reticulatis; paraphysibus filiformibus, clavula oblonga dilute colorata terminatis.

Hab. in terra adusta et cuniculis fœdata, Montmorency Galliæ.

28. **Ascobolus psittacinus** Quél. Enchir. p. 293, *Peziza* Quél. Jura II, 2158 p. 295, t. V, fig. 9. — Ascomate cyathiformi, ampio, 1 cm. lat., concavo-plano, granulato stipiteque tereti, brevi, amœne viridibus dein sulphureis, disco olivaceo-fusco; ascis cylindraceo-clavatis; sporidiis monostichis, ellipsoideo-oblongis, amethysteis v. lilacineis, dein brunneis, rivulosis.

Hab. ad humum in ollis floralibus Jura Galliæ.

29. **Ascobolus foliicola** B. et Br. Fung. of Ceyl. n. 972. — Cupulis 2159 extus luteo-ferrugineis; hymenio demum undulato, spadiceo; spo-ridiis pallide lilacinis longitudinaliter rugosis.

Hab. in foliis putridis in insula Ceylon. — Episporium senio in globulos minutos diffinditur.

30. **Ascobolus atro-fuscus** Phill. et Plow. in Grevillea II, p. 186, 2160 t. 24, f. 1, *Ascobolus viridis* Boud. Ascob. p. 27, t. V, f. 4 nec Curr., *Ascobolus carbonicola* Boud. Bull. Soc. Bot. Fr. 1877, p. 310. — Sessilis, confertus v. sparsus, nigro-brunneus, concavus, dein planus, margine crenulato, extus furfuraceus; ascis clavatis, octosporis; sporidiis late ellipsoideis, obtusis, 22 = 13, hyalinis, dein

purpureis, demum brunneis, episporio granulato v. verrucoso; paraphysibus linearibus, subinde furcatis.

Hab. in areis carbonariis vel in carbone ipso in Britannia et Gallia. — Ascoma 2-6 mm. lat. Variat luteo-virens, ferrugineo-furfuraceus vel fulvus vel nigro-purpurascens. — Var. **pruinosus** Boud. l. c. t. V, fig. 5: minutissime pruinosus, opace fulvo-brunneus, carne luteo-virente, margine tantum dentato. Iisdem locis in Gallia.

31. **Ascobolus Archeri** Berk. Fl. Tasm. p. 276. — Cupula undulata, 2161 sessili vino-fusca; ascis clavatis; sporidiis amethysteis, eleganter granulatis, 12-13 μ . longis, dein usque 16-17 μ .

Hab. in carbone in Tasmania (ARCHER). — Ascoma 3 mm. lat.

32. **Ascobolus fimiputris** Quél. Enchir. p. 293, *Peziza fimiputris* 2162 Quél. Jura II, p. 397, t. V, f. 30. — Ascomate sessili, majusculo, patellari, farinoso, fuscello, 5 mm. lat., margine crenulato, furfuraceo-albo; disco spadiceo; sporidiis ellipsoideo-oblongis, obscure lilacinis vix rivulosis.

Hab. ad humum hortorum, Jura Galliæ.

c) Species phytogenæ.

33. **Ascobolus lignatilis** A. S. Lus. p. 347, t. 6, f. 6, Fr. Syst. Myc. II, 2163 p. 162, Boud. Ascob. p. 25, t. V, fig. 1. — Sordide luteus, extus albido-furfuraceus, stipite brevi crassiusculo in cupulam planam marginatam dilatato; ascis clavatis, mamillato-operculatis, 8-sporis; sporidiis anguste ellipsoideis, minoribus, longitudinaliter striatis parceque reticulatis, fuligineis.

Hab. ad ligna pinea fabrefacta subterranea (aqua cloacina aliquando male inquinata) Niskiæ in Lusatia. — Reliquis major, gregarius, furfure cum *Ascob. furfuraceo* congruens. Stipes 3-4 mm. fere longus et crassus, obconicus, sursum sensim dilatatus in cupulam planam v. concaviusculam, 8-10 mm. latam, ascis creberimis nigro-punctatam. Color constans, e sordide luteo-subvirens (virore fere obsoleto).

34. **Ascobolus denudatus** Fr. Syst. Myc. II, pag. 164, Boud. Ascob. 2164 p. 26, t. V, f. 3. — Luteo-viridis, glaber, cupula turbinata sub-stipitata, disco plano; ascis amplis, stipitatis, 8-sporis; sporidiis subdistichis, oblongo-ovatis, æquilateralibus, demum conglutinatis, 16-18 = 8, episporio subtilissime longitudinaliter striato et parce reticulato amœne violaceo, demum opaco-fusco; paraphysibus filiformibus, furcatis, apice clavatis.

Hab. ad asseres vetustos pineos in latere ad terram vergente in Suecia, Britannia, Germania. — Gregarius. Stipes brevis, sursum dilatatus in cupulam planam, marginatam, luteo-viridem, extus nudam et glabram, 2 mm. latam nec ultra. Variat stipite æquali.

35. **Ascobolus Crouani** Boud. Ascob. p. 26, tab. V, f. 2. — Sessilis, 2165 luteo-virens, globosus, dein turbinatus, plus minusve expansus, 2-5 mm. lat., fibrillis albis adfixus, extus conspicue fulvo-furfuraceus, vetustus brunnescens, margine furfuraceo-denticulato, disco concavo dein applanato lucido; ascis clavatis, basi attenuatis, octosporis; sporidiis ellipsoideis, subminutis, violaceis, denique brunneis, episporio longitudinaliter creberrime rimuloso; paraphysibus gracilibus septatis, apice piriformibus ascisque gelatina luteo-virente obvolutis.

Hab. inter folia putrida ad basim trunci *Syringæ vulgaris*, Montmorency Galliæ. — *Ascob. lignatili* dimidio minor.

36. **Ascobolus incolor** Quél. X Suppl. p. 11, t. IX, f. 17. — Ascomatibus turbinatis, 200-300 μ . diam., teneris, crassis, glabris, albescentibus v. opalinis, margine crenulato, furfuraceo, griseo-carneo; hymenio concavo, dein plano, velo fugaci tenui prædicto, dein violaceo-punctulato; sporidiis ellipsoideo-oblongis, 25 μ . longis, venoso-reticulatis, violaceis.

Hab. in truncis *Brassicæ* putrescentibus Jura. — *Habitu Ombrophilam Clavum* in mentem revocat.

37. **Ascobolus Thwaitesii** B. et Br. Fung. of Ceyl. n. 970. — Carninus, extus albo-furfuraceus; sporidiis ellipticis echinulatis, 25 \approx 12.

Hab. in ligno putri, Central province, Ceylon. — Ab *A. lignatili* diversus.

38. **Ascobolus angulisperus** Boud. Bull. Soc. Bot. 1881, p. 92, t. II, 2168 f. 2. — Majusculus, tenuiter furfuraceus, subolivaceus hemisphaericus v. depresso, 5-7 mm. diam., extus granulis cupulâ saturatioribus conspersus; disco ascis punctato; ascis clavatis, octosporis, operculo subpapillato, 200-250 \approx 20; sporidiis ellipsoideis, violaceo-brunneis, 15-18 \approx 7-8, episporio depresso anguloso-verrucoso atque rimoso; paraphysibus filiformibus, septatis, clava 5-7 μ . crassa terminatis.

Hab. in scobe lignea putrescente, Montmorency Galliæ.

39. **Ascobolus cenangioides** Ces. Myc. Born. p. 11. — Cupulis gregariis, 1-1½ mm. latis, margine involuto, extus ferrugineis, disco brunneo, glabris; sporidiis fere opacis, nigrescentibus, levibus visis,

oblongis, e latere paullisper curvulis, obtusissimis, 9-10 \approx 3-4; paraphysibus numerosissimis filiformibus vix sursum incrassatis, obscure septatis.

Hab. parasiticus in *Xylaria cupressiformi*, Sarawak Bornei (BECCARI).

40. **Ascobolus Constantini** Roll. Bull. Soc. Mycol. IV, p. 56, t. XV, 2170 f. 1. — Stipitatus, carnosus; disco tenui, purpureo-brunneo, luteo-marginato, saepe umbilicato, subtus ochraceo; stipite sat brevi, etiam ochraceo, sursum dilatato; ascis cylindricis, breviter pedicellatis aut ad basim attenuatis, 100 \approx 15, 8-sporis; sporidiis elliptico-fusiformibus, 15 \approx 8, episporio violaceo longitrorsum rimoso rimis tenuibus; paraphysibus numerosis, gracillimis, hyalinis, simplicibus ascisque gelatina sulphurea immersis.

Hab. ad corticem putrescentem Rubi prope Parisios Galliæ.

II. *Dasyobolus* Sacc. Ascomata pilosella.

41. **Ascobolus immersus** Pers. Obs. I, p. 35, t. 4, f. 7, Sacc. Mich. 2171 I, p. 234, Nees Syst. f. 297, Fr. Syst. Myc. II, pag. 164, Boud. Ascob. pag. 36, t. VIII, fig. 17, *Ascob. gigasporus* De Not. Disc. p. 360. — Gregarius v. sparsus, subimmersus aut emersus, teretipiriformis dein planiusculus v. convexus, sordide pallido-virens v. fuscescens, circiter 0,5-0,9 mm. lat., extus pilis minutissimis continuis, hyalinis simplicibus v. fasciculatis obsitus; ascis octosporis, rarius 2-4-6-sporis, cylindraceo-ellipsoideis, basi breviter attenuatis, 180-250 \approx 60-95, apice truncatis, jodo dilute cœrulescentibus; sporidiis oblongo-ellipsoideis, longitrorsum remote rimulosis laxeque reticulatis, violaceo-fuscis, 60-80 \approx 30-48; paraphysibus æqualibus, inarticulatis, gracilibus.

Hab. supra stercus vaccinum et equinum in Suecia, Dania, Britannia, Germania, Belgio, Hungaria, Italia. — ***A. macrosporus** Cr. in Ann. Sc. nat. 4, VIII, p. 173, t. 4, B, Cav. Mat. Myc. Lomb. p. 8. A typo distinguitur ascis minoribus, apice rotundatis; sporidiis ellipsoideis, minoribus 40-50 μ . long. Ibidem. ridiisque maximis facillime dignoscitur.

42. **Ascobolus porphyrosporus** (Hedw.) Fr. Syst. Myc. II, pag. 163, 2172 Boud. Ascob. p. 36, Fuck. Symb. myc. App. II, p. 57, *Octospora porphyrospora* Hedw. Musc. frond. II, p. 25, t. 7, f. A. — Sessilis, subcylindricus, luride viridis, externe hirtulus; ascis clavato-saccatis, breviter stipitatis, 8-sporis, 240 \approx 20-30; sporidiis plerumque in asci superiori parte distiche conglomeratis, nec non

sæpe liberis, oblongo-ovatis, rectis, utrinque obtusis, monoplastis, episporio longitudinaliter distincte striato, pulcherrime violaceo, 22-24 = 10-12 (Fuck.).

Hab. in straminibus fimi veteris equini in Suecia, Gallia, Germania. — Gregarius, minor; junior nunc conoideus, nunc globosus; ille æqualiter in formam cylindricam, hic superne in cupulam rotundatam abit; sub lente extus hirtulus, deorsum vero glaber dilutior. Color sec. Fuck. initio albus dein diaphane aureus, demum brunneus.

43. **Ascobolus brunneus** Cooke in Hedwigia VI, 1867, p. 154. — 217:

Sessilis, minutus, globosus, dein depresso, extus pilosus, pallide ochraceus; ascis late clavatis, brevibus, octosporis; sporidiis ellipsoideis, initio hyalinis et granulosis, demum brunneis, 25-35 = 15; paraphysibus filiformibus.

Hab. in fimo vaccino, Shere Surrey Britanniæ.

III. Species minus notæ.

44. **Ascobolus? microscopicus** (Wallr.) Sacc. *Peziza microscopicica* 2174

Wallr. n. 2543. — Ascomatibus glabris, microscopicis, punctiformibus exilissimis, carnosulis, pallide flaventibus, dein discum planiusculum cerinum dilutius marginatum recludentibus, basi pilis arachnoideis suffultis.

Hab. ad fimum vaccinum in Thuringia in consortio. — *Lasioboli pilosi*, cuius forsitan initia incondita.

45. **Ascobolus Tetricum** Carestia in Rabenh. F. E. n. 1236 (non vidi) 2175 et in Bot. Zeit. 1869, p. 431 (nomen).

Hab. in stercore Tetraonis Tetricis in Italia. — Verisimiliter una alterave ex speciebus in fimo Tetraonum nascentibus et hic descriptis.

SACCOBOLUS Boud. Ascob. pag. 38 [1869] (Etym. *saccos* saccus et *ballo* ejicio). — Ascomata carnosula, sessilia, glabra, rarissime pilosella, plano-convexa. Discus ascis mox nitide nigro-punctatus. Ascii lati, breves, octospori, paraphysati. Sporidia ellipsoideo-oblonga, colorata, levia vel asperula, ad apicem ascii sacculo communi inclusa.

I. *Eu-Saccobolus*. Ascomata glabra.

1. **Saccobolus Kerverni** (Cr.) Boud. Ascob. p. 39, t. VIII, f. 19, E. 2176 Ch. Hans. F. fimb. dan. t. VI, f. 22, Ascob. *Kerverni* Cr. Finist. p. 57 et Ann. Sc. nat. 1858, pag. 193, t. XIII, f. B. — Minutus,

flavo-aureus, sessilis, hemisphæricus v. subcylindricus, disco piano vel convexo; ascis ovato-cylindraceis, sursum obtuse rotundatis, deorsum brevissime crassiuscule stipitatis, 85-105 \approx 26-35, in basidiis cylindraceis, apice subincrassatis, 45-50 \approx 6-8, acrogenis; paraphysibus filiformibus, 5 cr., granulosis viridulis obvallatis; glo-merulis sporidiorum sacculo inclusis, cylindraceo-ellipsoideis, 45-50 \approx 20; sporidiis subnavicularibus, 18-24 \approx 10-12, utrinque rotundato-truncatulis, primo violaceis, dein amœne vinoso-fuligineis.

Hab. in fimo vaccino et equino in Gallia, Italia, Britannia, Hungaria, Germania, Amer. austr.

2. **Saccobolus violascens** Boud. Ascob. p. 40, t. VIII, fig. 19, Gillet 2177
Discom. cum. ic. Phill. Disc. p. 296, t. IX, f. 55. — Sparsus v. aggregatus, minutus, 1-2 mm. lat., rotundatus v. plus minusve flexuosus, glaber, nitidus subgelatinosus, cinereo-violaceus, disco convexo, vix pallidore; ascis late clavatis, 66 \approx 15, basi attenuatis, operculo plano dehiscentibus, octosporis; sporidiis ellipsoideis, subacutis, subinde obliquis, hyalinis, demum cyaneo-violaceis vel atro-violaceis, levibus, 15-17 \approx 7-9, sacculo communi hyalino inclusis; paraphysibus septatis, clava violacea terminatis.

Hab. in fimo vaccino et Cuniculorum in Gallia, Britannia, Germania et Hungaria.

3. **Saccobolus depauperatus** (Berk. et Br.) Rehm Ascom. n. 90, Asco- 2178
bolus depauperatus B. et Br. Ann. Nat. Hist. n. 1084, tab. XIV, f. 6, Asc. Kerverni Karst. M. F. I, p. 78, nec Cr. — Minutus, planus, pallidus, dein vinosus; ascis brevibus, ovato-clavatis, basi abrupte breviter attenuato-stipitatis, octosporis, 40 \approx 18; sporidiis obtuse fusoideis, luride violaceis, levibus, 10-12 \approx 5-6,5, sacculo communi hyalino inclusis; paraphysibus filiformibus, septatis, tenuiter apice incrassatis.

Hab. in stercore ovino, equino et cervino in Britannia et Transylvania. — Ascoma 250 μ . lat., flavidum, mox vinosum.

4. **Saccobolus versicolor** Karst. Rev. mon. p. 123, Asc. versicolor 2179
Karst. M. F. I, p. 79. — Gregarius, sessilis, nudus, immarginatus, convexellus, primum incarnato-pallidus, dein rufo-carneus, demum fuligineus, 0,4-0,9 mm. lat.; ascis oblongatis, basi breviter attenuatis, 70-86 \approx 20-22, jodo cœrulescentibus, 8-sporis; sporidiis fusideo-oblängato-ellipsoideis, levibus, violaceo-fuscescentibus, 17-20 \approx 7-9 in sacculum oblongum apice ascorum collectis.

Hab. supra stercus vaccinum in Fennia australi. — An, teste Rehm, satis distinctus a *S. violascente*?

5. **Saccobolus obscurus** Cooke in Grev. IV, p. 112. — Subgregarius, 2180
subinde dense congestus, fuligineus vel atro-brunneus, disco con-
vexo, pallidiore; ascis clavatis, octosporis; sporidiis ellipsoideis, vio-
laceis, dein brunneis, rugosis, 12-14 = 7, sacculo communi hyalino
inclusis; paraphysibus filiformibus simplicibus.

Hab. in tela putrescente uda, Forden Montgomeryshire Britan-
niæ (COOKE). — Ascoma 200-900 μ . diam.

6. **Saccobolus dilutellus** (Fuck.) Sacc. *Ascobolus dilutellus* Fuck. in 2181
Hedw. 1866, p. 4, f. 7, a, b. c. — Ascomatibus gregariis sessi-
libus, 0,5 mm. lat., planis, diaphanis, dilutissime aquoso-violaceis,
disco paullulo obscuro, subtilissime punctatis; ascis stipitatis,
oblongo-ovatis, curvatis, 8-sporis; sporidiis in asco amplo semper
dense conglomeratis, glomerulis suborbicularibus, 24 μ . diam., ova-
tis, dilute violaceis, dein fuscis, 14 = 8; paraphysibus simplicibus,
plerumque clavatis, 1-3-septatis.

Hab. in fimo putri canino, Oestrich Rhenogoviæ.

7. **Saccobolus globulifer** Boud. Ascob. p. 42, t. IX, f. 21. — Hemi- 2182
sphæricus, glaber, vix nitidus, albus, dein pallide cinereo-purpu-
rascens, disco subplano; ascis crassis, brevibus, apice truncatis,
basi attenuatis, operculo plano dehiscentibus, octosporis; sporidiis
oblongo-ovatis, utrinque subacutis, pressione trigonis, maturis in
globulum rotundatum aggregatis, roseis in cinereo-violaceum ver-
gentibus, episporio verrucis minutis, præsertim ad margines prædi-
to; paraphysibus bacillaribus, septulatis, hyalinis.

Hab. in fimo Cuniculorum, Ecouen pr. Paris Galliæ.

8. **Saccobolus neglectus** Boud. Ascob. p. 41, t. IX, f. 20. — Pere- 2183
xiguus, globosus, glaber, subnitidus, albus vel dein cinereus vel
cinereo-purpureus, basi pallidior; disco fulvo, subinde vero ascis
exsertis nigricante; ascis minutis, latis, basi attenuatis, operculo
plano dehiscentibus, octosporis; sporidiis ellipsoideis, subacuminatis,
levibus, roseolis dein cinereo-violaceis nigrisque, 15-19 = 6-8, sa-
culo communi hyalino inclusis; paraphysibus simplicibus vel fur-
catis, apice vix incrassatis, hyalinis, septatis, muco parum conspi-
cuo obductis.

Hab. in fimo Cuniculorum, equino, asinino, vaccino
et ovino in Gallia et Britannia. — Variat albidus et cinereus
disco subpurpureo.

9. **Saccobolus Hansenianus** Speg. Mich. I, p. 234. — Ascomatibus 2184
primo conoideis, durissimis, flavo-viridulis, vertice atro-vinoso, dein
applanatis undique flavo-viridulis sed disco ob ascos exsiliens brun-

neo-punctato; ascis amplis saccatis, elliptico-fusoideis, deorsum brevissime stipitatis, apice truncatis, 180-210 \approx 67-75, basidiis cylindraceis, vix clavulatis, 40-45 \approx 10-12, suffultis, paraphysibus filiformibus viridulis obvallatis; glomerulis sporidiorum elliptico-ovati, 85-90 \approx 30, sacculo crassissimo inclusis; sporidiis ovato-ellipsoideis, inæquilateralibus, utrinque rotundato-truncatis, 35-40 \approx 25, primo opace violaceis, dein intense fuligineis.

Hab. in fimo vaccino et equino in umbrosis herbidis, Conigliano Ital. bor.

10. **Saccobolus Boudieri** Oud. Hedw. 1882, p. 166 et Bijd. IX, p. 23. 2185
 — Ascomatibus minimis, glabris, levibus; ascis clavatis, 100 \approx 25; sporidiis singulis violaceis, in glomerulum ovalem vel oblongo-ovalem, 42 \approx 18, conglutinatis, pressione trigonis, episporio verrucis minutissimis exasperato, 16 \approx 7.

Hab. in fimo Amsterdam Nederlandiæ (VAN LEDDEN). — Color ascomatis non indicatus.

II. *Eriobolus* Sacc. Ascomata albo-pilosa.

11. **Saccobolus cæsariatus** Renny in Phill. Disc. p. 297. — Minutus 2186 hemisphæricus, albus, pilis hyalinis, septatis, tenuibus, flexuosis ob-situs; disco convexo, ascis exsertis papillato; ascis late clavatis; sporidiis ellipsoideis, levibus, purpureis, sacculo communi hyalino inclusis; paraphysibus filiformibus, obtusiusculis, simplicibus vel ramosis.

Hab. in fimo? in Britannia. — *Lachnellam* quamdam candidam imitatur.

Sect. 2. *Hyalosporæ* Sacc.

- CUBONIA** Sacc. nov. gen. (Etym. a cl. prof. J. CUBONI, Instituti phytoto-pathologici romani directore), *Lasioboli* et *Ascophani* spec. Auct. — Ascomata sessilia hemisphærico-obconica, glabra vel pilosa, disco plano, ascis papillato. Asci clavati vel subpiriformes, octospori, paraphysati. Sporidia perfecte sphærica, asperula, hyalina.

1. **Cubonia brachyasca** (March.) Sacc. *Lasiobolus brachyascus* March. 2187 Fung. copr. Belg. p. 41. — Sparsa, primo obconica deinde applanata, sessilis, alba, extus dense longeque pilosa, pilis inferne incrassatis, 200-280 \approx 6-8, continuis vel basi vix septatis, hyalinis vestita, 120-200 μ . diam., margine obsoleto; disco plano; ascis crystallinis valde prominulis eximie papillato; paraphysibus gra-

cilibus, hyalinis, simplicibus vel 2-4-divisis, valde arcuatis, inferne articulatis, in fragmenta bacillaria abeuntibus et ascis vix æquantibus, apice late rotundatis, inferne breviter stipitatis, 40-45 = 19,5-21,5, jodo non cœrulecentibus; sporidiis subdistichis, perfecte globosis, hyalinis, 8,8-9,2 μ . diam. minute asperulatis.

Hab. in fimo canino, socio *Ascobolo aeruginoso*, Werbomont (Liège) Belgii.

2. **Cubonia Boudieri** (Renn.) Sacc. *Ascoph. Boudieri* Renny et Phill. 2188 Disc. p. 304. — Minuta, sessilis, hemisphærica, glabra, alba; disco plano, papillato; ascis late cylindraceo-clavatis, octosporis; sporidiis globosis, asperis, hyalinis; paraphysibus filiformibus obtusis.

Hab. in fimo cuniculorum, Hereford Britanniæ (RENNY).

ASCOPHANUS Boud. Ascob. p. 51 (1869) [Etym. *ascos* ascus et *phaino* appareo], *Ascoboli* et *Pezizæ* spec. Auct. — Ascomata carnosula, sessilia, scutellata, plana v. convexula, glabra v. rugosa (non setosa). Asci lati, 4-8-spori, operculo apice dehiscentes, e disco exsilientes, varie paraphysati. Sporidia ellipsoidea, subhyalina, continua, levia vel vix granulata.

I. Species fimicolæ.

1. **Ascophanus argenteus** (Curr.) Boud. Ascob. pag. 55, t. XI, f. 32, 2189 *Ascobolus* Curr. in B. et Br. Ann. N. H. n. 1088, t. XVII, f. 32. — Gregarius, minutissimus, oculo inermi vix visibilis, subpiriformis, argenteo-albus; ascis oblongo-ovoideis, basi acutatis, octosporis; sporidiis ellipsoideis, hyalinis, 12 = 7-8; paraphysibus apice clavatis.

Hab. in fimo vaccino, Elthan Britanniæ.

2. **Ascophanus lacteus** Cooke et Phill. Grev. V, p. 119, Disc. p. 306. 2190 — Sparsus, initio obconicus dein expansus, concaviusculus, demum convexus, lacteus, glaber; ascis clavatis, octosporis; sporidiis subdistichis, ellipsoideis, hyalinis, levibus, 10 = 5; paraphysibus filiformibus.

Hab. in fimo vaccino, Shrewsbury Britanniæ. — Ascoma 1- $\frac{1}{5}$ mm. lat.

3. **Ascophanus microsporus** (B. et Br.) Phill. Disc. p. 307, *Ascobolus* B. et Br. Ann. N. H. n. 1087, t. XVI, f. 28, *Asc. Coemansii* Boud. Ascob. p. 54. t. X, f. 30. — Exiguus, sessilis, sordide albus vel flavidо-brunneus, depresso; ascis late clavatis sæpe stipite tenui obliquo auctis, 8-sporis; sporidiis ellipsoideis, hyalinis, levi-

bus, 7-8 = 3,5; paraphysibus filiformibus apice eximie capitatis, endochromate flavo-viridi.

Hab. in fimo vaccino et ovino in Gallia et Britannia. — Ascoma $\frac{1}{2}$ -1 mm. lat. Habitus fere *Ascoph. granuliformis* sed fructificatio diversa. Sec. B. et Br. sporidia tandem violacea fiunt, quod Boudier haud observavit.

4. **Ascophanus incanus** (Phill.) Sacc. *Ascobolus* Phill. Grev. V, p. 117, 2192 tab. 88, f. 10. — Sparsus, turbinatus, dein planus, disco depresso incano-griseus, margine membranaceo dein evanescente; ascis late clavatis, 8-sporis; sporidiis ovoideis, hyalinis, 29 = 14; paraphysibus linearibus.

Hab. in fimo vaccino, Blue Canon Sierra Nevada Californiae. — Sporidia interdum utrinque papillata; operculum ascorum evidens.

5. **Ascophanus Aurora** (Cr.) Boud. *Ascob.* p. 58, tab. XI, f. 36, *Peziza Aurora* Cr. Fin. p. 53. — Sparsus, minutus, aurantio-ruber nitidulus, applanatus, disco distincte papillato; ascis clavatis, octosporis; sporidiis ellipsoideis, hyalinis, levibus, 10 = 6,5; paraphysibus filiformibus, simplicibus vel divisis, curvis, aurantio-granulosis.

Hab. in fimo vaccino in Gallia, Britannia et in Agro tridentino. — In exemplaribus tridentinis ascii sunt oblongo-cylindracei 80 = 16; sporidia disticha oblongo-ellipsoidea, 1-guttata, hyalina, 14-15 = 7-8; paraphyses filiformes sursum furcatæ.

6. **Ascophanus nitidus** (Fuck.) Sacc. (nec Spög.), *Ascobolus* Fuck. 2194 Hedw. 1866, p. 4, f. 4 a, b, c. — Ascomatibus gregariis, seminis papaveris magnitudine, lentiformibus, omnino levibus, nitidis, pulcherrime miniatis siccitate excavatis, *Nectriæ Pezizæ* similibus, colore immutato diaphanis; ascis oblongis, brevissime oblique stipitatis, octosporis; sporidiis ovatis, hyalinis; paraphysibus simplicibus, filiformibus, apice curvatis, multiguttulatis, miniatis.

Hab. ad fimum equinum in silva Estrich Rhenogoviæ.

7. **Ascophanus cinnabarinus** Spög. Mich. II, p. 258. — Primo cylindraceo-conicus melleo-fulvus, dein cupuliformis, vividissime cinnabarinus, disco saturiore $\frac{1}{4}$ - $\frac{1}{5}$ mm. diam.; ascis cylindraceis longissime pedicellatis, octosporis 150 = 8-9; paraphysibus apice crasse clavatis 10 μ . cr., atque intense rubro-fuligineis, jodi ope opace cœruleoscentibus; sporidiis monostichis, hyalinis, ellipticis, eguttulatis, 10-12 = 5.

Hab. in fimo vaccino in alpinis Cadore Ital. bor., ubi, etsi

minutus ob vivacitatem coloris, facilime detegitur, socio *Ascoph.*
Holmskioldii E. C. Hans.

8. *Ascophanus amethysteus* Quél. Esp. nouv. Champ. Fr. 1878, p. 291. 2196

— Ascomatibus globulo-lentiformibus, teneris, glabrescentibus,
sub lente granulosis, translucidis, hyalino-roseis; sporidiis ellipso-
deis, 12 μ . longis, hyalino-roseis.

Hab. in fimo in silvis Juræ.

9. *Ascophanus vicinus* Boud. Ascob. p. 56, t. XI, f. 33. — Sparsus, 2197

raro aggregatus, $\frac{1}{2}$ -1 mm. lat. ochraceo-carneus, convexulus v.
applanatus; disco minute papillato; ascis oblongis, amplis, basi at-
tenuatis, octosporis; sporidiis ellipsoideis utrinque rotundatis, hya-
linis, 30-32 \approx 20-22; paraphysibus apice clavatis, septatis, hya-
linis.

Hab. in fimo vaccino in Gallia, Belgio et America boreali.

10. *Ascophanus Holmskjoldii* Hans. F. fim. dan. p. 12, t. VI, f. 1-8. 2198

— Ascomatibus cinereo-rubentibus, furfurellis, sessilibus, hemisphæ-
ricis vel subcylindraceis, disco convexo papillato, $\frac{1}{2}$ -1 mm. lat.;
ascis elongato-fusoideis, p. spor. 180-200 \approx 30-45, stipite 100 μ . long.,
paraphysibus filiformibus jodo cœrulescentibus obvallatis, octospo-
ris; sporidiis distichis oblongo-ellipsoideis 30-36 \approx 13-16, minute
asperulis, utrinque appendiculâ rotundatâ auctis stratoque mucoso
obvolutis, flavid-guttulatis, hyalinis.

Hab. in fimo vaccino et damarum in Dania, Belgio, Italia.

— Sec. Hans. loc. cit. ex appendiculis setulæ fasciculatæ oriun-
tur, quæ vero in sporidiis junioribus tantum observandæ sunt.

11. *Ascophanus subgranuliformis* Rehm in Voss Mater. Krains V, 2199

p. 224, fig. 7 a, b. — Gregarius pallide luteus, levis, $\frac{1}{2}$ -1 mm.
lat., ascis cylindraceis, breve stipitatis; sporidiis monostichis, ob-
longo-ellipsoideis, 30-33 \approx 12-14 (ex icon), hyalinis, levibus; pa-
raphysibus filiformibus, apice capitatis, non curvatis.

Hab. in fimo vaccino exsiccato in fagetis Carnioliae, 1500 m.
alt. — Ab *Asc. granuliformi* Boud. differt ascis cylindricis, pa-
raphysibus non curvatis.

12. *Ascophanus granuliformis* (Cr.) Boud. Ascob. p. 55, t. X, f. 31, 2200

Pat. Tab. f. 173, *Ascobolus* Cr. Fin. p. 56, *Peziza* Karst. M. F.
I, p. 50. — Sparsus sphæricus, flavid-ochraceus, pellucidus, cen-
tro opacus, levis, disco planiusculo, papillato, extus glaber; ascis
brevibus et latis, 90-115 \approx 12-18, octosporis; sporidiis ovoideis, ob-
tusis, uniguttatis, levibus, 12-15 \approx 7-10; paraphysibus filiformibus
apice subincrassatis.

Hab. in fimo vaccino, equino, rarius cunicolorum in Gallia, Britannia, Germania, Belgio, Italia, Fennia, Amer. merid. — Ascoma $\frac{1}{2}$ -1 mm. diam.

13. **Ascophanus ochraceus** (Cr.) Boud. *Ascob.* p. 57, tab. V, fig. 34, 2201

Ascobolus Cr. Fin. p. 57, *Ascoph. nitidus* Speg. Mich. I, p. 235, nec Fuck. — Glaber, ochraceus, minutus, $\frac{1}{2}$ -1 mm. lat., disco convexo, ascis minute punctato; ascis amplis, octosporis, cylindra-CEO-subfusoideis, apice subattenuatis; sporidiis ovato-oblongis, levibus, hyalinis vel luteolis, 16-17 \approx 8-9; paraphysibus simplicibus raro e basi bi- trifidis sursum curvatis et subincrassatis.

Hab. in fimo vaccino, equino, ovino et leporino in Gallia, Britannia, Italia et Belgio. — Variat ascomate pallido et fere albo.

14. **Ascophanus minutellus** Karst. Rev. Monogr. p. 121, *Peziza mi-* 2202

nutella Karst, M. Fenn. I, p. 51. — Sparsus, sessilis, immarginatus, planus, testaceo-pallidus aut albidus, 0,2-0,3; ascis fusoideo-elongatis, 80-100 \approx 16, octosporis; sporidiis mono- vel distichis, ellipsoideis, eguttulatis, 15-18 \approx 8-9; paraphysibus filiformibus, simplicibus, inarticulatis, 1,5-2 μ . cr., apice clava subovoidea, obliqua, 4-5 μ . crassa.

Hab. supra fimum ovinum vetustum in Mustiala Fenniae.

15. **Ascophanus vilis** Karst. et Starb. Rev. Myc. 1887, p. 150 et Symb. 2203

XXIII, p. 3. — Sessilis, immarginatus, planiusculus, testaceo-pallescens, siccus fuscescens, 0,2 mm. lat.; ascis cylindraceis 120 \approx 10-12, 8-sporis; sporidiis monostichis, ellipsoideis, eguttulatis, 15-18 \approx 8-9; paraphysibus filiformibus, conglutinatis, apice incurvatis, vix incrassatis.

Hab. in fimo equino ad Mustiala Fenniae (K. STARBAECK). — Ab *Asc. minutello* sporidiis monostichis et paraphysibus haud clavatis distinctus.

16. **Ascophanus cinereus** (Cr.) Boud. *Ascob.* pag. 59, t. XI, f. 37, 2204

Ascobolus Cr. Finist. p. 56, *Peziza* Karst. M. F. I, p. 59. — Gregarius, sessilis, crassiusculus, carnosus, disco plano vel convexiusculo, ab initio pallidus, dein cinereus vel fuliginosus, demum saepe nigricans, fere marginatus, interdum obliquus et irregularis, 1 mm. lat.; ascis cylindraceo-clavatis, basi tenuata stipitiformi, 120-180 \approx 16-25, jodo coerulecentibus, 8-sporis; sporidiis monostichis, cylindraceis vel ellipsoideis, eguttulatis, 14-23, plerumque 18-20 \approx 8-11; paraphysibus filiformibus, simplicibus.

Hab. supra fimum equinum in Gallia, Italia, Germania, Bri-

tannia, Belgio, Fennia. — Sec. Rehm sporidia vetusta brunnescunt et utrinque papillam hyalinam præbent.

17. **Ascophanus cinerellus** (Karst.) Speg. F. Arg. II, n. 97, *Pezizula* 2205
cinerella Karst. M. F. p. 51. — Sparsus vel subgregarius, sessilis, convexellus, margine flexuoso-repando (saltem in statu sicco), cinereo-pallidus circ. 0,3-0,4 mm. lat.; ascis cylindraceo-clavatis, 38-42 \approx 6-7, jodo non tinctis, 8-sporis; sporidiis mono- vel distichis, ellipsoideis vel sphæroideo-ellipsoideis, eguttulatis, 5-7 \approx 3-4; paraphysibus non visis.

Hab. supra sterlus vaccinum et equinum in Fennia, Belgio et Amer. australi. — Species facillime cognita, *Mollisiam fallentem* Karst. in memoriam revocans.

18. **Ascophanus antarcticus** Speg. Fung. Patag. p. 58. — Sparsus hemisphærico-cupulatus, minutus 300-350 μ . diam., ater, levissimus glaberrimus, disco concaviusculo non vel vix pallidore, carnosulo-subcarbonaceus, contextu excipuli dense minutissimeque merenchymatico, olivaceo; ascis cylindraceis, subtorulosis, apice obtuse rotundatis, postice in stipitem crassum ac breviusculum productis, 65-70 \approx 7-8, octosporis, a paraphysatis, muco fuscescente tenacissimo immersis; sporidiis recte vel oblique distichis, ellipticis, 10-11 \approx 5-6,5, utrinque obtusis, hyalinis non vel grosse 1-guttatis.

Hab. in stercore vetusto Rhei Darwinii in altiplanite, Meseta vocata, secus Rio S. Cruz Patagoniæ. — Jodi vis nulla.

19. **Ascophanus punctiformis** (Karst.) Sacc. *Ascobolus* Karst. F. Fenn. exs. n. 655, *Pezizella polyspora* * *punctiformis* Karst. M. F. I, p. 82. — *Rypar. polysporo* extus similis sed ascis octosporis, oblongato-clavatis, 35-50 \approx 10-14; sporidiis ellipsoideis 8-10 \approx 4-5, hyalinis.

Hab. in fimo equino prope Abo et Mustiala Fenniæ.

20. **Ascophanus fallax** (Auersw.) Sacc. *Ascobolus fallax* Auersw. Hedw. 1868, p. 52, *Pezizula crustacea* v. *fallax* (Auersw.) Karst. M. F. I, p. 81. — Microscopicus, depresso-hemisphæricus, (ut videtur) immarginatus, lividus vel e luteo fuscescens, glaber; ascis semiprojectis ut punctulis albis inspersus; ascis parvis, 30 \approx 8-10, clavatis 8-sporis; sporidiis ovalibus et omnino cum præcedentium congruentibus.

Hab. in stercore canino, Rosenthal Cassel Germaniæ et in Fennia. — Affinis *Ryp. crustaceo* sed octosporus.

21. **Ascophanus subfuscus** Boud. Ascob. p. 52, t. 10, f. 28. — Confertus, rarius sparsus, minutissimus, hemisphæricus, extus glaber,

brunneus, basi pallidior, siccus atro-brunneus, margine inæquilatero inflexo, disco plano, pallido vel pallide fulvo; ascis brevibus, late clavatis, octosporis; sporidiis oblongo-ellipsoideis vel subsulcatis, hyalinis, 10×5 ; paraphysibus late clavatis, apice brunneis.

Hab. in fimo canino, felino, vaccino et humano in Gallia, Britannia, Belgio, Germania. — *Ascoma* 100-200 μ . lat.

22. **Ascophanus Cesatii** (Carest.) Sacc. *Ascobolus* Car. in Rab. F. E. ²²¹⁰ ed. II, n. 976, *Pezizula Cesatii* (Carest.) Karst. M. F. I, p. 83. — Cupulis madore molliusculis, patellulatis, disco pallide carneo, extus fuscis; ascis clavulatis, curvulis, numerosis, 8-sporis, 32×8 ; sporidiis hyalinis, ovalibus vel subrotundatis, 8μ . long.; paraphysibus simplicibus, capillaribus, rigidulis.

Hab. supra fimum Tetraonis Tetricis in Italia boreali, Fennia, in stercore anserino in Beeren Eiland (sporid. ellipsoidea, $8-10 \times 5-6$).

23. **Ascophanus tetraonalis** Peck Bull. N. Y. Mus. 1887, II, p. 22. — ²²¹¹ Sessilis, 2-4 mm. lat., extus cinereus, margine subinde undulatoflexuoso, disco nigricans vel nigro-brunneus; ascis cylindraceis, apice truncatis; sporidiis monostichis ellipsoideis, levibus, hyalinis, $15-18 \times 7$.

Hab. in excrementis Tetraonis, Catskill Mount. Amer. bor. — *Ascoph. gallinaceo* subaffine.

24. **Ascophanus minutissimus** Boud. Ascob. pag. 53, t. 10, f. 29. — ²²¹² Omnia minutissimus, 100-200 μ . lat., sparsus, hemisphæricus v. subplanus, brunneo-fulvus, margine nigro, siccus totus nigricans; ascis tereti-clavatis, minoribus basi attenuatis, octosporis; sporidiis ovoido-fusoideis, hyalinis, $7-9 \times 4$; paraphysibus simplicibus, apice piriformi, luteo-virente.

Hab. in fimo equino et ovino, Montmorency Galliæ, in Britannia et Belgio.

25. **Ascophanus conformis** (Karst.) Sacc. *Pezizula conformis* Karst. ²²¹³ M. F. I, p. 83. — Confertus, fuscidulus, glaber, disco pallidiori planiusculo, 0,1-0,2 mm. lat.; ascis clavatis, $40-70 \times 10-12$, parte sporif. $26-28 \mu$. long., octosporis; sporidiis distichis, ellipsoideis, eguttulatis, incoloribus, $7-8 \times 4$, paraphysibus nondum visis.

Hab. ad fimum vaccinum in sphagnosis deustis, Pellinsuo prope Mustiala Fenniæ. — Teste Karst., est subspecies *Rypar. polyspori*, sed asci octospori.

26. **Ascophanus humosoides** Peck Bull. N. Y. St. Mus. n. 2, p. 22. ²²¹⁴ Minutus, vix 1 mm. lat., sessilis, sparsus vel gregarius, ex au-

rantio vinosus, disco plano vel convexulo, breve marginato; ascis brevibus cylindrico clavatis; sporidiis ellipsoideis, levibus, 20-25 = 12; paraphysibus filiformibus, sursum crassiusculis.

Hab. in excremento cuniculi cuiusdam silvatici, Catskill Mount Was. Amer. bor. — Ascoma filamentis hyalinis paucis matrici affixum.

27. **Ascophanus Oudemansii** March. Fung. coproph. p. 39. — Sparsus 2215 vel gregarius, basi filamentis albis, subtilissimis, evanescentibus, adfixus, primo globoso-piriformis, demum planiusculus, sessilis, glaber, purpurascens, 2,5-4 mm.¹ diam., margine lato denticulato-fimbriato tandem eximie reflexo; disco plano, dein leniter concavo, subundulato valde crystallino-papillato; paraphysibus ascos subæquantibus gracilibus, septatis, hyalinis, simplicibus vel ramosis, apice paullo incrassatis; ascis cylindraceis, rectis, superne vix attenuatis, breviter stipitatis, 180-215 = 17-21, jodo non cœrulescentibus, 8-sporis; sporidiis monostichis, 26-28 = 14-16, ellipsoideis, incoloribus, levibus, plerumque nucleo conspicuo.

Hab. supra fimum Leporum et Cuniculorum, socio *Ascophano sexdecimsporo*, Aywaille Belgii. — Affinis *A. difformi*, sed ascis cylindraceis jodo non tinctis, sporidiis majoribus mox distinctus.

II. *Polygenæ*, in tela, lana, corio, rarius in fimo et humo nascentes.

28. **Ascophanus carneus** (Pers.) Boud. Ascob. pag. 60, t. XII, f. 38, 2216 *Ascobolus* Pers. Syn. p. 666, nec Myc. Eur., Fries S. M. II, p. 165. — Sparsus vel rarius subaggregatus, minutus, sessilis, 1-1,5 mm. lat., roseus vel carneus, glaber, initio globosus dein applanatus, disco convexo, immarginato, applanato; ascis late clavatis deorsum attenuatis, saepius rosellis, octosporis, 100-120 = 20-24; sporidiis ellipsoideis, levibus vel minutissime granulosis, hyalinis, initio muco obvolutis, 16-18 = 9-10; paraphysibus obtusis, clavatis, septatis.

Hab. in fimo vaccino, in tela et lana putrescentibus, in Suecia, Germania, Britannia, Gallia, Hungaria, Belgio, Italia, Fennia, Amer. bor. — Var. **Cuniculi** Boud. Asc. loc. cit. fig. 39, Pat. Tab. An. fig. 380: cupulis aurantiis vel aurantio-carneis, 1 mm. lat.; paraphysibus hyalinis. In stercore Cuniculorum in Britannia et Gallia.

29. **Ascophanus saccharinus** (Berk. et Curt.) Boud. Ascob. p. 61, t. 12, 2217 f. 40. — Sparsus vel consertus, disco hemisphærico, dein expanso

et subplano saccharino-nitido, rubro-lateritus v. salmonicolor, siccus margine pallidior; ascis amplis, octosporis; sporidiis ellipsoideis, hyalinis, 17-20 μ . long.

Hab. ad fimum, corium vetustum et telam putrem in Gallia, Britannia. — Ascoma basi hyphis albidis affixum. Potius var. *A. carnei*. Ceterum *A. carneus*, *saccharinus*, *diformis*, *testaceus* nimis inter se affines.

30. **Ascophanus diformis** (Nyl.) Boud. *Ascob.* pag. 62, Karst. Rev. Monogr. p. 121, *Pez. diformis* Karst. Fenn. I, p. 61, *Ascobolus diformis* Nyl. Obs. p. 85. — Gregarius vel confertus, saepe confluens, sessilis, primum convexus, demum planiusculus, carneus vel dilute subaurantiacus, basi filamentis intricatis, albis adfixus 0,5-0,6 mm. lat.; ascis oblongato-clavatis, 20 μ . cr., jodo cœrulecentibus, 8-sporis; sporidiis distichis, ellipsoideis, eguttulatis, 18-22 = 10-14; paraphysibus simplicibus, mollibus, articulatis, sursum vix vel levissime incrassatis, 4-6 μ . cr.

Hab. in corio et panno lineo in Fennia, in fibris ligneis putridis Pircuniæ in Argentina. — (Asci 150-160 = 25-35; sporidia 20 = 10). Teste Phill., non differt ab *Ascophano testaceo*. Sec. Rhem est var. *A. carnei*.

31. **Ascophanus testaceus** (Moug.) Phill. Disc. p. 310, t. IX, f. 58, 2219 *Ascob. testaceus* Moug. Wallr. Fl. crypt. IV, pag. 513, B. et Br. Ann. N. H. n. 1082, t. 14, f. 5, Fuck. Symb. App. II, p. 58, f. 23, *Peziza test.* Moug. in Fr. Elench II, p. 12, *Helotium* Berk. Outl. p. 372. — Cupulis testaceo-rufescensibus, carnosis, gregariis sparsisve, sessilibus, primo perfecte globosis, clausis, demum ore lacerato magis magisque dilatatis, postremo hemisphæricis, tympaniformibus, 2-3 mm. latis, orbicularibus seu irregularibus, extus glabris seu junioribus paucitrachis, margine distincto, acuto, erecto, subtiliter lacerato, vix pallidiore, disco plano concolore, subtiliter punctulato; ascis oblongo-cylindraceis, breviter stipitatis, 8-sporis, 170 = 18; sporidiis oblique monostichis, oblongo-ovatis, uniguttulatis, continuis, rectis, hyalinis, 24 = 12 (Fuck.), 18-20 = 9-11 (Phill.); paraphysibus simplicibus, anguste clavatis, clavulis fuscis.

Hab. in tela putrida et in stercore murino et Cuniculi in Rhenogovia, Gallia, Britannia. — *Patellaria testacea* Fr., ex ipso auct. (Summa p. 366) est *Lachnea scutellata*.

32. **Ascophanus miniatus** (Preuss) Sacc. *Ascob. miniatus* Preuss Hoyerw. n. 155. — Sessilis, glaber, carnosus, subconvexus, immar-

- 536 Discomyceteæ, Ascoboleæ, Hyalosporæ, Lasiobolus.
 ginatus, dilute miniatus; ascis clavatis; sporidiis ovatis, albis; paraphysibus simplicibus, supra capitato-clavatis, miniatis.
Hab. in cortice Juglandis in hortis Hoyerswerda.
33. **Ascophanus consociatus** (B. et Br.) Phill. Disc. p. 312, *Ascobolus* 2221
 B. et Br. Ann. Nat. Hist. n. 1491, t. II, f. 7. — Gregarius
 subhemisphæricus, extus rugoso-granulatus, pallide flavus vel albidus; ascis late clavatis, brevibus, octosporis; sporidiis distichis,
 late fusoideis, intus granulosis, 10 = 6; paraphysibus filiformibus.
Hab. ad Chætosphaeriam cupuliferam vetustam Land-
 ridge Brit. — Ascoma 100-150 μ. lat.; sporidia luteola.
34. **Ascophanus pallens** Boud. in Bull. Soc. B. et M. tr. 1888, pag. 2222
 XLVIII, t. II, f. 2. — Minutus, 1-3 mm. latus, albidus aut albi-
 do-cinereus, hemisphæricus aut lenticularis, submarginatus, extus
 levus, disco plano aut convexo, ascis prominentibus longe papilla-
 to; paraphysibus hyalinis, saepius continuis ad apicem incrassatis,
 ad basim divisis et septatis, apice 6-8 μ. cr.; ascis maximis, am-
 plis, clavatis, octosporis 290-350 = 35-45, operculo lato leniter um-
 bonato; sporidiis maximis, oblongo-fusiformibus, hyalinis, intus
 nullo modo granulosis, 40-45 = 15-20.
Hab. ad terram argillosam cum Confervis in umbrosis silva-
 rum Ecouen Galliæ.
35. **Ascophanus Guernisacii** Cr. Finist. p. 56, tab. ultima, fig. 2. — 2223
 Brunneus, basi albicans, hemisphæricus, levus, 2-3 mm. lat., mar-
 gine membranaceo, lobato; ascis subcylindraceis, amplis, octospo-
 ris; sporidiis cymbiformibus, hyalinis, inæqualiter granulosis; pa-
 paraphysibus simplicibus, filiformibus contextu crasso, reticulato,
 colorato.
Hab. ad terram schistosam et in muris antiquis, Finistère
 Galliæ.
- LASIOBOLUS** Sacc. Cons. gen. Disc. p. 8 (1884), Karst. Rev. p. 122,
 Boud. Class. Disc. p. 20 (Etym. *lasios* pilosus et *bolus* pro Asco-
 bolo), *Ascophani* spec. Auct. — Ascomata carnosula, sessilia, pla-
 na vel convexa, distincte setosa vel pilosa. Asci oblongi vel clava-
 ti, 4-8-spori, paraphysati e disco erumpentes. Sporidia ellipsoidea
 vel oblonga, continua, hyalina, typice levia.
1. **Lasiobolus equinus** (Müll.) Karst. Rev. pag. 122, *Peziza equina* 2224
 Müll. in Flor. dan. t. 779, f. 3, *Pez. papillata* Pers. Syn. Fung.
 p. 650, Nees Syst. f. 276, *P. diversicolor* Fr. Syst. myc. II, p. 88,
P. cervina Pers. Myc. cur. I, p. 254, *Ascobolus papillatus* Wallr.

Flor. Crypt. Germ. II, pag. 513, *Lasiob. papillatus* Sacc. Consp. Gen. Disc. p. 8, *Ascophan.* Boud. Ascob. p. 62, t. XII, f. 41. — Cupulis gregariis, obconice prominulis, ab initio pallido-luteis, dein plus minusve aurantiacis v. fulvescentibus, pilis erectis, albido-incoloribus strigosis, 0,3-0,5, mm. lat.; ascis oblongato-clavatis, 22-33 mm. crass., jodo non tinctis; sporidiis octonis, ellipsoideis, 18-24 \approx 11-14; paraphysibus 2-3,5 μ . crass.

Hab. in fimo equino, vaccino, suino, Cuniculorum, murino in Suecia, Fennia, Lapponia, Belgio, Germania, Gallia, Amer. bor. — Pili cupulæ subulati, inarticulati, 220-260 \approx (basi) 14-22. — ***L. ciliatus** (Berk.) Sacc. *Ascobolus* Berk. Engl. Fl. V, p. 374, Ann. N. H. n. 1083, t. 14, f. 7, an Fr., *Ascophanus* Boud. Ascob. p. 63, Phill. Disc. p. 312: sessilis, subhemisphæricus, levis, aurantiacus ad basim setulis continuis, albis cinctus; disco plano, margine tumido, albo, glabro; ascis late clavatis; sporidiis ellipsoideis, levibus, hyalinis, 20-22 \approx 10; paraphysibus filiformibus, aurantio-granulosis. In fimo vaccino et equino in Britannia, Germania, Amer. bor. et Africa australi. Quid spectet *A. ciliatus* Fr. S. M. II, pag. 164 (margine albo ciliato), ignotum. — ***L. pilosus** (Fr.) Sacc. Consp. gen. Disc. p. 8, *Ascobolus* Fr. S. M. II, p. 164, Pat. Tab. an. f. 487, *Ascophanus* Boud. Ascob. pag. 64, t. XII, f. 42-44: gregarius, sessilis, obconice prominulus, granuliformis, inferne pilis albidis, sparsis strigosus, primum vulgo pallidus, dein fulvo-testaceus vel plus minusve aurantiacus, minutissimus; ascis oblongatis, basi stipitato-constrictis, 100-120 \approx 22-34, jodo non tinctis, 8-sporis; sporidiis ellipsoideis, 19-23 \approx 10-14; paraphysibus gracilescensibus. In fimo caprino, capreolino, ovino, vaccino, equino, canino, in Suecia, Germania, Britannia, Gallia, Hungaria, Italia, Amer. bor. et merid. Quoad formas varias cfr. Boud. loc. cit.

2. **Lasiobolus raripilus** (Phill.) Sacc. *Ascobolus raripilus* Phill. in Grevillea VII, p. 23. — Gregarius vel confertus, minutus, sessilis, glaber, primo globosus, dein hemisphæricus, disco convexo, pallido, vitellino-flavo, extus pilis paucis pallidis, septatis, rectis parce obsito; ascis late clavatis, octosporis; sporidiis ovatis, 25 \approx 14, levibus, hyalinis; paraphysibus linearibus, simplicibus, apice clavatis.

Hab. in fimo vaccino in California (HARKNESS).

3. **Lasiobolus ruber** (Quél.) Sacc. *Ascophanus ruber* Quél. 12 Suppl. 2226 p. 13, t. VII, f. 13 et in Grevillea 1879, p. p. — Ascomate globuloso, dein hemisphærico, 1 mm. diam., rufulo, setulis subtilibus,

538 Discomyceteæ, Ascoboleæ, Hyalosporæ, Lasiobolus.

albis nitentibus hirto, hymenio plano, papillato, rufo-aurantiaco; sporidiis ellipsoideis, 20 μ . longis, hyalinis.

Hab. in fimo vaccino in pascuis alpinis Juræ. — *L. ciliato* similis sed margine integro et sporidiis hyalinis differt.

4. **Lasiobolus diversisporus** (Fuck.) Sacc. *Ascob. diversisporus* Fuck. 2227

S. M. p. 289, *Humaria* Speg. in Mich. I, p. 236. — Cupulis gregariis, usque ad 1 mm. latis, $\frac{1}{2}$ mm. altis, primo globosis, demum vertice applanatis, submarginatis, diaphanis, flavo-rubris, extus sparse pilosis, pilis rectis, hyalinis, continuis, cupulæ diametrum subæquantibus; ascis stipitatis, oblongis, 4-8-sporis, longitudine valde varia, plerumque 212 \approx 21; sporidiis inordinatis, ovatis vel ovato-oblongis, magnitudine in eodem asco item valde varia, 20-24 \approx 12-16 hyalinis; paraphysibus filiformibus, ramosis (?).

Hab. in fimo vaccino in Rhenogovia et Ital. bor. — In forma italica (ex Conegliano) sunt ascomata viridulo-testacea, campanuliformia, sessilia, undique setis hyalinis ornata; setulis unicellularibus, e basi bulbillosa acutissimis, crasse tunicatis; ascis saccato-cylindracei deorsum in stipitem filiformem brevem abrupte attenuati, apice obtuse rotundati, 150-180 \approx 30-38, a paraphysati (?); sporidia inordinate monosticha, oblongo-ellipsoidea, utrinque obtusa, 22-26 \approx 12-15, eguttulata, hyalina.

5. **Lasiobolus albicans** (Fuck.) Sacc. *Ascobolus albicans* Fuck. in 2228

Hedwigia 1866, p. 3, f. 2, a, b, c. — Ascomatibus carnosis, rubescensibus, late obconicis, usque ad 4 mm. lat., 2 mm. alt., planis concavisve, disco argute marginato paullo pallidiore, siccitate albo-pruinoso, extus margineque pilis brevissimis obsitis; ascis longissime stipitatis, oblongis, octosporis; sporidiis ovatis, hyalinis; paraphysibus apice bi-tripartitis hyalinis.

Hab. ad terram inter muscos, in fimi equini vicinia, in pinetis Budenheim Rhenogoviæ.

6. **Lasiobolus avicularius** Karst. Rev. p. 122, *Peziza* Karst. M. F. I, 2229

pag. 74. — Gregarius, obconicus, fere immarginatus, pilis erectis hyalino-albidis strigosus, hyalino-albidus, punctiformis, disco planiusculo; sporidiis fusideo-oblongatis, eguttulatis, 10-11 \approx 4.

Hab. supra fimum ovium e. g. *Tetraonum*, prope oppid. Abo Fenniæ.

7. **Lasiobolus hirtellus** Karst. Rev. p. 122, *Peziza* Karst. M. F. I, 2230

p. 74. — Gregarius sessilis, planiusculus, pallidus, pilis erectis, albido-incoloribus strigosus, 0,5 mm. lat.; ascis 50-100 \approx 12, jodo

non tinctis; sporidiis ellipsoideis, 10-12 = 6-7; paraphysibus gracilescensibus, 2-3 μ . cr.

Hab. in fimo Tetraonum prope Mustiala Fenniæ.

RYPAROBUS Boud. Ascob. p. 47 (Etym. *ryparos* sordidus et *bios* vita) *Thecotheus* Boud. Ascob. p. 45, *Ascozonus* Renny. — Ascomata carnosula, sessilia, scutellata v. disciformia, minuta v. minutissima glabra v. ciliatula. Asci crassiusculi sed breves, rarius amplissimi, 16-myriospori, operculo apice dehiscentes, e disco exsilentes, varie paraphysati. Sporidia ellipsoidea, oblonga v. subfusoidea, continua, hyalina, quandoque in glomum fere congregata.

I. *Eu-Ryparobius*. Asci minuti v. mediocres.

1. **Ryparobius crustaceus** (Fuck.) Rehm Ascom. p. 52, *Ascobolus* 2231 *crustaceus* Fuck. in Hedw. 1866, p. 4, *Asc. Cookei* Crouan Flor. Finist. p. 56, suppl. f. 3, (1867) *Asc. polysporus* Auersw. in Hedw. 1868, p. 51, *Pezizula crustacea* Karst. M. F. I, p. 81, *Ryparobius Cookei* Boud. Ascob. p. 48, t. IX, fig. 24, Phill. Disc. p. 299, t. IX, f. 57. — Aggregatus, sæpissime confluens, sessilis, depresso-sphæroideus, glaber, fuscus, disco pallidiore, 0,1 mm. lat.; ascis paucis, obovoideo-oblongatis, basi paullo curvatis, 60-100 = 20-28, jodo non tinctis; sporidiis 48-64 incoloribus, congregatis, ovoideis aut oblongato-ovoideis, 7-10 = 4,5-5,5; paraphysibus parcis, gracilescensibus, apice subclavato.

Hab. in fimo canino, suino, equino, vaccino putrescente in Germania, Fennia, Gallia, Italia, Belgio, Britannia, Fuegia. — ***myriadeus** Karst. M. F. I, p. 81, ?*Ascob. caninus* Auersw. in Hedw. l. c.: typo simillimus sed ascis 24-34-sporis, clavatis, 44 = 11 recedens. Cum typo in Fennia et Germania.

2. **Ryparobius caninus** (Auersw.) Sacc. *Ascobolus* Auersw. Hedw. 2232 1868, p. 51. — Microscopicus, depresso-globosus, fuscus v. brunneus, glaber, acute marginatus, disco pallidiore; ascis clavatis, polysporis 44 = 11; sporidiis ovalibus, hyalinis, monoblastis, levibus 7-8 = 4, 24-32 in quovis asco.

Hab. in stercore canino, Rosenthal Cassel. — Cfr. præcedentem.

3. **Ryparobius polysporus** (Karst.) Sacc. Mich. II, p. 571, *Pezizula* 2233 Karst. M. F. I, p. 82. — Sparsus v. gregarius, sessilis, glaber, fuscidulus, 0,1 mm. lat.; ascis paucis ellipsoideis 118-130 = 50, jodo

non tinctis; sporidiis ellipsoideis, conglobatis, 150-200, hyalinis, 6-3; paraphysibus gracilibus.

Hab. in fimo equino pr. Abo et Mustiala Fenniæ, in fimo vaccino Newfield N. J. Amer. bor. (forma ascis 70-80 ± 18-20; sporidiis 5-3,5-4) et in Argentina Amer. merid.

4. **Ryparobius brunneus** Boud. Ascob. p. 47, t. IX, fig. 23. — Confer. 223 fertissimus, minutissimus, perexiguus, glaber, fulvo-brunneus, disco concolore, subconvexo; ascis oblongis, latis, basi acutatis; sporidiis 32, oblongo-ovoideis, minutis, hyalinis, utrinque subacutis; paraphysibus parcis, septatis, apice vix incrassatis.

Hab. in stercore humano, vaccino et asinino, Montmorency Galliæ.

5. **Ryparobius felinus** Boud. Asc. p. 49, tab. X, fig. 45. — Confer. 223^a tus aut aggregatus, minutissimus, fulvo-cinnamomeus, disco convexo, saturatiore, margine vix conspicuo; ascis amplis, oblongis, basi attenuatis; sporidiis 64, oblongo-elongatis, hyalinis; paraphysibus parcis, septatis, hyalinis, apice incrassatis.

Hab. in stercore felino circa Montmorency Galliæ — Affinis *Ryp. brunneo*.

6. **Ryparobius myriosporus** (Cr.) Boud. Ascob. p. 50, tab. X, fig. 27, 223^b *Nectria myriospora* Crouan Fin. p. 37, tab. ult., fig. 15, *Chilonectria* Sacc. Syll. II, p. 454. — Minutissimus, congestus, rarius sparsus, $\frac{1}{10}$ - $\frac{1}{5}$ mm. lat. rufescens, rotundatus, glaber versus marginem ruber, disco plano, aereis exsertis papillato; ascis amplissimis, ovoideis, basi vix attenuatis, operculo amplissimo dehiscentibus; sporidiis numerosissimis, 200-250, ovoideis, minutis, hyalinis, apice subacutis, initio ascum replentibus, demum in asci apice conglobatis.

Hab. in stercore canino, Montmorency, in stercore equino Finistère Galliæ.

7. **Ryparobius saccoboloides** Speg. Mich. I, p. 474. — Dense gregarius, minutissimus, 80-100 μ . diam., difficilime nudo oculo perspiciens, sordide roseo-fuscens, disco obscuriore virescente, primo globosus, hemisphæricus, subsuperficialis; contextu excipuli minutissime ac densissime parenchymatico, dilute roseo-fuscidulo; ascis ovoideis v. elliptico-clavatis, superne obtusis, inferne in pedicellum crassum ac nodulosum desinentibus, 40-50 ± 20-25, 60-120-sporis, a paraphysatis (?); sporidiis in glomerulo 35-40 ± 15-20 elliptico, tenaciter muco connexis, etiam extra ascos, ellipticis v. ovoi-

deis, utrinque obtusiusculis, continuis, hyalinis, luce refracta dilute chlorinis, crassiuscule tunicatis, $5-6 \approx 2\frac{1}{2}-3$.

Hab. in fimo canino vetusto putrescente, Conegliano Ital. bor. — Iodi ope asci et sporidia rubro-fuliginea evadunt, stipites autem et bases ascorum ita colorantur ut opaci fiant.

3. **Ryparobius pachyascus** Rehm Asc. fasc. XIX, pag. 5, n. 914. — 2238
— Gregarius, raro sparsus e globoso depresso, 60-100 μ . diam., subimmersus, tenuis, diaphanus, flavidus, rarius brunneo-flavus; ascis ovoideis, basi acutatis, 4-8, rarius 1-16 in quoque ascomate, polyporis, 77-90 \approx 38-40; paraphysibus parcis ramosis, curvulis cito evanidis; sporidiis ovoideis, 6 \approx 3, hyalinis.

Hab. in fimo equino et Cuniculorum in Austria (ZUKAL).

— Apex ascorum senio operculatim secedens; sporidia initio strato mucoso sphærico simul obvoluta.

9. **Ryparobius albidus** Boud. Bull. Soc. Mycol. IV, pag. XLIX, t. II, 2239
f. 3. — Minutissimus sed pro genere major, albidus, sparsus aut gregarius, 0,2-0,4 mm. diam., primo hemisphæricus, dein lenticularis, immarginatus extus et supra pallidus vix cinerescens, disco placo dein convexo, ascis papillato; paraphysibus cylindricis, subspisis, bene septatis, ad apicem vix 4 μ . cr.; ascis late clavatis, hyalinis, 32-40-50 \approx 5-6; sporidiis maturis ad apicem ascorum in massam ovatam conglobatis, hyalinis, levibus, oblongo-fusiformibus, intus non granulosis, 10-15 \approx 5-6.

Hab. ad sterlus vaccinum in pratis, loco dicto «Foret de Carnelle» pr. Montmorency Galliæ.

10. **Ryparobius dubius** Boud. Ascob. tab. X, fig. 26. — Sparsus, per- 2240
exiguus, sæpe immersus, globulosus, pallide griseus vel fulvus, disco convexo, pellucido, immarginato; ascis oblongis vel oblongo-ovatis, hyalinis, basi obtusis vel subattenuatis; sporidiis 100-128, ellipsoideis, hyalinis, subacutis; paraphysibus nullis vel obsoletis.

Hab. in fimo equino, ovino et Cuniculorum in Gallia, Britannia et Belgio. — Sec. Rehm ascom. n. 105, asci 40-45 \approx 24, circiter in quoque ascomate seni; sporidia ellipsoidea, 6 \approx 4. An idem ac *Rypar. myriosporus* Boud.?

11. **Ryparobius sexdecimporus** (Cr.) Sacc. *Ascobolus* Cr. Ann. Sc. 2241
Nat. 1858, p. 195, tab. XIII, fig. 21-26, *Ascoph.* Boud. Ascobol. p. 57, tab. XI, fig. 35. — Sparsus, minutus, sessilis hemisphæricus, convexus vel planus, glaber, albus dein griseus, demum flavidus-albus; ascis oblongis, latis, basi acutatis; sporidiis 16, ellip-

psoides, hyalinis, levibus, 11-12 = 6-7; paraphysibus simplicibus v. furcatis; apice vix incrassato, recto vel curvulo.

Hab. in fimo equino, vaccino in Gallia, Britannia, Belgio, Germania. — Ascoma 1-2 mm. diam.

12. **Ryparobius hyalinellus** (Karst.) Sacc. *Pezizula hyalinella* Karst. 2242

M. F. I, pag. 83. — Gregarius, valde cenvexus, fere sphæroideus, hyalinus, immarginatus, glaber, latit. vix 0,1 mm.; ascis oblongato-clavatis, 55-100 = 18-28; sporidiis 72, sphæroideo-ellipsoideis, hyalinis, 5 = 3; paraphysibus filiformibus, apice ovoideis, 6 μ . cr.

Hab. supra fimum Tetraonum in silva Haarankorpi prope Mustiala Fenniae. — ***promiscuus** Karst. M. F. I, p. 83: *hyalinello* similis, ascis vero 50-62 = 16-18 sporidiisque 32, chlorino-hyalinis, 6 = 4 diversa. Cum specie.

II. *Thecotheus* Boud. Asci magni, valde exsilientes.

13. **Ryparobius Pelletieri** (Cr.) Sacc. Mich. I, p. 605, *Ascobolus Pel-* 2243

letieri Cr. Finist. p. 56, Ann. Sc. Nat. 1857, p. 173, tab. IV A, fig. 1-4, Pat. Tab. An. fig. 172, *Thecotheus* Boud. Ascob. p. 46, t. IX, fig. 22, A. *Solms-Laubachii* Rabh. Bot. Zeit. 1862, p. 198, *Pezizula Pelletieri* Speg. Mich. I, 238. — Ascomatibus primo sub-sparsis globosis, dein patellaribus, candidissimis vel dilute cinereo-roseis, $\frac{3}{4}$ - $1\frac{1}{4}$ mm. diam., extus pruinosis; ascis cylindraceo-clavatis, deorsum attenuatis, breve stipitatis, apice obtuse rotundatis, 200-300 = 50-55, 30-32-sporis; sporidiis tetrahexastichis elliptico-fusiformibus 27-30-35 = 14-18, utrinque acutiuscule rotundatis, crassiuscule tunicatis, hyalinis.

Hab. in fimo equino in umbrosis in Gallia, Italia, Germania, Britannia et Belgio.

14. **Ryparobius Winteri** March. Fung. coproph. p. 39, t. IV, f. 1-7. — 2244

Sparsus, rarius subgregarius, globoso-cylindricus, albus v. albocyanellus, glaber, 400-500 = 200-300, margine obsolete dentato, tandem infexo; disco concolore plano, ascis parce papillato; paraphysibus numerosis, hyalinis, linearibus, 1-2 μ . crass., septatis, ramosis, inferne non raro crassioribus, articulatis, ad apicem curvatis, ascos superantibus; ascis oblongato-cylindraceis, subsessilibus, rectis vel deorsum curvatis, operculo convexo dehiscentibus, sporidia circ. 256 (?) includentibus, jodo haud cœrulescentibus, 160-210 = 45-55; sporidiis ellipsoideis, utrinque rotundatis, 10-11 = 5-6 hyalinis, nucleo magno et conspicuo.

Hab. in fimo damarum, Tervueren Belgii. — Cupulis cylin-

draceis albis, sporidiis ratione majusculis, grosse nucleatis mox dignoscitur.

III. *Ascozonus* Renny. Ascii (saltem in plerisque) operculo magno conico, subinde et longitr. fisso, transverse dehiscentes. Ascomata ciliata v. pilosula.

5. **Ryparobius subhirtus** (Renny) Phillips. Disc. p. 302, *Ascozonus* 2245
Renny Journ. Bot. 1874, p. 357, tab. CLV, fig. 4-7. — Minutus
sessilis, hemisphæricus, albus, subdiaphanus pilis brevibus inæqua-
libus sæpe fasciculatis, conspersus, margine ciliato, disco plano,
papillato; ascis latissimis, oblongis basi acutatis; sporidiis circ.
128, ellipsoideis, hyalinis, levibus, initio in massam ovoideam col-
lectis; paraphysibus non visis.

Hab. in stercore Cuniculorum, Hereford Britanniæ (RENNY).

6. **Ryparobius Crouani** (Renny) Phill. Disc. pag. 300, *Ascozonus* 2246
Crouani Renny in Journ. Bot. 1874, p. 356, tab. 154, fig. 6-10
nec *Ascob. Crouani* Cooke = *Humaria*. — Perexiguus, initio al-
bus, dein albidus, fragilis, sessilis, hemisphæricus, glaber, excipulo
strato unico cellularum formato, margine setulis brevibus subacu-
minatis, rugulosis uniseriatis ciliato; ascis late clavatis; sporidiis
32, oblongo-fusoideis, utrinque obtusis, hyalinis, levibus; para-
physibus non visis.

Hab. in stercore Cuniculorum, Hereford Britanniæ (RENNY).

— Ascoma 200 μ . diam.

7. **Ryparobius Woolhopensis** (Renny) B. et Br. Ann. Nat. Hist. 2247
n. 1395, tab. IX, fig. 12, *Ascozonus* Renny Journ. Bot. 1874,
p. 356, tab. CLIII. — Minutus, sparsus, initio albus dein sordidus,
crassiuscule stipitatus, tuberculatus, margine setuloso-fimbriatus
dein nudus, excipulo basi vesiculoso; ascis late clavatis; sporidiis
64, late fusoideis, hyalinis, levibus, 17-18 \times 5; paraphysibus fili-
formibus obtusis, simplicibus.

Hab. in stercore avium hyphis Mucoris commixtus, Here-
ford Britanniæ (RENNY). — Ascoma 400 μ . lat.

8. **Ryparobius argenteus** B. et Br. Ann. Nat. Hist. n. 1394, t. IX, 2248
f. 11, Grev. t. 22, f. 4, *Ascozonus cunicularius* Renny Journ. Bot.
1874, p. 355, tab. CLV, fig. 1-3. — Perexiguus, initio globosus,
dein applanatus, argenteo-albus, pilis teneris ciliatus; ascis latis,
brevibus 100 μ . longis operculatim dehiscentibus; sporidiis 64, fu-
soideis hyalinis, levibus, 17 \times 7,6; paraphysibus filiformibus, fur-
catis, apice leniter incrassatis.

Hab. in fimo Cuniculorum, hyphis Mucoris plerumque af-

fixus, Hereford Britanniæ (RENNY), in Belgio (MARCHAL). — Ascoma 100 μ . lat. — An hoc pertineat *Comesia cunicularia* (Boud.) valde verisimile.

19. Ryparobius Leveilleanus (Renny) Phill. Disc. p. 301, *Ascozonus* 2249

Renny Journ. Bot. 1874, p. 356, tab. CLIV, fig. 1-5. — Minutissimus, breve stipitatus, argenteo-albus, stipite cellulis prominulis, rotundis formato, margine setulis brevibus inæqualibus, uniseriatis ciliato, disco papillato; ascis latissimis, basi acutatis, operculo umbonato; sporidiis 64-96, oblongo-fusoideis, utrinque obtusis, hyalinis, levibus, paraphysibus non visis.

Hab. in fimo Cuniculorum, Hereford Britanniæ (RENNY). — Ascomata 200-300 μ . lat. Excipulum cellulis minutis, applanatis formatum.

20. Ryparobius parvisporus (Renny) Phill. Disc. pag. 303, *Ascozonus* 2250

Renny Journ. Bot. 1874, pag. 356, tab. CLVI, fig. 1-5. — Peregrinus, fragilis, carnosulus, albus, nitidulus, dein sordidus, subvinosus, margine setulis subulatis, asperulis, 1-2-seriatis, inæqualiter fimbriato; disco plano, dein ascis prominentibus punctato; ascis late clavatis; sporidiis 16-24, ellipsoideis, hyalinis, levibus; paraphysibus filiformibus, obtusis, septatis.

Hab. in stercore Cuniculorum, Hereford Britanniæ.

21. Ryparobius niveus (Fuck.) Sacc. *Ascobolus niveus* Fuck. in Hedw. 2251

1866, 1, 4, f. 3 a, b, c et Symb. myc. pag. 289. — Ascomatibus sparsis, punctiformibus, usque ad 1 mm. lat., planis vix excavatis, hyalino-diaphanis, niveis, extus margineque pilis concoloribus subtilissime puberulis; ascis stipitatis, oblongo-ovatis, curvatis, 64-sporis, demum operculatim dehiscentibus; sporidiis farctis, ellipticis, hyalinis, continuis; paraphysibus nullis.

Hab. ad fimum caninum putridum in M. Rabenkopf Rheno-goviæ.

22. Ryparobius Asteriscus Karst. Discom. nov. in Notis. ur Sällsk. 2252

pro Faun. et Flor. Fenn. Förh. XIII, p. 446, Rev. monogr. atque Synops. Ascomycet. fenn. pag. 122. — Subgregarius, sessilis, subsphæroideus, disco planiusculo, vitreo-pellucido, margine pilosellus, minutissimus 0,1-0,2; ascis subsessilibus, obovoideo-oblongatis 94-96 = 24-33; sporidiis 48-60, conglobatis, utrinque attenuatis seu acutatis, simplicibus, eguttulatis, hyalinis, 10-16 = 4; paraphysibus parcis, gracilibus, cito pereuntibus, apice incrassatis.

Hab. in stercore vulpino vetusto prope Mustiala Fenniæ.

IV. *Moutonia* Sacc. Ascomata monoasca.

23. **Ryparobius monoascus** Mout. Ascom. Liège p. 141. — Sparsus, 2253 plerumque immersus, hemisphæricus, 150-200 μ . diam. contextu membranaceo, polygonio, subhyalino, asco in quoque ascomate unico, ovoideo, 200 \approx 150-170, sessili, e disco emergente, multisporo; sporidiis (ultra 500) ovatis, hyalinis, 5-6 $\frac{1}{2}$ \approx 4-4 $\frac{1}{2}$.

Hab. in fimo Capreolorum, Angleur, Beaufays Belgii ¹⁾.

GYMNODISCUS Zuk. Ein. Ascom. pag. 8, tab. I, f. 1 (1887) [Etym. *gymnos* nudus et *discus*]. — Ascomata carnosо-ceracea, initio concava dein convexula, margine et excipulo nullo, hypothecio globoso-discoideo parenchymatico. Asci clavati, leniter exslientes, polyspori. Sporidia elliptico-oblonga, continua, hyalina, initio muco hyalino obvoluta. Paraphyses bacillares, subinde furcatæ. — Est *Ryparobius* excipulo carens, ad *Gymnoascaceas* nutans.

1. **Gymnodiscus neglectus** Zuk. Ein. Ascom. p. 8, t. I, f. I a, c. — 2254 Solitarius, albidus, 200-250 μ . lat. et alt., glaber; ascis clavatis, breve stipitatis, apice rotundatis, 86 \approx 21, operculo secesso, truncatis, polysporis; sporidiis ex ellipsoideo breve fusoideis, polystichis, 10-11 \approx 3, initio strato mucoso hyalino subgloboso obvolutis; paraphysibus interdum furcatis.

Hab. in fimo, Wien Austriæ (ZUKAL).

Fam. 5. **Dermateæ** Fr. Summa p. 345.

Ascomata urceolata, cupulata vel demum applanata, sessilia, rarius stipitata, suberosa, coriacea vel cornea, extus saepius furfuracea, typice brunnea vel atra, saepissime cæspitoso-erumpentia. Asci 4-8-spori vel myriospori, typice paraphysati. Sporidia continua vel septata, hyalina, rarius colorata. — Familia ægre limitanda, in proximas *Pezizeas*, *Patellarieas* etc. sensim abiens: genera quoque difficilia ex eo quod plures species habitu simillimæ, fructificatione generice diversa (nempe octospora vel myriospora, hyalospora vel didymospora) utuntur; dubium ergo an de polymorphismo agatur. Dolet eximium opus Rehmianum de Discomyc. germanicis serius pervenisse, ut eo ubique uti potuissem; species vero ejusdem omnes ad diem editas, quantum potui, recepi.

1) Teste cl. Heimerl (Niederöst Ascob. p. 30, opella egregia sub prelo accepta), *R. monoascus* Mout. ad gen. *Thelebolum* est referendus et totum hoc genus *Ascoboleis* adscribendum, præeuntibus quoque observationibus cl. H. Zukalii (Denkschr. K. Akad. Wien 1885). Cfr. Heim. l. c. ubi *Theleb. stercoreus* Tode, *Th. nanus* Heim. et *Th. Zukalii* Heim. describuntur una cum plur. aliis novis *Ascoboleis*.

Clavis analytica generum.

Sect. 1. Hyalosporæ. Sporidia ovoidea, oblonga v. fusoidea, hyalina.

* Stipitatæ, majores.

Midotis. Ascomata auriformia, subverticalia, coriacea.

Urnula. Ascomata urceolato-turbinata, coriacea.

** Subsessiles, minores.

Dermatea. Ascomata subsuberosa, mox patellari-aperta. Asci octospori.

Cenangium. Ascomata coriaceo-subcornea, urceolato-scutellata. Asci octospori.

Tympانis. Ascomata *Cenangii*. Asci myriospori vel myriospori simul et octospori.

Ameghinella. Ascomata membranaceo-coriacea, stromate tuberculato, initio hypodermico inserta. Asci 8-spori.

Ephelina. Ascomata ceraceo-subcoriacea, stromate crustaceo effuso inserta. Asci 8-spori.

Laquearia. Ascomata coriaceo-subcornea, dimidiata, infra deficiens. Asci 8-spori.

Sect. 2. Phæosporæ. Sporidia ovoidea vel oblonga, fusca.

Hymenobolus. Ascomata coriacea initio tecta et clausa, dein erumpentia et ore lacero cupulari-aperta. (Ad *Phacidieas* prope accedit).

[*Phæangium.* Sporidia fusca, cetera *Cenangii*. Vide Sect. III. *Cenangii*].

Sect. 3. Didymosporæ. Sporidia ovoidea, oblonga v. fusoidea, 1-septata, hyalina v. fusca.

Cenangella. Ascomata coriacea vel subcornea ex urceolato patellaria. Sporidia hyalina.

[*Phæangella.* Sporidia fusca, cetera *Cenangellæ*. Vide Sect. II. *Cenangellæ*].

[*Tryblidium.* Ascomata hysterioidea. Sporidia fusca. Cfr. Syll. II, p. 740].

Sect. 4. Phragmosporæ. Sporidia oblonga v. fusoidea, 2-pluriseptata, hyalina v. colorata.

b.594 *Scleroderris.* Ascomata coriacea vel subcornea, glabra, ex urceolato patellaria. Sporidia hyalina.

[*Phæoderris.* Sporidia fusca, cetera *Scleroderridis*. Vide Sect. II. *Scleroderridis*].

Crumenula. Ascomata ceraceo-membranacea, pilosula, urceolata. Sporidia hyalina.

[*Tryblidiella.* Ascomata hysterioidea. Sporidia fusca. Cfr. Syll. II, p. 757].

Sect. 5. Scolecosporæ. Sporidia filiformia, continua v. septata.

pg.601 x *Godronia.* Ascomata subcoriacea, urceolato-calyciformia.

Pocillum. Ascomata clavato-turbinata, fibrosa, disco truncato. *pg. 605*

Crinula. Ascomata clavata, stipite corneo, hymenio submucoso, initio convexo conidiophoro, demum disciformi ascophoro. *pg. 606*

Sect. 1. Hyalosporæ Sacc.

MIDOTIS Fries. Syst. Orb. Veg. p. 363 (1825), Elench. II, pag. 29 (Etym. *Mida* et *ous*, *otos* auris), *Wynnea* Berk. et Curt. F. Mex. p. 424. — Ascomata elongata, auriformia, coriacea, sursum divisa, basi in stipitem plus minus distinctum connata. Hymenium inferum ab excipulo subiude discretum. Asci teretes, octospori, paraphysati. Sporidia ovoideo-oblonga, continua, hyalina.

1. **Midotis gigantea** (B. et C.) Sacc. *Wynnea gigantea* Berk. et Curt. 2255 Fung. of Mexic. n. 5, t. XII, Cooke Myc. f. 93. — Stipite communis subcylindrico, crasso, rugoso, suberoso, intus albo, sursum ramoso, in cupulas auriformes multiplices quandoque proliferas diviso; cupulis extus glabris, marginibus inflexis, cum hymenio rufis; ascis subcylindricis, octosporis; sporidiis magnis, subcymbiformibus, 25×14 , utrinque obtusis.

Hab. in lignis putridis? in Mexico. — Stipes communis 7-8 cm. long., 18 mm. cr., cupulæ 6-7 cm. lat.

2. **Midotis macrotis** (Berk.) Sacc. *Peziza* Berk. Dec. fung. n. 356, 2256 *Wynnea* Berk. Journ. Linn. 1886, p. 424, Cooke Myc. f. 94. — Cupulis elongatis, obliquis, auriformibus, basi connato-ramosis, coriaceo-subcartilagineis, hepatico-rufis, extus glabris, hymenio purpurascente; ascis cylindricis, filiformi-paraphysatis, octosporis; sporidiis ellipsoideis, hyalinis, 25×12 .

Hab. in ligno putri in India et Mexico. — Ascoma 12-13 cm. alt.

3. **Midotis heteromera** Mont. Syll. Crypt. n. 642, *Peziza* Mont. Cent. 2257 II, n. 18, t. 6, f. 3. — Fasciculata, basi ramosa aut prolifera, coriacea, fragilis, stipitata; stipite brevi ramoso rubiginoso strigoso, in cupulam dilatato inæqualem, uno latere versam, subauriformem, disco umbrino ascophoro; ascis filiformi-clavatis, octosporis; sporidiis oblongis cymbiformibus, uniseriatis, $10 \times 3,5$.

Hab. ad ligna in silvis sinnamariensibus Guianæ (LEPRIEUR), S. Domingo et Cuba (BERK.).

4. **Midotis verruculosa** B. et C. Cub. Fung. n. 705. — Cupula obliqua 2258 extus lateritia verrucosa; hymenio livido-fusco; sporidiis 10×5 .

Hab. in ligno putri in insula Cuba. — A *M. heteromera* dif-

fert superficie verruculosa nec tantum furfuracea sporidiisque crassioribus.

5. **Midotis regularis** Cooke et Phil. Grev. IX, pag. 109. — Erum-²²⁵⁹
pens; stipite brevi, ramoso; ramulis brevissimis; cupulis concavis,
fuscis, punctatis; margine infexo; hymenio obscuriore levi; ascis
cylindraceis; sporidiis ellipticis 10-13 \approx 4-5; paraphysibus linea-
ribus.

Hab. in ligno putri, Rio Janeiro Brasiliæ.

6. **Midotis plicata** Phill. et Harkn. Calif. p. 4. — Ascomatibus obli-²²⁶⁰
quis, externe granulosis, brunneo-nigrescentibus, hymenio conco-
lori, subplicato; ascis cylindraceis, octosporis; sporidiis ovato-el-
lpticis, leniter curvatis, binucleatis, 7 \approx 5-7; paraphysibus non visis.

Hab. in ligno Umbellulariæ californicæ in California.
— Affinis *M. regulari*, a qua differt sporidiis minoribus, curva-
tis, binucleatis atque disco longitudinaliter plicato.

7. **Midotis Lingua** Fr. Elench. II, p. 29. — Solitaria, suberecta, ob-²²⁶¹
ovato-oblonga, utrinque plana, margine leviter deflexo, 5 cm. usque
alta, stipite brevissimo laterali instructa; stipite 2-4 mm. longo,
glabro sublaccato-nitente fragili, margine ascomatis rufo-badio ab
hymenio distinto, sed in pileum abeunte, pagina sterili, levi,
glabra, azona, badio-fusca, subfuliginosa, subinde laccato-nitente;
contextu floccoso, albo, primo molli, sicco cartilagineo; hymenio
molli, absque papillarum vestigio, nigricante, siccitate rimoso-fo-
veolato, sub lente subpruinoso (unde sporidia haud elastice explodi-
videntur).

Hab. ad bases truncorum in Vallisia (SCHLEICHER). — Fingas
Otideam leporinam resupinatam, sic ejus habebis ideam.

8. **Midotis Patella** Fries. Nov. Symb. Mant. p. 231. — Cupularis, ses-²²⁶²
silis, postice oblique adnata, supra glaberrima cervina, hymenio
concavo brunneo; ascis exakte linearibus, octosporis; sporidiis el-
lipsoideis, ut plurimum biocellatis.

Hab. ad trunhos prostratos in Mexico (LIEBMANN). — Cupula
coriacea *Auriculariæ* instar, 25 mm. lat. Hymenium in adultis a
cupula facillime secedens. An hujus generis?

URNULA Fr. Sunima p. 364 (1849) [Etym. *urna*] ¹⁾. — Ascomata sti-

1) Teste cl. Cooke in litt. hoc ducenda **Urnula rhytidia** (Berk.) seu
Peziza rhytidia Berk. (supra n. 331) ob excipulum extus fibrosum brunneum,
medie fibrosum virens, interius compactius, denique pallidius, ascophorum.

pitata, urceolata subclausa, dein ore rotundo vel laciniato dehiscentia, coriacea, fusca vel furfuracea vel pubescentia. Asci cylindracei, octospori, paraphysati; sporidia ellipsoideo-oblonga, continua, hyalina.

1. **Urnula Craterium** (Schw.) Fr. Novæ Symb. p. 122, *Peziza* Schw. 2263

Carol. p. 91, t. I, f. 7-11, Fr. Syst. Myc. II, p. 74, *Cenangium* Fr. Elench. II, p. 21. — Maxima, longe stipitata, subcæspitosa, nigro-badia, subclausa, intus cava, subiculo tomentoso; ascis longissime cylindraceis, 8-sporis, 400-500 \approx 14-25; sporidiis monostichis oblongis hyalinis, 23-33 \approx 10-13; paraphysibus ramosissimis, septatis.

Hab. ad ramos dejectos in Carolina, Ohio et in America arctica (DRUMMOND) et in Hungaria. — Vegetatione superficiali ab *Cenangiis* differt, sed velo pulveraceo convenit. Passim laciniatim dehiscit. Cfr. Hedw. 1883, p. 182, ex quo patet *P. adustam* Schulz. et *P. lugubrem* Kalchbr. huc referendas esse.

2. **Urnula Hartii** Berk. Fung. Arct. Exped. p. 16. — Capula cyathiformi, extus cum stipite spadiceo velutina; margine infexo; hymenio pallidiore; sporidiis ellipticis binucleatis, monostichis, 15 \approx 7.

Hab. inter muscos Upernavik in regione arctica (H. C. HART et FEILDEN). — Ascoma 6 mm. lat., stipes 4 mm. alt. Affinis *U. Craterio* sed minor. Habitus *Geopyxidis Ciborii*.

3. **Urnula minor** Fr. Nov. Symb. p. 122. — Coriacea, subfasciculata, primo turbinato-clavata, extus albo-pruinosa, stipite brevi nigro, dein dehiscens, ore obliquo lacero dentato, disco urceolato pallido.

Hab. ad terram in Guinea (AFZELIUS). — Ab *U. Craterio* differt statura multiplo minori, cupula turbinata, extus e pruina albida et, quantum e paucis speciminibus videre licet, defectu insignis mycelii byssacei. Stipes admodum tenuis, 2 mm. paullo ultra longior et sensim (nec abrupte ut in *U. Craterio*) in clavam dilatatus, nudus, niger, subinde pauci basi juncti; cupula membranaceo-coriacea, primo omnino clausa, dein ore indefinito lacera hiascens, ore obliquo dentato-lacero, extus e pruina pallida. Asci et sporidia nulla adsunt, sed floccorum s. paraphysium densa congeries exacte similis *U. Craterii*.

4. **Urnula Microcrater** (Hazsl.) Sacc. *Craterium Microcrater* Hazsl. 2266

Eing. Disc. p. 17 et Magyar. Szab. Disc. t. 9, f. 31. — Ascomate clavato-piriformi, stipitato, apice truncato, excavato, initio clauso,

demum margine erecto fisso aperto, atro-brunneo, extus furfuraceo; stipite brevi, cylindraceo, basi atro-radiculoso; ascis cylindricis, substipitatis, filiformi-paraphysatis, octosporis; sporidiis oblongo-ellipsoideis, monostichis, hyalinis, levibus.

Hab. in ramis terra infossis, mycelio atro-brunneo insidens, in Hungaria et Slavonia. — Totus fungus 10-12 cm. alt., 5-6 cm. lat.

5. **Urnula terrestris** (Niessl) Sacc. *Podophacidium terrestre* Niessl 2267
Beitr. p. 163, t. VII, f. 56. — Gregaria; ascomata substipitato, turbinato vel sycioideo, 2-3 mm. diam., 1-2 mm. alt., coriaceo-membranaceo, badio, primum clauso demum laciniato, disco undulato, sulphureo; ascis clavatis, inferne attenuatis stipitatis, apice late rotundatis, 8-sporis, 124-136 = 8-9; sporidiis monostichis, oblongis, inaequilateralibus, simplicibus, continua vel interdum nucleo 1-2 diviso, hyalinis, 11-13 = 4-5; paraphysibus ascorum longitudine, apice bifurcatis.

Hab. ad terram nudam in silvis prope Gratz Stiriae. — A typo *Urnulae* vix stipite breviore distingui videtur.

DERMATEA Fr. Summa Veg. p. 362 (1849), [Etym. *derma* corium].

— Ascomata erumpenti-superficialia, urceolato-scutellata, mox aperta, subtuberosa, saepius e basi communi cæspitosa. Ascii paraphysati, octospori. Sporidia ellipsoidea vel oblonga, continua, hyalina. — Hinc *Peziculæ* illinc *Cenangio* affinis, a prima substantia magis suberosa, ab altera substantia vix coriacea, colore plerumque lætiore, disco magis aperto, sporidiis saepius majoribus differet; sed revera species plures hucusque ambiguæ, genera haud acute limitata demostrant.

1. **Dermatea Cerasi** (Pers.) Fr. Summ. Veg. Scand. p. 362, Tul. Carp. 2268 III, p. 156, t. XIX, f. 13-16, Karst. M. F. I, p. 223, *Peziza* Pers. Disp. p. 35, *Cenangium* Fr. Syst. Myc. II, p. 180, Sacc. F. it. fig. 1309. — Stromate cortici immerso, carnosu-suberoso, continuo, e parenchymate luteo-virente et fusco-virgato composito, crasso; cupulis per epidermidem erumpentibus, solitarie vel saepius gregatim stromati insidentibus, primitus planiusculis margineque elevato sinuoso-repando, demum plus minus convexis et margine reflexo vel evanescente, furfure citrino-virente conspersis disco e cæsio-virente fusco vel tandem nigrescente, suberosis, 2-4 mm. lat.; ascis cylindraceo-clavatis, 85-90 = 10, 8-sporis apice jodo non tinctis; sporidiis plerumque distichis, oblongatis vel aciculari-ob-

longatis, curvulis vel subrectis, eguttulatis, chlorinis, 14-20 \approx 4-5,5; paraphysibus filiformibus, conglutinatis, apice clavato, fuscidulo v. subfusco.

Hab. in ramis exsiccatis Cerasorum in Suecia, Fennia, Gallia, Germania, Italia, Belgio, Austria, Hungaria, Britannia, Amer. bor. — Status secundarii: *Micropera Drupacearum* Lév. var. *Cerasi* et *Dothichiza dubia* (Pers.).

2. **Dermatea pulcherrima** Fuck. Symb. App. II, p. 56. — Cupulis cæ-
spitosis, subcoriaceis, 2-8 mm. latis, subsessilibus, primo concavis,
demum magis explanatis, margine involuto, extus pulveraceis, pul-
cherrime læte flavo-viridibus, disco flavo-fusco; ascis stipitatis,
cylindraceis, 8-sporis, 94 \approx 8; sporidiis oblique monostichis, fusi-
formibus, curvatis, continuis, biguttulatis, hyalinis, 14 \approx 3; para-
physibus simplicibus, filiformibus.

Hab. in ramis Cerasi avium pr. Hattenheim Rhenogoviæ. —
A *D. Cerasi* ascomatibus maximis distinguenda.

3. **Dermatea Padi** Fr. Summa Veg. p. 362, Tul. Carp. III, p. 160, 2270
Karst. M. F. I, pag. 224, *Peziza Cerasi* β. *Padi* A. et S. Lus.
p. 345. *Cenang. fallax* Fr. — Cæspitosa vel subsolitaria, erum-
pens, epidermide cincta, globosa clausa, mox subpatellaris, disco
plano olivaceo-nigricante, extus rufo-olivaceo-furfuracea; spori-
diis ellipsoideo-oblongis, utrinque 1-guttatis, hyalinis.

Hab. in ramis emortuis Pruni Padi in Suecia, Fennia, Ger-
mania, Gallia. — Status secundarii: *Micropera fallax* (Wahl.)
[Cfr. Karst. loc. cit.], *Dothichiza Padi* S. et R. et *Sphaeronema*
brunneo-viride Fuck.

4. **Dermatea Ariæ** (Pers.) Tul. Cap. III, p. 160, *Peziza* Pers. M. E. 2271
I, p. 325, *Tympanis* Fr. Syst. Myc. II, pag. 175 et Elench. II,
p. 18, *T. inconstans* Fr. Summa p. 400, *Cenangium inconstans*
Fuck. Symb. pag. 268, *Dermatea* Tul. — Ascomatibus sessilibus,
sparsis, duris, nigris, primum subrotundis clausis, dein vertice dilata-
tatis orbiculato-deplanatis, ambitu marginante subelevato cinctis
majusculis subgregariis; ascis cylindraceo-clavatis, apice incrassa-
tis, 8-sporis, 70-90 \approx 8-9; sporidiis oblongis curvulis, 2-guttula-
tis, hyalinis, 10-14 \approx 3-4; paraphysibus sursum furcatis, septatis,
clavulatis, 6 μ . cr.

Hab. ad cortices Sorbi Ariæ, S. Aucupariæ et Mespili
in Suecia, Germania, Gallia, Helvetia. — Status pycnidicus est
Micropera Sorbi (Fr.) Sacc. seu *Sphaeria conica* A. et S.

5. **Dermatea Cydoniæ** Schw. Syn. Am. bor. n. 1973. — Sparsa, sim- 2272

plex, sessilis, applanato-adpressa, difformis aut subrotundo-undulata, extus furfure albido crasso pulverulento tecta, marginibus tumidis, arcte inflexis; disco fusco-badio aut subcinnamomeo 4-6 mm. lat.

Hab. rara in emortuis ramis *Cydoniæ*, Bethlehem Amer. bor.

6. **Dermatea Corni** Phil. et Hark. Grev. XIII, p. 22. — Cæspitosa v. 2273 sparsa, minuta, primo tecta, demum erumpens, turbinata, furfuracea, vitellina, margine distincto, disco plano vel subdepresso, obscuriore; ascis late clavatis; sporidiis oblongo-ellipticis, 25-30 = 8; paraphysibus filiformibus, superne clavato-incrassatis, furcatis.

Hab. in ramis *Corni* in California (HARKNESS).

7. **Dermatea Pseudoplatani** Phill. in Grev. XVII, p. 45. — Cæspito- 2274 sa, erumpens, sessilis v. substipitata; disco initio convexo dein subdepresso, albo demum pallide flavo-brunneo; ascis late clavatis 8-sporis; sporidiis oblongis vel oblongo-ellipsoideis, triguttatis demum (spurie?) triseptatis 15-17 = 5-7; paraphysibus apice clavatis.

Hab. in cortice *Aceris Pseudoplatani*, Spark Hill Britanniae (W. H. GROVE). — Ascoma $\frac{1}{2}$ -1 mm. lat. *Derm. lividæ* proxima videtur. Colore albo mox distincta species. «*Potius Dermatella*».

8. **Dermatea olivacea** Ellis Bull. Torr. Cl. 1877, p. 133. — Ascoma- 2275 tibus cæspitosis, erumpentibus, in stipitem brevem desinentibus, olivaceo-furfuraceis, disco plano, margine obtuso; ascis latis obtusis, breviter stipitatis, 125-135 = 20-25; sporidiis oblongis, distichis, obtusis, 35 = 12-15, primo protoplasmate griseo dein saturatiore farctis; paraphysibus linearibus, crispulis, simplicibus vel parce ramosis, apicibus vix incrassatis.

Hab. in ramis *Ilicis glabræ* emortuis, South Jersey Americæ borealis.

9. **Dermatea Xanthoxyli** Peck 31 Rep. St. Mus. p. 47. — Ascomati- 2276 bus dense cæspitosis, minutis, mutua pressione sæpe inæqualibus, brunn eo-lilacinis extus margineque albis et villoso-pruinosis, disco piano vel concaviusculo; ascis brevibus clavatis; sporidiis stipatis continuis subcylindricis, leniter curvis, obtusis, hyalinis.

Hab. in ramis emortuis *Xanthoxyli americanæ*, West Troy Amer. bor. — Cæspituli vix 2 mm. lat. sæpe ex cupulis 10-12 compositi.

10. **Dermatea acicola** Briard et Sacc. Fung. Alg. et Gall. n. 15 in 2277 Rev. Mycol. 1885. — Ascomatibus erumpentibus, gregariis obconicis, tenacellis, minutis, $\frac{1}{4}$ mm. diam., totis nigris, glabris disco sub-

plano, breve marginato; ascis clavatis, breve stipitatis, subtuber-
losis, 120-144 \approx 22-24, octosporis; paraphysibus filiformibus, ramu-
losis, sursum leniter incrassatis obvallatis; sporidiis ellipsoideo-ob-
longis, continuis granulosis 1-3 guttulatisque, dein eguttulatis, 20-
26 \approx 12-14, hyalinis.

Hab. in pag. sup. foliorum Juniperi communis pr. Troyes Galliæ (BRIARD). — Etsi pro genere minuta species, tamen ab hujus typo non abhorret:

11. **Dermatea juniperina** Ellis Amer. Nat. 1883, p. 193. — Erumpens, 2278
sessilis, orbicularis, fuliginoso-nigra, disco udo vix pallidiori, mar-
gine obsoleto circiter $\frac{1}{4}$ mm. crass. sicca deorsum contracta et
fere stipitata; ascis clavato-cylindraceis, 100-114 \approx 15-18; para-
physibus filiformibus superne vix incrassatis; sporidiis subdistichis,
ellipsoideis, subhyalinis, crasse uninucleatis, 18-20 \approx 7-12.

Hab. in foliis emortuis Juniperi communis, Decorah Iowa Amer. bor. (G. W. HOLWAY).

12. **Dermatea Pini** Phill. et Hark. Grev. XIII, p. 22. — Cæspitosa 2279
vel sparsa, subsessilis, globosa, demum patelliformis, ceraceo-co-
riacea, furfuracea, ochraceo-incarnata, margine indistincto, disco
concolori, applanato; ascis clavatis, octosporis; sporidiis oblongo-
ellipticis, 3-4 nucleatis, 17-20 \approx 6-8; paraphysibus linearibus, su-
perne incrassatis, furcatis.

Hab. in cortice Pini in California (HARKNESS).

13. **Dermatea phyllophila** Peck 31 Rep. Stat. Mus. p. 47. — Asco- 2280
matibus minutis suborbicularibus, margine flexuoso, siccis, subfi-
brosis brunneis, extus pilosellis, erumpentibus, epidermide rupta
cinctis et partim tectis, udis expansis et discum planum, pallidum
vel sordide album revelantibus; ascis oblongo-clavatis, obtusis ses-
silibus; sporidiis late ellipsoideis, subhyalinis 7-8 \approx 5-6 plerumque
uninucleatis; paraphysibus apice incrassatis ascos leniter supe-
rantibus.

Hab. in foliis emortuis sed adhuc stantibus Abietis balsa-
meæ Summit Amer. bor. — Discus udus pruinosis.

14. **Dermatea cinnamomea** Cooke et Peck Disc. U. S. II, p. 24 et II, 2281
p. 28, Rep. Stat. Mus. p. 67 nec DC. (*Pezicula*). — Cupulis sub-
cæspitosis erumpentibus, subsessilibus, epidermide rupta cinctis, sub-
coriaceis, extus pulverulentis cinnamomeis, margine involuto, disco
subplano sicco leniter anguloso brunneo; ascis elongato-clavatis,
sporidiis anguste ellipsoideis, continuis 12-13 μ . longis.

Hab. in ramis emortuis Populi, Sandlake Amer. bor.

15. **Dermatea australis** (Speg.) Sacc. *Cenangium australe* Speg. F. 2282

Fueg. n. 345. — Stromatibus cortice innatis, sparsis v. laxe gregariis, primo peridermio tectis, dein, eo fisso, transverse erumpentibus, ellipticis 3-5 mm. long., 1-3 mm. lat. vel pulvinatulis 1 mm. crass., superne applanatis truncato-marginatis, fusco-purpureis, saepius pulvere subviolascente inspersis; cupulis stromate immersis, primo clausis (an medio minute papillulatis?), dein applanato-expansis 0,5-1 mm. diam., saepe subconfluentibus, non v. parce obtuseque marginatis, non v. vix exsertis, e violascenti purpurascensibus; ascis cylindraceis, antice truncato-rotundatis, postice breviter crasseque attenuato-pedicellatis, 90-180 \approx 14-18 octosporis; paraphysibus filiformibus longioribus dense obvallatis; iodi ope tunica ascorum apice dilute violaceo-cœrulecente; sporidiis transverse v. oblique v. recte monostichis, ellipticis saepius utrinque obtusatis 16-24 \approx 7-14, crasse tunicatis, granuloso-farctis v. grosse 2-guttulatis, primo hyalinis dein chlorinulis v. dilute olivascentibus.

Hab. vulgata ad ramos dejectos putrescentes Fagi betuloidis in silvis in Staten island et Fuegia. — Species fabrica naturaque melius Bulgarieis adscribenda, vegetatione tamen cenangoidea donata. An novum genus?

16. **Dermatea lobata** Ellis Bull. Torr. Cl. 1876, p. 108. — Superfi. 2283
cialis sparsa, globosa, primo clausa, dein expansa, aperta et disco convexo, margine lacerato-lobato, extus rufo-brunnea atque rugosa, squamulis innatis vestita; ascis cylindraceis; paraphysibus superne guttulato-incrassatis; sporidiis monostichis, breviter oblongis vel ovoideis, subhyalinis, 12-15 \approx 10.

Hab. in ramis Quercus albæ dejectis sed nondum putridis Newfield, New Jersey Amer. bor.

17. **Dermatea tabacina** Cooke Disc. U. S. II, p. 24, Rehm Ascom. 2284
n. 359. — Ascomatibus cæspitosis, erumpentibus, stipitatis, primo subglobosis, dein cupulatis, subinde coriaceis, margine infexo, lateraliter compressis castaneo-brunneis, vix furfuraceis; disco pallido, in sicco hysteriformi; ascis clavatis, 10 \approx 18; sporidiis elongatis, obtusis, subellipsoideis, distichis rectis vel curvulis, 1-cellularibus, hyalinis, 7,5 \approx 6; paraphysibus crassiusculis, articulatis, 5 μ . crassis, sursum brunneis clavulatis.

Hab. in Quercus albæ ramis vivis Unit. Stat. Amer. bor.
— Sec. Cooke sporidia 10-11 \approx 7, fusiformia, intus granulosa.

18. **Dermatea purpurascens** E. et E. Jour. Myc. 88, p. 100. — Erumpens, sessilis v. substipitata $1\frac{1}{2}$ - $3\frac{1}{4}$ mm. alta, sparsa vel ascomati-

bus 2-4 aggregatis, disco convexo vel plano, subimmarginato, $\frac{3}{4}$ -1 mm. diam., e sordide albo rufo-purpurascente extus obscurior, inferne in basim stipitiformem contracta; ascis clavato-cylindraceis, 115-120 \approx 18-22; paraphysibus crassis superne vix incrassatis; sporidiis distichis oblongis, leniter curvulis, 25-30 \approx 8-11, utrinque obtusis, crasse 4-nucleatis subhyalinis.

Hab. in ramulis emortuis Castaneæ, West Chester Amer. bor. — Affinis *D. olivaceæ*, a qua differt in primis colore disci.

19. **Dermatea pruinosa** E. et E. Jour. Myc. 88, p. 100. — Gregaria, 2286 sessilis vel subsessilis, 2-2,5 mm. crass. sordide brunnea, extus albo-pruinosa, disco livido-albo; ascis cylindraceis, 55 \approx 5; paraphysibus filiformibus simplicibus vel ramosis; sporidiis subdistichis allantoideis, hyalinis, 8-10 \approx 2- $2\frac{1}{2}$.

Hab. in cortice emortuo, Colorado Amer. bor. (D. A. COCKE-RELL). — Affine *D. fusco-sanguineæ*, a qua differt colore, magnitudine atque forma sporidiorum.

20. **Dermatea inclusa** Peck. 30 Rep. Stat. Mus. pag. 62. — Minuta, 2287 sparsa, erumpens, sessilis, epidermide rupta, arcte cincta, marginata, margine farinoso v. furfuraceo; disco-plano v. concavo subochraceo; ascis latis oblongo-cylindraceis; sporidiis distichis vel inordinatis, oblongo-ellipsoideis saepe curvulis hyalinis, 28-35 μ . long.

Hab. in truncis emortuis Salicis Maryland Amer. bor. — Sporidia 1-nucleata v. granulosa. Fungus uodus intus rhabarbarinus.

21. **Dermatea rufa** Cooke Grev. VIII, pag. 72. — Sparsa, erumpens, 2288 fusco-rufa, carnosocoriacea; cupulis in stipitem productis, disco excavato, rufo, 0,12-0,3 mm. diam., extus farinaceo, aureo-fusco; carne concolori; ascis clavatis; sporidiis fusiformibus, rectis vel curvulis, 35-40 \approx 5-6, quadrinucleatis, hyalinis; paraphysibus filiformibus.

Hab. in corticibus, Natal Africæ austr. — Madefacta colorem aureo-brunneum exsudat.

22. **Dermatea carnea** C. et Ellis Grev. V, p. 32, t. 75, f. 9. — Cæspitosa, 2289 minuta, ochraceo-carnea, erumpens, substipitata, turbinata demum excavata, extus furfuracea, pallidior; ascis clavatis, octosporis; sporidiis ellipsoideis, saepe (genuine?) 3-septatis, hyalinis, 20-22 \approx 8; paraphysibus filiformi-clavulatis.

Hab. in ramis corticatis Aceris rubri, Viburni, Carpinii americanæ, New Jersey et West Albany Amer. bor. — Ascoma 0,2-0,5 mm. lat. «An rectius *Dermatella*?»

CENANGIUM Fr. Syst. Myc. II, pag. 177 ex p. (1822) [Etym. *cenos* cavus et *angeion* vas), *Dermatea*, *Phibalis*, *Encœlia* Auct. ex p. — Ascomata erumpenti-superficialia, saepius ex stromate communis cæspitosa, globulosa, clausa, dein aperta urceolata vel scutellata, coriacea v. subcornea extus typice furfuracea, plerumque fusca v. atra. Ascii paraphysati 4-8-spori. Sporidia oblonga v. fusoidea, continua, hyalina. Sæpe adsunt pycnidia et spermogonia. — Cum gen. *Dermatea* confluit, hinc species etiam sub hoc genere quærendæ.

I. *Eu-Cenangium*. Ascomata minora, duriora.

1. **Cenangium Prunastri** (Pers.) Fr. Syst. Myc. II, pag. 180, *Peziza* 2290

Pers. Tent. pag. 35, *Dermatea* Fr. Summa p. 362, Rehm Disc. p. 261, *Tympinis* Wallr. Crypt. p. 427. — Subcæspitosum, subcorneum, nudum, nigricans, cupulis primo subulæformibus, dein apertis concavis substipitatis; ascis clavatis, 8-sporis, 60-80 = 12; sporidiis oblongis vel elongato-ovatis, obtusis, 1-cellularibus, 1-2 nucleis magnis præditis (an demum 2-cellularibus?), hyalinis, distichis, 12-14 = 4-4,5; paraphysibus filiformibus, superne furcatis.

Hab. in ramis siccis Pruni domesticæ, spinosæ, Americanæ, Armeniacæ in Suecia, Germania, Gallia, Hungaria, Britannia, Belgio, Italia, Amer. bor. — Status pycnidicus est *Sphaeronema spurium* (Fr.) Sacc. Etiam transversim erumpens, subcæspitosum et primitus sphæriæforme, adultum profunde excavatum, vix 2 mm. latum; hymenium virens.

2. **Cenangium polygonium** Fuck. Symb. App. II, p. 55, p. p. *Dermatea polygonia* Rehm Disc. p. 263. — Cupulis ut in *Tymp. polygonia* (quam cfr.); ascis tereti-clavatis 110-120 = 20, 8-sporis; sporidiis oblongo-ellipsoideis obtusis, rectis, continuis, hyalinis, 15-20 = 6-8, distichis; paraphysibus furcatis, apice elliptico-clavatis, brunneis, 5 μ. cr.

Hab. in ramis siccis Piri Mali pr. Oestrich Rhenogoviæ. — An forma octospora *Tymp. polygoniae*?

3. **Cenangium Syringæ** (Rehm) Sacc. *Dermatea Syringæ* Rehm Ascom. 2292

Lojk. n. 58 (1873). — Plerumque gregarium, ex rimis epidermidis erumpens, primitus globosum, dein irregulariter expansum, rugoso-marginatum, fuscum, 0,5-1,5 mm. diam.; ascis clavatis, 8-sporis, 50-60 = 5; sporidiis ellipticis vel oblongato-ovoideis, 1-cellularibus, saepè 2-nucleatis hyalinis, distichis, 8-9 = 2-2,5; paraphysibus superne clava fusca 4-5 μ. cr. instructis.

Hab. ad ramum siccum *Syringæ vulgaris* prope Plavise-vicza com. Szörény in Hungaria (LOJKA).

4. **Cenangium pulveraceum** (A. et S.) Fr. *Syst. Myc.* II, p. 181, *Peziza* Alb. et Schw. *Lus.* pag. 342, t. VIII, f. 2, *Phibalis* Wallr. *Crypt.* n. 2408. — Ascomatibus gregariis nigricantibus, extus scobbe furfuracea cinerea obtectis, primum subglobosis, dein in basin stipitiformem productis turbinatis, vertice semipatulis, 1 mm. lat.; disco flavidō, plano; ascis clavatis 8-sporis; sporidiis elongatis, hyalinis, continuis, 5-8 = 2; paraphysibus filiformibus (Phill.).

Hab. ad corticem Betulæ, Pruni, Ilicis, Fagi in Germania, Belgio, Britannia, Ceylon Amer. bor. — Cfr. Rehm Disc. p. 229. Sec. Fuck. ascomata adsunt spermogonica sporulis allantoideis 4 = 1 referta.

5. **Cenangium vagabundum** (P. et B.) Sacc. *Tympinis vagabunda* 2294 Pass. et Belt. *Fung. Sic.* n. 17. — Cupulis sparsis, solitariis, erumpentibus, sessilibus vel turbinato-substipitatis, corneis, primo clausis, dein apertis, margine involuto, rubiginoso, fuscis, nitidis striatulis; ascis cylindrico-clavatis, basi longe attenuatis, jodo immutatis, 112-125 = 10-12, sed saepe multo brevioribus, 8-sporis (?); sporidiis oblongo-ellipticis, biguttulatis, hyalinis, 12 = 5 in ascorum lumine plerumque imperfecte perspicuis ibique guttulis eorum sporidia globosa numerosa simulantibus; paraphysibus filiformibus stipatis ascos æquantibus, apice subcoalitis, sed non fucatis.

Hab. in ramulis aridis dejectis Rosæ, Rubi, Pistaciæ Terebinthi prope Monostalla Siciliæ.

6. **Cenangium bicolor** (Ell.) Sacc. *Tympinis bicolor* Ellis Amer. Nat. 2295 1883, pag. 193. — Cæspitosum v. simplex, sessile, margine angusto, subundulato, extus pallide rufum, disco atro plano, $\frac{3}{4}$ -1 mm. diam.; ascis subcylindraceis, 75 = 8-9; paraphysibus obtusis, sursum indistincte incrassatis; sporidiis uniseriatis, oblongo-ellipsoideis, saepe uno apice acutis, 2-4-nucleatis, lutescentibus, 13-15 = 3,5-4.

Hab. in ramulis emortuis (*Aceris?*), Decorah, Jowa Amer. bor.

— Affine *Tymp. acerinae*.

7. **Cenangium clandestinum** Rehm Disc. p. 223. — Erumpens, seriatim cæspitosum, globosum dein scutellatum extus atro-brunneum, margine pallidore, siccum nigrum et convolutum, 0,5-1 mm. lat., ceraceo-coriaceum; disco plano griseo; ascis clavatis 45-50 = 6-8, octosporis; sporidiis ellipsoideis obtusis rectis curvulisve continuis hyalinis, distichis 9-11 = 3; paraphysibus filiformi-clavulatis,

Hab. in ramulis siccis *Aceris campestris* in Franconia Bavariæ.

8. **Cenangium Magnoliæ** B. et C. North Am. Fung. n. 766. — Cæ-
spitosum, apertum marginatum, nigrum; ascis amplis; sporidiis ma-
gnis allantoideis, 25-30 = 12-15.

Hab. in ramis *Magnoliæ glaucæ*, *Perseæ* et *Lauri*, Ala-
bama, Darien, Gainesville Amer. bor.

9. **Cenangium? Sambuci** (Cr.) Sacc. *Tympanis Sambuci* Cr. Finist. 2293
pag. 43. — Ascomatibus 1 mm. diam., lutescentibus, translucidis,
sessilibus subcylindraceis; ascis dehiscentibus, octosporis; sporidiis
subfusiformibus, parum curvatis.

Hab. in ramulis emortuis *Sambuci*, Finistere Galliæ.

10. **Cenangium Sophoræ** Pass. Erb. critt. ital. II serie, n. 1173. — 2299
Cupulis cæspitosis, minutis erumpentibus, atris; ascis oblongo-
clavatis, 8-sporis; sporidiis distichis, elliptico-oblongis, rectis vel
curvis, minutis, hyalinis; paraphysibus non visis.

Hab. in ramis siccis *Sophoræ pendulæ* in Horto botanico
Parmæ Ital. bor.

11. **Cenangium Sarothamni** Fuck. Symb. p. 270. — Cupulis gregariis 2300
cæspitosisve, substipitatis, primo clausis, umbilicatis, demum apertis,
applanatis, extus fuscis, subpulveraceis, demum atris, 2 mm. latis,
disco nigro-fusco, margine elevato, atro cinctis; ascis oblongis,
stipitatis, antice obtusis, seorsum attenuatis, 8-sporis, 18 = 8-9;
sporidiis subdistichis oblongo-ovatis, hyalinis, continuis, 8-10 = 3-4.

Hab. in ramulis siccis *Sarothamni scoparii* pr. Eberbach
Rhenogoviæ et in Arduennis (FUCKEL, LIBERT).

12. **Cenangium Andromedæ** (Schw.) Fr. Syst. Myc. II, p. 282, *Peziza* 2301
Schw. Carol. n. 1261. — Solitarium, minutum, nudum, nitens, ni-
gerrimum, cupula subglobosa, stipite longiusculo; sporidiis fusoi-
deis vel sigmoideis, tenuibus, pluriguttulatis (Berk.).

Hab. in cortice *Andromedæ arboreæ Carolinæ*. — Spar-
sum, interdum etiam fasciculatum, magnitudine capitis acicularum
minimarum. Substantia fere cornea. Cupula sphæriæformis, mar-
gine valde involuto, disco urceolato. *Tympanidi* affine.

13. **Cenangium Vaccinii** (Fuck.) Sacc. *Dothiora* Fuck. Symb. App. III, 2302
pag. 29. — Patellis erumpentibus, demum valde prominentibus,
gregariis, ramulos plerumque totos occupantibus, fusco-nigris, 1 mm.
diam., rotundatis sed plerumque ellipticis irregularibusque, mar-
gine elevato, tumido, vertice concavis, primo carbonaceis concolo-
ribus, demum irregulariter fissis et disco sordido exposito; ascis

Discomyceteæ, Dermateæ, Hyalosporæ, Cenangium. 559

clavatis, 8-sporis, 74-12; sporidiis farctis, oblongo-ovatis, utrimque obtusis, parum curvatis, continuis, hyalinis, 12-6 mm. crass.

Hab. in caulis ramisque siccis Vaccinii uliginosi, S. Moritz Ober-Engadin Helvetiæ. Habitus hinc *Sporomegatis degenerantis*, illinc *Dothioræ ellipticæ*, sporidiis vero diversissimum.

14. **Cenangium cæspitosum** (Fuck.) Rehm Disc. p. 223 *Dermatea* 2303
Fuck. Symb. p. 277. — Cupulis erumpentibus, dense cæspitosis, difformibus, usque ad 2 mm. latis, extus margineque cinereo-pulverulentis, asperis, disco atro, excavato; ascis linearibus, 8-sporis, 60-8; sporidiis cylindraceis, rectis, continuis, hyalinis, 10-2.

Hab. ad ramos corticatos Coryli Avellanæ. — Habitus *Cen. populnei* sed minus.

15. **Cenangium Coryli** Corda Ic. V, p. 78, t. IX, f. 63. — Aggregatum, superficiale, minutum; cupulis globulosis, dein applanatis, substipitatis discoque nigricantibus; ascis clavatis; paraphysibus tenuissimis obvallatis; sporidiis senis (semper?) oblongo-ellipsoideis, hyalinis, 9-10 μ . longis.

Hab. in ramis emortuis Coryli in Bohemia. — Sec. Kickx Crypt. Fl. I, pag. 473 stat. pycnidicus est *Catinula turgida* (Fr.). An forma oligospora *Tymanidis corylinæ?*

16. **Cenangium sticticum** (B. et C.) Sacc. *Tymanidis stictica* B. et C. 2305
North Am. Fung. n. 763. — Punctiforme; cupulis concavis, margine elevato; ascis clavatis; sporidiis fusiformibus, curvatis, quadrinucleatis.

Hab. in ramis Salicis babylonicae et forte Pini et Quercus in Amer. bor.

17. **Cenangium glabrum** (Wallr.) Rehm Disc. p. 225, *Phibalis glabra* 2306
Wallr. Crypt. n. 2412. — Ascomatibus hypophloeodeis, subsolitariis, mox epidermidem scissam superantibus sessilibus, membranaceis tenacellis, extus glabris dilute-fuscescentibus, dein discum albo-pallescentem marginibus inflexis cinctum recludentibus; ascis clavatis, apice rotundatis, 45-50-6-8, 8-sporis; sporidiis oblongo-clavatis, obtusis, rectiusculis, eguttulatis, hyalinis, 6-9-2, distichis; paraphysibus filiformi-clavulatis, apice brunneolis, 5-6 μ . cr.

Hab. ad ramulos quercinos juniores languidos in Thuringia et Franconia.

18. **Cenangium turgidum** Fr. Syst. Myc. II, p. 236, *Peziza quernea* 2307
Schw. Carol. n. 1265. — Aggregatum, obovatum, subrugosum, nigro-fuscescens, ore connivente stellatim infexo, subiculo corticali tumido; sporidiis distichis, fusoideis, rectis, 4-guttulatis, hyalinis.

Hab. ingenti copia ad ramos et ramulos querneos eorumque excrescentias occupat in Carolina Amer. bor. — Ascomata furfuracea, primo clausa dein aperta, 4-6 mm. lat. *C. fuliginoso* utique affine, sed tubercula, cui insident cupulæ, multo minora magis prominent, subhemisphærica, intus tota lignosa (nec fungosa) et corticis colore; cupulæ minores minus prominent. *Dichæna* facile dicendum. Fr. l. c.

19. **Cenangium salignum** Fuck. Symb. p. 268, nec Auct. — Cupulis ut 2308 in *Tymp. saligna* (quam cfr.); ascis anguste clavatis, 8-sporis, 36 \approx 4; sporidiis cylindraceis, curvatis, continuis, 6 \approx 1,5.

Hab. in ramis siccis Salicis capræ prope Estrich Rhenogoviæ et in Belgio (LAMBOTTE).

20. **Cenangium spermatiosporum** (Nyl.) Sacc. *Tympinis* Nyl. Obs. 2309 p. 70, Karst. M. T. I, p. 227. — Erumpens, solitarium v. cæspitosum, nigricans, mox apertum, concavum vel planiusculum, margine subinæquali, nudum, 0,6-0,9 mm. lat.; ascis elongato-clavatis, 11 μ . cr., octosporis; sporidiis breve ellipsoideis, continuis, 6-7 \approx 4; paraphysibus filiformibus, apice incrassatis, fuscis, conglutinatis.

Hab. in cortice Populi tremulæ emortuo in Fennia. — Adsunt spermogonia cæspitosa, nigra, minuta, sporulis allontoideis 3-4 \approx 0,5. An status octosporus *Tymp. populinæ*?

21. **Cenangium subnitidum** Cooke et Phill. in Grev. III, p. 186. — 2310 Cupulis gregariis erumpentibus turbinatis, demum patellatis, bruno-nigricantibus, disco marginato-depresso dein plano vel convexo, extus nudi sopacis vel nitidulis; ascis clavato-cylindraceis, 8-sporis; sporidiis anguste ellipsoideis, curvis, biguttatis, pallide succineis, 15 \approx 5.

Hab. in ramis emortuis deiectis Alni, Trefriw North Wales Forres Britanniæ. — Adsunt pycnidia conica vel subsphærica sæpe solitaria; stylosporis anguste fusoideis, curvis, hyalinis, 10-15 \approx 1.

22. **Cenangium rubiginellum** Sacc. *C. rubiginosum* Cooke in Rav. 2311 Fungi Amer. n. 635, nec Fr. — Sparsum v. gregarium, rubiginosum; cupulis erumpentibus, coriaceis, breviter stipitatis, glabris, disco pallidiore; ascis cylindraceis; sporidiis ellipticis vel utrinque leniter acuminatis, hyalinis, nucleatis, 16 \approx 8.

Hab. in ramis Ostryæ virginicæ in Carolina Amer. bor. (RAVENEL).

23. **Cenangium Abietis** (Pers.) Rehm Disc. pag. 227, *Peziza Abietis* 2312

Pers. Syn. p. 671, *P. cervina* Pers. Syn. p. 647 (non *Cyphella digitalis*), *Cen. ferruginosum* Fries Syst. Myc. II, p. 187, *Trybl. pineum* Pers. Myc. Eur. p. 332. — Gregarium, coriaceo-membranaceum, subsessile, rugosum, subpruinosum, cervino-nigrescens, ore compresso inflexo, humido patulo, disco lutescente; ascis clavatis, supra rotundatis, crasse stipitatis, 60-80 \approx 10-12, 8-sporis; sporidiis ellipsoideis vel ovato-ellipsoideis, 11-13 \approx 5-7, distichis, continuis, 1-2-guttatis; paraphysibus filiformi-clavatis, virescenti-flavidis.

Hab. ad ramos exsiccatos Pinis silvestris, P. Strobi, Abietis balsamiferæ, A. pectinatæ etc. in Suecia, Germania, Belgio, Gallia, Italia, Britannia, Amer. bor. in Quercu Carolinae (SCHWEIN.). Status secundarii noti. — Species memorabilis, versiformis. Prima ætate simillima *Sphaerice* cuidam cæspitosæ peritheciis subrotundis reniformibusque, pulvere ferrugioso tectis; dein magis explicata, 2-4 mm. lata, subnuda, stipite brevissimo et affinium more in cortice latente, sicca compressa connivente-clausa, sed saepe etiam angulata; humida aperta subrotunda, in lacinias plures divisa, disco olivaceo-lutescente. Color exterior opacus, varius. — β . **strobilinum** A. et S. p. 362: sparsum e badio nigricans. In strobilis Abietis in Gallia et Germania.

24. **Cenangium laricinum** (Pass.) Sacc. *Tympanis laricina* Pass. Erb. 2313 critt. ital. II. ser. n. 1284, vix *Cenangium laricinum* Fuck. — Cupulis erumpentibus, cæspitosis nec raro solitariis, primo clausis, dein apertis, disco concavo vel plano marginato, dissectis intus albis; ascis elongato-clavatis attenuato-stipitatis, ad apicem rotundatis, longitudine et crassitie variis sed ut plurimum 65 \approx 10 octosporis, jodo immutatis; sporidiis mono- vel distichis elliptico-navicularibus integris ad polos nucleatis, hyalinis, 10 \approx 5; paraphysibus obscuris.

Hab. in ramis siccis dejectis Laricis europææ in silvis Tressero Ital. bor. (ANZI). — Spermatia (vel stylosporæ?) subfusiformia vehementer arcuata, hyalina, medio spurie septata, 15 \approx 2,5.

25. **Cenangium acicolum** (Fuck.) Rehm Disc. p. 228. — Gregarium, 2314 erumpens, turbinatum, brevissime crasse stipitatum extus flavo-brunneum, glabrum siccum furfuraceum et convolutum 1-3 mm. lat., ceraceo-coriaceum, disco planiusculo concolore pallidius marginato; ascis clavatis 75-90 \approx 10-12, 8-sporis; sporidiis ellipsoideis rectis continuis hyalinis, 12-14 \approx 3,5-4,5 distichis; paraphysibus filiformibus, sursum clavulatis flavo-brunneolis.

- 562 Discomyceteæ, Dermateæ, Hyalosporæ, Cenangium.
Hab. in acubus putridis *Pini silvestris* in Germania.
26. **Cenangium farinaceum** (Pers.) Rehm Disc. p. 226, *Peziza* Pers. 2315
Syn. p. 672, *Pez. pinicola* β. *cæspitosa* Fr. Syst. Myc. II, p. 113,
Pseudophacidium decorticans Rehm in Myc. March. n. 854. —
Cæspitosum, innato-erumpens, seriatum, ceraceo-coriaceum, glo-
bosum clausum dein subapplanatum, apertum, subsinuosum, brun-
neo-nigricans, griseo-pulvereum, disco nigro, 1-1,5 mm. lat.; ascis
clavatis, apice rotundatis, 75-90 ± 10-12, 8-sporis; sporidiis sub-
clavatis rectis curvulisve, continuis, 2-guttatis, 15-17 ± 5-6, disti-
chis; paraphysibus filiformibus 2 μ. cr., subcyaneis.
Hab. in ramis siccis *Pini silvestris* in Germania.
27. **Cenangium leptospermum** B. et C. North Amer. Fungi n. 768. — 2316
Fasciculatum, minutum, nitidum, subglobosum, disco punctiformi;
sporidiis elongato-fusiformibus, arcuatis, pluri-nucleatis, nucleis
globosis.
Hab. in ramis *Abietis Douglasii*, Alabama Americæ borea-
lis (PETERS).
28. **Cenangium helotoides** Mout. et Sacc. (sp. nova). — Ascomatibus 2317
sparsis obconicis breve stipitatis, totis olivaceo-nigris, $\frac{1}{3}$ mm. lat.;
disco concavo, margine diu inflexo subtecto; ascis clavatis, 8-sporis,
120-130 ± 20; sporidiis distichis ellipsoideo-cymbiformibus, conti-
nuis, hyalinis, granulosis, 18-22 ± 8-8,5; paraphysibus filiformibus
ascos paullo superantibus superne atro-olivaceis.
Hab. in pagina sup. foliorum adhuc pendulorum *Juniperi*
communis pr. Liege Belgii.
29. **Cenangium acuum** C. et Peck in Grev. VII, p. 40. — Ita, teste 2318
Cooke, reformanda *Mollisia Pinastri* (C. et P.) Sacc., supra de-
scripta p. 327.
30. **Cenangium Sabalidis** (Ell. et Mart.) Sacc. *Dermatea* Ell. et Mart. 2319
Amer. Nat. 1884, p. 1147. — Ascomatibus tenuibus, gregariis vel
sparsis, brunneis, 250-450 μ. diam.; ascis subsessilibus, 35 ± 5;
sporidiis bi-tristichis, oblongis vel clavato-oblongis, lutescentibus,
2-nucleatis, 4-6 ± 1-1,2.
Hab. in petiolis emortuis *Sabal serrulatae*, Cove springs
Americæ borealis.
31. **Cenangium pelidnum** (K. et C.) Sacc. *Dermatea pelidna* K. et 2320
C. Grev. IX, pag. 25. — Cæspitosum, subcoriaceum; cupulis (pres-
sione mutua) irregularibus, lète badiis, furfuraceis 3 mm. diam.,
disco atro-brunneo vel fuligineo; ascis minimiis, cylindraceis, 30 μ.
longis; sporidiis linearibus 4 μ. longis.

Discomyceteæ, Dermateæ, Hyalosporæ, Cenangium. 563

Hab. in ramis decorticatis, Roschberg Afr. austr. (MAC-OWAN).

— Affine *C. furfuraceo* et *C. populneo*.

32. **Cenangium pezizoides** Peck 31 Rep. Stat. Mus. p. 48. — Cupulis 2321 sparsis, minutis erumpentibus, sessilibus v. basi angusta affixis, levibus nigris; ascis oblongo-clavatis; sporidiis stipatis, oblongo-ellipsoideis 20-25 μ . long. sæpe grosse 1-nucleatis, subinde curvulis.

Hab. in petiolis foliorum Cassandræ calyculatæ, socio *Cen. Cassandræ*, Center Amer. bor. — A *C. Cassandræ* mox distinctum.

33. **Cenangium Cassandræ** Peck 31 Rep. Stat. Mus. p. 48. — Cupulis 2322 oblongis v. hysteriformibus, erumpentibus, epidermide rupta, arcte cinctis; ascis oblongo clavatis; sporidiis linearibus curvis muco obvolutis, leniter coloratis 28-30 μ . long.

Hab. in petiolis foliorum Cassandræ calyculatæ, Center Amer. bor.

34. **Cenangium lichenoideum** B. et Br. Fung. of Brisb. p. 404, t. 45, 2323 f. 9. — Cæspitosum, cinereum; cupulis turbinatis, stipitatis, verrucis cinereis irregularibus vestitis, perfectis, margine crasso, incurvo, cinereo, striato, junioribus absque ordine squamosis; hymenio levi rufo-brunneo; ascis octosporis; sporidiis ellipsoideis, 32-36 μ . long.

Hab. ad truncos Brisbane Australiæ. — Ascomata 1,2-2,2 mm. lata.

35. **Cenangium brasiliense** Mont. Cent. VII. n. 95. — Gregarium, 2324 erumpens; excipulo coriaceo initio cum stipite obpiriformi clauso pallido, hirto, dein cupulari-aperto immo toto explanato, disco cinnamomeo nudo; ascis longe clavatis inter paraphyses capillares nidualantibus, octosporis; sporidiis fusiformibus.

Hab. in cortice arborum provinciæ Goyaz in Brasilia (WEDDEL).

36. **Cenangium fulvo-tingens** B. et C. North Am. Fungi n. 764. — 2325 Fasciculatum, fulvo-tingens; cupulis tenuibus extus venosis subtiter pulverulentis, congestis, compressis; sporidiis oblongis curvulis, suballantoideis, 6-7 μ . long.

Hab. in ligno sicco, quod colore fulvo maculatur, in Pensylvania Amer. bor. (MICHERER).

37. **Cenangium Aparines** (Wallr.) Fuck. Symb. pag. 271, *Tympanis* 2326 *Aparines* Wallr. Flor. Crypt. n. 2342. — Ascomatibus sparsis, primum hypophlœodeis, dein epidermidem scissam infarcentibus, sessilibus, subrotundis, extus fuligine tenui adspersis asperulis nigricantibus diu clausis, subglobosis, denique in discum dilute cine-

reum carnosulum limbo infexo cinctum reclusis; ascis clavatis, 8-sporis; sporidiis fusoideis 4-5-guttulatis, hyalinis, continuis.

Hab. ad caules sicclos Galii Aparines in agro Badensi (KNEIFF), Thuringiaci (WALLR.) et Rhenogovia (FUCK.).

38. **Cenangium Pustula** Rehm Disc. pag. 225. — Gregarium, innato-²³²⁷ erumpens, globosum dein scutellatum extus brunneolum, siccum griseo-albo-furfuraceum, striatum et convolutum 1-3 mm. lat. membranaceo-ceraceum; disco planiusculo griseo-brunneo; ascis clavatis supra rotundatis, 36-45 \approx 6-8, octosporis; sporidiis distichis elliptico-oblongis obtusis rectis curvulisve, continuis, hyalinis, 6-8 \approx 2,5-3; paraphysibus filiformibus 3 μ . cras. ascos superantibus.

Hab. in foliis siccis Eryngii campestris et Convallariæ majalis in Thuringia. — An diversum a sequente?

39. **Cenangium Umbellatarum** Ces. in Rabenh. H. Viv. II ed. u. [312]²³²⁸ et in Bot. Zeit. 1857, p. 93. — Primitus stictidiforme, immersum, pulverulentum, mox reclusum, limbo dentato v. lacero; disco primum luteo v. fuscello, demum fusco v. badio; ascis e tubuloso clavatis tenuibus, dense stipatis, 44 μ . longis; paraphysibus parcis rigidulis filiformibus æquilongis intermixtis; sporidiis minutis, 6-8 μ ., ovatis, obtusis, (spurie?) 1-septatis.

Hab. in caulis Oreoselini et Eryngii prope Vercelli Ital. bor.

40. **Cenangium Colensoi** Berk. Fl. New Zeal. pag. 201. — Sparsum,²³²⁹ primitus oblongum, demum orbiculari-expansum, fuscum, hymenio pallido subcarneo; ascis oblongis; sporidiis minutis, elliptico-subcymbiformibus 6-7 μ . hyalinis.

Hab. in foliis putridis Phormii (ut videtur) in Nova Zelandia (COLENSO).

41. **Cenangium microspermum** Sacc. et Ell. Mich. II, p. 571. — Ascomatibus cæspitoso-erumpentibus, calyculatis, sessilibus, subcoriaceis, vix $1/5$ mm. diam., fusco-cinereis, extus setuloso-olivascentibus; excipulo parenchymatico olivaceo; ascis teretiusculis breve stipatis, 40 \approx 4-5; paraphysibus filiformibus obvallatis, octosporis; sporidiis distichis cylindraceo-clavulatis, 5 \approx 1,5, hyalinis, rectis v. leniter curvis.

Hab. in Valsaria didymospora parasiticum ad Ilicis opacæ ramulos, Jona N. J. (ELLIS). — A *Cenangio pulveraceo* (A. S.) certe diversum.

II. *Encælia* Fr. Ascomata typice majora, tenui-coriacea, demum varie contorta.

42. **Cenangium furfuraceum** (Roth) De Not. Prof. Disc. p. 30, *Peziza* 2331

Roth Cat. II, p. 257, t. IX, f. 3, *Phibalis* Wallr. Crypt. n. 2413, *Encælia* Karst. M. F. I, p. 218, *Pez. Coryli* DC. Fr. VI, p. 21.

— Ascomatibus sessilibus cæspitosis, carnoso-coriaceis, majusculis, saepius difformibus, extus furfure ferrugineo-pallido obtectis, dein discum cinnamomeo-nigrum margine involuto cinctum recludentibus; ascis clavulatis 81-100 = 6-8, paraphysatis, octosporis; sporidiis cylindraceis curvis, 2-guttulatis (quasi Allantosporarum) 10-12 = 3-4, hyalinis.

Hab. ad ramos coryleo et alneos in Germania, Helvetia, Gallia, Belgio, Fennia, Britannia, Rossia. — Ascoma 5-15 mm. lat.

43. **Cenangium Fuckelii** Sacc. Miscell. myc. I, p. 8, *Dermat. furfuraceo* 2332

Fuck. Symb. p. 278 et Sacc. F. Ven. IV, p. 29, nec Auct.

— A *C. furfuraceo* genuino differt ascomatibus multo minoribus, vix 1-1½ mm. diam.; ascis cylindraceis; sporidiis ellipsoideo-oblongis, rectis, 10-11 = 6-8; structura tamen ascomatis et indumenti simillima.

Hab. in ramis *Coryli* *Avellanæ* in Rhenogovia (FUCKEL) et in Pruno in Ital. bor. — Icon in *Fung. ital.* n. 1312 pessime fucata, nam non ochracea sed stramineo-furfuracea prætereaque ascomata nimis aperta.

44. **Cenangium populneum** (Pers.) Rehm Disc. p. 220, *C. populinum* 2333

Tul. Carp. III, p. 171, Sacc. F. it. t. 1310 (f. *Populorum*), *Peziza fascicularis* Alb. et Schw. Consp. Fung. nisk. p. 315, t. 12, f. 2,

Fr. Syst. Myc. II, p. 75, *Peziza populnea* Pers. Syn. Fung. p. 671, *Dermatea fascicularis* Fr. Summ. Veg. Scand. p. 362, *Encælia*

Karst. M. F. I, p. 217, *Peziza crispa* Sow. Engl. Fungi t. 425.

— Cupulis cæspitose aggregatis, raro solitariis, cortici interiori adnatis, sed epidermide secedente superficialibus, sessilibus, coriaceo-membranaceis, difformibus, subfarinaceis, fusco-nigrescentibus, rugosis, disco concavo, obscure spadiceo, dein pallidiore, 5-15 mm. lat.; ascis cylindraceo-clavatis, 70-92 = 8-9; sporidiis distichis, oblongatis vel elongatis, ut plurimum curvulis, eguttulatis, 11-16 = 3-4; paraphysisibus sursum incrassatis.

Hab. ad truncos et ramos putres Populi tremulæ, rarius Sorbi, Salicis, Fraxini, Corni in Suecia, Lapponia, Fennia,

Gallia, Germania, Hungaria, Belgio, Italia, Sibiria, Britannia, America boreali.

45. **Cenangium Ellisii** Sacc., *Cenang. purpureum* Ellis Bull. Torr. 2334 Cl. 1876, pag. 108, nec Fr. — Cæspitosum erumpens, irregulare, saturate atro-purpureum, sordide albescente furfuraceum; ascis clavato-cylindraceis; paraphysibus sursum leniter incrassatis; sporidiis phaseoliformibus, hyalinis, obtusis, 20 ± 8 .

Hab. in Viburno emortuo, Newfield, New Jersey Amer. bor.

— *C. furfuraceo* simile sed minus. Verisimiliter var. *C. populnei*.

46. **Cenangium tiliaceum** (Fr.) Karst. Rev. p. 145, *Peziza* Fr. Syst. 2335 Myc. II, p. 76, *Dermatea* Fr. Summa pag. 362, *Encælia tiliacea* (Fr.) Karst. M. F. I, p. 218, Bresad. Trid. p. 93, t. 105. — Subcæspitosum, sessile, ceraceum, cinnamomeum, subtus rugosum nudum; ascis tereti-clavatis, longe stipitatis, $110-130 \pm 7-8$, octosporis; sporidiis distichis cylindraceis, obtusis, leniter curvis, pluri-guttulatis, hyalinis, $12-18 \pm 3-4$; paraphysibus filiformibus apice clavulatis, aureo-granulosis.

Hab. in ramis exsiccatis Tilæ, rarius Mali in Suecia, Fennia, Gallia, Germania, Italia. — Erumpens, cæspitosum, sed et solitarium, 6-12 mm. lat. plane sessile, extus corticatum quasi, ceterum nudum glabrum dilutius, disco opaco fusco-cinnamomeo, in solitariis plano, margine integerrimo, sed cupulæ in cæspitosis, ob mutuam pressionem concavæ, undulato-repandæ.

47. **Cenangium Ulmi** Tul. Carp. III, p. 170, t. XIX, f. 18-22, *Dermatea* Fuck. Symb. App. II, p. 56. — Gregarium, ceraceo-coriacum initio globulosum clausum, magnitudine seminis sinapeos, rubiginosum, innato-erumpens, e basi communi cupulis 2-4 orientibus, denique scutellatum, tortuosum, inæqualiter marginatum, disco flavo v. brunneo, 2-4 mm. lat.; ascis cylindraceis, 8-sporis, apice rotundatis, $6,5 \mu$. cr.; sporidiis cylindraceis, curvulis, continuis, $10-15 \pm 2,5-3$, saepe 2-guttulatis; paraphysibus filiformibus, clavulatis.

Hab. in ramis corticatis siccis Ulmi in Gallia et Rhenogovia.
— Sec. Fuck. asci $108 \pm 10-11$, sporidia $14-16 \pm 4$. Spermogonia adsunt, teste Tul.

48. **Cenangium Carpini** Rehm Disc. p. 221. — Atro-brunneum, coriaceum extus striatum, siccum flavidobrunneum, erumpens, primo globosum in cæspitulos subrotundos ex 8-12 cupulis constantes depositum, epidermide laciniata cinctum, dein inæquale patellatum; disco plano brunneo, 2-4 mm. lat.; ascis tereti-clavatis, apice rotundatis, $75-80 \pm 8-9$, octosporis; sporidiis elongatis obtusis cur-

vulis continuis, hyalinis, 12-15 = 3-3,5, distichis; paraphysibus filiformibus apice clavulatis 4 μ . crass. brunneolis.

Hab. in ramis siccis *Carpini Betuli* in Carniola (Voss). —

A *Cen. populneo* differt cupulis multo minoribus, a *Cen. Ulmi* colore flavidо-brunneo.

49. **Cenangium impudicellum** Karst. Rev. p. 146, *Encælia impudicella* Karst. M. F. I, p. 219. — Solitarium vel cæspitosum, erumpens, ex sphæroideo-hemisphæricum, tandem explanatum, difforme, leniter furfuraceum, demum subnudum, primitus griseum, dein subcinnamoineo-fuscum, marginem versus dilutius, siccum fuligineo-fuscum vel nigricans, breviter stipitatum vel sessile, coriaceum, disco ex lutescente pallido, siccitate subcinnamomeo, levi vel ruguloso, margine integerrimo, 3-5 mm. lat.; ascis cylindraceo-subclavatis, 40-50 = 5, apice jodo non cœrulecentibus; sporidiis distichis vel oblique monostichis, elongatis, utroque apice paullulum attenuatis, simplicibus, curvulis, 6-9 = 1,5-2; paraphysibus haud bene discretis.

Hab. ad corticem ramorum Abietis excelsæ dejectorum exsiccatorum in Mustiala et prope oppidum Wasa Fenniæ. — Est forsitan *Peziza tristis* Saut. in Flora, 1841, p. 309, teste Karst.

50. **Cenangium inculcatum** (Karst.) Sacc. *Encælia inculcata* Karst. M. F. p. 220. — Solitarium, sparsum vel subgregarium, immersum, dein plus minus emergens, subsphæroideum vel dein depresso-sphæroideum, primitus clausum, dein vertice in laciniæ irregulares complures acutas tenerrimas dehiscens, ore constricto, atrum, disco pallido, demum nudum, circiter 0,7 mm. lat., coriaceo-membranaceum; ascis et paraphysibus cohærentibus; sporidiis elongatis vel subfusoideo-elongatis, simplicibus vel spuriæ tenuiter uniseptatis, rectis vel quadantenus incurvatis, incoloribus, 9-12 = 2,5-3,5; hymenio jodo intense cœrulecente.

Hab. in cortice Abietis excelsæ, Mustiala Fenniæ. — Nuper ab Auctore (Rev. p. 146) ad *Odontotrema* relatum.

51. **Cenangium russum** (B. et Br.) Sacc. *Peziza russa* B. et Br. Fung. of Ceyl. n. 940. — Cupulis subcoriaceis rugulosis granulatis, vivide rubro-brunneis, margine appendiculis moniliformibus ornato; ascis 50 μ . longis; sporidiis allantoideis 7-8 μ . longis.

Hab. ad cortices, Central province, Ceylon.

52. **Cenangium cumulatum** (Quél.) Sacc. *Encælia cumulata* Quél. X, Suppl. p. 14, tab. IX, fig. 14. — Ascomatibus turbinatis, compressis, dein varie contortis, 400 μ . diam., subtiliter tomentosis, pallide lute-

- 568 Discomyceteæ, Dermateæ, Hyalosposæ, Cenangium.
scentibus, ochraceis; hymenio plano vel irregulari, pulverulento,
concolori; sporidiis fusoideis, 3-4 μ . long. curvatis.
Hab. seriatim in rimis corticis Aceris Pseudoplatani, Jura.
53. **Cenangium striatum** Mout. et Sacc. (sp. nov.) — Ascomatibus 2342
dense, late gregariis, obconico-cupulatis, tabacino-fuscis, disco con-
cavo, 2-2,5 mm. lat., longitud. striatulis; ascis cylindraceo-oblon-
gis, 160 \approx 16; sporidiis ovato-oblongis, 26-30 \approx 9-10.
Hab. in ramis exsiccatis Betulæ albæ, pr. Liege Belgii. —
Ascomata initio cortice relaxato velata.
54. **Cenangium Bloxami** (Phill.) Sacc. *Encælia* Phill. Disc. pag. 338. 2343
— Gregarium, sæpe cæspitosum, breve stipitatum, cyathiforme, co-
riaceum, brunneo-nigricans, minute verrucosum; disco luride brun-
neo, margine subundulato, carne atro-purpureo-brunnea; ascis cla-
vatis, octosporis; sporidiis ellipsoideis, 3-4 \approx 2; paraphysibus fili-
formibus concretis.
Hab. in ligno putri in Britannia (BLOXAM). — Ascoma 2-8 mm.
lat. — Ex hymenio sæpe exsiliunt stylosporæ fusoideæ, uniseptatæ,
basidiis filiformibus suffultæ.
55. **Cenangium Reichenbachii** (Rabenh.) Rehm Disc. p. 229, *Peziza* 2344
Reichenbachii Rabenh. D. cr. Fl. p. 367, n. 2831. — Cæspitosum,
erumpens, 5-8 mm. alta, 5-15 mm. lat., primo rotundatum, clau-
sum, demum scutellatum e coriaceo carnoso-molle, extus farinosum,
albido-griseum, disco plano, æquali v. tortuoso, atro-brunneo, mar-
gine tenui plerumque gibboso; ascis cylindraceo-clavatis, octospo-
ris, flavescentibus; paraphysibus numerosis, filiformibus; sporidiis
ovoideis, nubilosus.
Hab. in ramis emortuis muscosis ad «Schandau» in Helvetia,
Saxonia. — An, teste Rehm, stirps bulgariacea?
56. **Cenangium ustale** (B. et C.) Sacc. *Peziza ustalis* B. et C. North 2345
Am. Fungi n. 718. — Congestum, irregulare, extus rufum, subti-
liter tomentosum; hymenio spadiceo; stipite cylindrico brevi.
Hab. ad ramos dejectos, Alabama Amer. bor. (PETERS).
57. **Cenangium Ravenelii** (B. et C.) Sacc. *Peziza Ravenclii* B. et C. 2346
North Am. Fung. n. 717. — Parvum, subsessile, concavum, flexuo-
sum, extus albidum, subtiliter tomentosum, intus rufescens; stipite
brevissimo.
Hab. in Patellaria nigro-cinnabarina in Carolina sup.
Amer. bor. — Ascoma 2 mm. lat.
58. **Cenangium purpureum** (Hedw.) Sacc. *Octospora purpurea* Hedw. 2347
Musc. fr. II, p. 63, t. 22, *Peziza caiycina* Nees Syst. f. 286,

Peziza purpurea Fr. S. M. II, pag. 77. — Subcoriaceum, infundibuliforme, extus pruinatum, obsolete purpureum; disco sanguineo; stipite brevi; ascis (ex icono Hedw.!) tereti-clavatis; sporidiis ellipsoideo-oblongis, hyalinis.

Hab. supra basin residuam truncorum *Quercus* putrilaginosam Lipsiæ, Upsaliæ, Ostrogothiæ, Sueciæ. — Stipes vix attinens, leviter striatus, infima basi nigricans, sursum in patellam aper-tam ampliatus. Extus fuscescens, tenerrime albido-papillosum, unde senescens, præcipue versus marginem, pruinatum; substantia albida, solidiuscula.

59. **Cenangium nitidum** (Saut.) Sacc. *Peziza nitida* Saut. Pilz. Salzb. 2348 p. 8. — Carnosum, stipite brevi crasso; ascomate extus roseo, subgloboso, disco coccineo, margine tenui inflexo.

Hab. ad ramos emortuos, prope Mittersill Salzburg. — Asco-ma 4–10 mm. diam. et pariter longum. *C. purpureo* affine.

60. **Cenangium tegulare** (Weinm.) Sacc. *Peziza Weinm.* Syllog. p. 112. 2349 — Sessile, carnosò-coriaceum, brunneum, demum explanatum, mar-gine repando, extus pruinatum.

Hab. in virgultis putrescentibus locis umbrosis silvaticis Petro-poli Rossiæ (WEINMANN). — Species luculenter distincta; ex habitu et loco inter *Encælia*s recensenda; solitaria aut subcæspitosa, tenax, sicca saltim subcoriacea; junior subglobosa, mox campanu-lata, demum explanata, 1,3–2 cm. et ultra lata, dilute brunnea aut olivascens, sicca obscure brunnea. Margo patens, repandus, non quidem lacero-fissus, sed sub lente crenulatus.

61. **Cenangium triste** (Saut.) Sacc. *Peziza tristis* Saut. Pilz. Salzb. 2350 p. 8. — Sessile, solitarium, coriaceum, nigrum, margine involuto integro.

Hab. in truncis Pin i putrescentibus prope Mittersill. — Affine *C. purpureo*. Ascoma 2–10 mm. lat.

62. **Cenangium rubiginosum** (Fr.) Sacc. *Pez. rubiginosa* Fr. Elench. 2351 II, p. 7. — Subcarnosum, persistens, turbinato-sessile, extus pur-pureo-pruinatum, margine tumido integro, disco urceolato nigro.

Hab. ad ligna putrida, Petropoli Rossiæ (WEINMANN). — Disci colore præcipue a *C. purpureo* recedit. Elevato-sessile, basi haud applanatum, 2–5 mm. altum et latum, regulare, margine subin-fexo integro. Sub disco nigro conspicitur stratum ascorum am-plorum albicans.

63. **Cenangium fissum** (Wallr.) Rehm Disc. p. 222, *Peziza fissa* Fr. 2352 Syst. II, 75, *P. fascicularis* β. *corylina* Wahlenb. Suec. II, 993,

570 Discomyceteæ, Dermateæ, Hyalosporæ, Cenangium.

P. lacera Pers. Myc. I, 227, *Phibalis fissa* Wallr. Crypt. n. 4683.

— Ascomatibus hypophlœodeis, mox epidermidem lacero-scissam infarcentibus subeuntibusque, subcæspitosis, sessilibus, membranaceis fragilibus, primum clausis, extus scobe farinosa dilute cinnamomea s. cervina adspersis, dein hiantibus discumque album recludentibus.

Hab. ad ramos coryleo in Suecia et Germania. — An diversum a *C. Coryli* Corda?

64. **Cenangium helvolum** Jungh. Crypt. Java p. 30 (*Peziza, Encælia*). ²³⁵

— Ochraceo-gilvum, cupulatum, stipitatum, extus granuloso-farinaceum, margine granuloso-crenato, 4 mm. latum, cupulatum, intus leve fusco-gilvum, extus strato farinaceo crassiusculo obductum, pallide gilvum, basi in stipitem brevem concolorem attenuatum.

Hab. in culmis Bambusæ arundinaceæ gregatim juxta articulos, in silvis Montis Merapi Javæ. — In statu siccato cupula (marginibus conniventibus) clausa, rima longitudinali aperta.

III. *Phæangium* Sacc. Sporidia fusca.

65. **Cenangium phæosporum** Cooke Grev. XII, pag. 44, Phill. Disc. ²³⁵

p. 346. — Cupulis sparsis erumpentibus, rufo-brunneis, breve stipitatis, extus delicate furfuraceis, in stipitem obscuriorem attenuatis, margine subincurvato, disco subconcolore; ascis elongato-clavatis, octosporis; sporidiis monostichis pallide brunneis, late ellipsoideis, levibus, 12-8; paraphysibus crassis, apice clavatis brunneis.

Hab. in cortice Aceris, Exeter Britanniæ (E. PARFITT). — Ascoma 1 mm. lat.

66. **Cenangium tetrasporum** (Ellis) Sacc. *Dermatea tetraspora* Ellis ²³⁵

Bull. Torr. Cl. 1876, p. 108. — Erumpens, simplex vel subcæspitosum, subsessile, marginatum, 2,5 mm. lat., disco plano vel vix concavo, obscure brunneo, rugoso, margine tenui et pallido; ascis brevibus, clavato-cylindraceis, latis, tetrasporis; sporidiis ovato-oblongis, nonnullis sphæroideis, brunneis, 20-25 = 15-20, crasse unicellularis, intus primo granulosis.

Hab. in ramulis nondum cæsis Quercus coccineæ, Newfield, New Jersey Amer. bor.

67. **Cenangium punctoideum** (Cooke) Sacc. *Tympanis punctoidea* ²³⁵

Cooke Disc. U. S. II, p. 30. — Minutum, erumpens, punctiforme atrum, coriaceum, sæpe seriatum; ascomatibus solitariis vel 2-3 acervatis, compressis, irregularibus, breviter stipitatis; disco con-

Discomyceteæ, Dermateæ, Hyalosporæ, Cenangium.

571

cavo, in sicco lateraliter compresso; ascis subcylindraceis; sporidiis ellipsoideis, brunneis.

Hab. in corticibus, United States Amer. bcr.

38. **Cenangium lignicolum** Preuss Hoyersw. n. 341. — Gregarium, 2357 nudum, subcorneum, nigrum; cupulis primo subglobosis, conniventibus, dein planis; stipite crasso, brevissimo, dein atro; ascis cylindricis, octosporis, paraphysibus mixtis; sporidiis ovatis, nigris.

Hab. ad truncos arborum frondosarum, Hoyersverda.

IV. Species *Eu-Cenangii*, fructificatione ignota, dubiæ.

39. **Cenangium apertum** Schw. Syn. Am. n. 2001. — Sparsum simplex, cupulis 2-4 mm. lat., primum clausis, pulvinato-rotundis extus nigro-ferruginosis, pulverulentis, demum late apertis, marginibus irregulariter inflexis, saepe ruptis; disco pallido.

Hab. in ramulis emortuis Hydrangeæ, Bethlehem et Salem Amer. bor. (SCHWEINITZ et MICHERER). — Affine est *C. ferruginoso*.

70. **Cenangium faticens** Schw. Syn. Am. n. 2005. — Minutum (majus tamen in Syringa) fasciculatum, prorumpens, sessile; cupulis globosis primum clausis, interdum difformibus, pulvere copioso testis, demum omnino fatiscentibus.

Hab. in cortice Syringæ et evolutius in Moro alba, Bethlehem Amer. bor. — Pulvere deterso apparent cupulæ applanatae atro-nitentes, demum irregulariter apertæ.

71. **Cenangium acutum** (Schum.) Fr. S. M. II, p. 188, *Hyster. acutum* 2360 Schum. Saell. p. 151. — Simplex, elliptico-oblongum, naviculare, leve, nigro-fuscum, disco umbrino-fusco.

Hab. in cortice Sambuci nigræ in Saellandia. — Sparsum, vix prominens, ut videtur nudum.

72. **Cenangium Cephalanthi** (Schw.) Fr. S. M. II, p. 188, *Peziza* 2361 Schw. Carol. n. 1256. — Simplex, sessile, compressum, subpulverulentum, nigro-fuscum, margine lacero involuto.

Hab. plerumque transversim erumpens in cortice Cephalanthi occidentalis emortuo in Carolina, Alabama, Greenbush America borealis. — Optima dicitur species, *Colpom. quercino* minor et obscurior; disco in humidis subrotundo, cinerescente. Non ita prominet.

73. **Cenangium laminare** Fries V. A. H. l. c. p. 560, Kunz. Myc. 2362 Heft. 2, p. 58. — Cæspitosum, plicato-lobatum, lobis planiusculis nigricantibus albido-pruinosis, apice cavis subclausis.

Hab. ad ramos dejectos putrescentes Robiniae? in Suecia et

572 Discomyceteæ, Dermateæ, Hyalosporæ, Cenangium.

Germania. — Substantia fere membranacea *C. Abietis*, cum quo etiam superficie convenit pulverulenta et forma compressa. Specimina plura basi concrescunt, inde subramosum v. in plicas quasi divisum, quæ hactenus clausæ modo repertæ sed apice cavæ et circum circa hymenio vestitæ.

74. **Cenangium striatum** (Bert.) Fr. S. M. II, p. 188, *Pez. striata* 2363

Bertier apud Spreng. in Vet. Ac. Handl. 1820, p. 50. — Simplex, sessile, compressum, naviculare, atrum, extus transversim sulcato-striatum.

Hab. in cortice Guilandinæ Bonduccellæ, Guadalupa. — *Hysterio* simile et quoad substantiam reliquis firmius, sed omnibus Cenangi characteribus gaudet. Cupula 4 mm. longa, subnuda, extus striis confertis parallelis exarata; quando plura specimina aggregata passim angulata; margine involuto. Discus concavus, nigrascens.

75. **Cenangium Viburni** (Schw.) Fr. S. M. II, p. 185, *Pez. Viburni* 2364

Schwein. Carol. n. 1264. — Subcæspitosum, stipitatum, rugosum, opacum, atrum, margine crasso rimoso-dehiscente.

Hab. ad cortices Viburni, Jadhin, Carolinæ. — An *Blitrydium?* *C. Pinastri* simile, tamen certe distinctum; stipes 2 mm. usque longus, subinde ramosus, crassus; cupula non patellato-collapsa, initio cinereo-pulverulenta, dein omnino nuda, in lacinias 4-5 subæquales dehiscens, disco intusque etiam atra.

76. **Cenangium viticolum** (Schw.) Fuck. Symb. p. 240, *Peziza Schw.* 2365

Carol. n. 1263, *Tympanis* Fr. Syst. Myc. II, p. 176. — Minutum, sessile, hemisphæricum, rugulosum, opacum, fuscescenti-nigrum, disco marginato; ascis maturis nondum visis (Fuck.).

Hab. ad ramos exsiccatos Vitis in Carolina (Schw.), Gallia (Fr.), Germania (Fuck.). — Species distinctissima, habitu fere *Pleosporæ herbarum*, substantia moliori (tremelloso-coriacea); sparsa, solitaria, innata, disco primo impresso punctiformi obtuse marginato, dein dilatato subelapso.

77. **Cenangium bicolor** (Wallr.) Sacc. *Phibalis bicolor* Wallr. Crypt. 2366

n. 2400. — Ascomatibus hypophlœodeis, mox epidermidem scissam penetrantibus, e strato pseudostromatico carnoso convexulo cæspitose erumpentibus, extus tegmine furfuraceo stipato albo-cinereo vestitis, primum subrotundis basi subæqualibus clavis sphæriæ-formibus, dein in basin productis subturbinatis vertice in osculum circinatum, demum in discum planum fusco-atrum margine tenui cinctum reclusis, scutellaribus.

Hab. ad Rhamni Frangulæ ramulos sicclos in Thuringia.

-- Ascomata 2-6 in cæspite stratoque coccineo epidermidis juncta.

78. **Cenangium Staphyleæ** Schw. Syn. Am. bor. n. 1989. — Sparsum 2367 vel gregarium, primum epidermide tectum minutum, sphæriæforme, demum prorumpens, nigrum, apice aperto irregulari; disco indistincte pulverulento.

Hab. passim in junioribus ramis Staphyleæ, Bethlehem Americae borealis.

79. **Cenangium palmatum** Schw. Syn. Am. bor. n. 1986. — Cæspitum 2368; cæspitibus pugilliformibus declinatis, cupulis nigris subcylindricis, quasi ex tuberculo cæspitosim orientibus variis sibi ipsis impositis, apice primum in pseudostiolum abeuntibus, demum aperiensculis, et quasi poro hemisphærico concavo minuto excavatis, intus pulveraceis, 2-4 mm. altitudine.

Hab. tractu longitudinali prorumpens ex cortice Rhododendri maximi, Bethlehem Amer. bor.

80. **Cenangium arcticum** (Ehrenb.) Fr. S. M. II, p. 186, *Trybl. arcticum* Ehrenb. Fung. Cham. n. XXXIII, t. 20, f. 17. — Solarium, minimum, nigrum, suborbiculare, cupula in laciniis tres obtusas dehiscente, demum trigonum, disco albido.

Hab. in caulis Andromedæ tetragonæ, non solum ad sinum S. Laurentii, sed et in Lapponia etc. — Primo clausum, oblongum, glabrum, dein suborbiculare et subhemisphæricum, subpedicellatum, granulosum. Maturum tribus laciniis obtusis dehiscendo formam trigonam acquirit; discus griseo-albidus.

81. **Cenangium Rhois** Schw. Syn. Am. bor. n. 1982. — Cæspitosum, 2370 albido-pulverulentum, sphæriæforme, demum apertum; cæspitulis minutis, fere semper circumdati pulvere ex cortice orto.

Hab. non infrequens in Rhois glabræ ramis, Salem et Bethlehem. — Affine *C. Aucupariæ* et cum hoc confusum.

82. **Cenangium clavatum** (Schw.) Fr. S. M. II, p. 179, *Pez. Ceratosorum* Schwein. Carol. n. 1176. — Cæspitosum, subnudum, fusco-nigricans; cupula subclavata connivente, disco nigrescente.

Hab. in ramis emortuis Pruni serotinæ in Carolina, rarissime. — Cæspites e paucis individuis componuntur, sed cupulæ subsessiles, reliquorum majores et magis evolutæ. Caro flava, in siccis vero fusca.

83. **Cenangium Cratægi** Schw. Syn. Amer. n. 1898. — Sparsum, sessile, repando-effusum, subtremelloso-corneum, atrum aut atro-fu-

574 Discomyceteæ, Dermateæ, Hyalosporæ, Cenangium.

scum, primo globosum, demum apertum, nigro-pulveraceum, pulvere compacto; disco semper impresso.

Hab. passim in ramulis Cratægi, Bethlehem Amer. bor.

84. **Cenangium Spirææ** (Schw.) Berk. North Am. Fungi p. 143, *Der. mea Spirææ* Schw. Syn. Amer. bor. n. 1975. — Subsimplex, aut 2-3 in cæspitum aggregatis, cupulis siccis conniventibus madidis apertissimis, extus brunneo-fuscis, furfuraceo-strigosis; disco badio minore, numquam 4 mm. diam. excedente.

Hab. in ramis Spirææ opulifoliæ rarer, Bethlehem et in Carolina super. Amer. bor.

85. **Cenangium Rosæ** Schw. Syn. Am. n. 2002. — Cæspitosum, pro- 2374 rumpens; cupulis dense aggregatis, corneis, nigro-nitentibus, mar- gine tuberoso, disco plano aperto, cupulis difformibus.

Hab. prorumpens rarius ex cortice Rosæ corymbosæ, Beth- lehem Amer. bor.

86. **Cenangium enteroxanthum** Schw. Syn. Am. n. 2014. — Longi- 2375 tudinaliter seriatum aut irregulariter prorumpens, sessile, hyste- riiforme, subrotundum aut ovatum, plerumque clausum aut rima longitudinaliter apertum, extus olivaceo-nigrum rugosum, intus colore luteo-virente, pulverulentum, præsertim sub disco subaperto.

Hab. parvulum sed distinctissimum in ligno Pomorum, Beth- lehem Amer. bor. — Crusta quadam ligno commixto nigra, ex qua prorumpit.

87. **Cenangium pirinum** Schw. Syn. Am. n. 2013. — Cupulis grega- 2376 riis aut sparsis, atro-fuscis, nitentibus, plerumque transversim aut varie sitis, semper basim versum in stipitem contractis, more *Lophii* siccitate conniventibus, sed etiam late hiascentibus, hysteriformibus, marginibus inflexis, discum planum atro-pulverulentum exhibentibus, interdum scutellam Lichenis referens, rotundato- apertum.

Hab. longe lateque obtegens frustula corticis Piri corona- riæ, Nazareth Amer. bor.

88. **Cenangium Juglandis** B. et C. North Am. Fungi n. 769. — Cæ- 2377 spitosum, globosum, pruinatum, apice nigrum, serius omnino denu- datum.

Hab. in ramis Juglandis in montibus Virginiae Amer. bor.

— Asci nondum observati.

89. **Cenangium sphæriæmorphum** Schw. Syn. Am. bor. n. 1991. — 2378 Gregarium, rarius confluens, epidermide innatum aut prorumpens, interdum quasi fasciculatum, plerumque autem forma obovata tur-

binata, fusco-nigrum, rugosum, punctatum, non stipitatum, disco rarius subaperto, intus ferrugineum, 1 mm. diam.

Hab. in ramis emollitis *Quercus albæ*, Bethlehem Amer. bor. — Ramum agmine vasto longe lateque diffuso occupat.

90. **Cenangium confusum** Schw. Syn. Am. bor. n. 1990. — Sparsum 2379 vel confluens, ramos aut juniores truncos pedatiter obtegens sub epidermide crusta quadam effusa fusco-nigra insidens, cupulis pro-rumpentibus, irregularibus, difformibus, sphæriæmorphis, conspurcatis extrorsum sporidiis quasi profluentibus, ceterum sæpe globosis demum apice aperto, ore aut pezizoideo aut sphæroideo aut irregulari.

Hab. frequens in cortice ramorum *Quercus albæ* in terra jacente, Salem et Bethlehem Amer. bor.

91. **Cenangium fibrisedum** Schw. Syn. Am. n. 2000. — Minutissimum, fasciculatum et sparsum, etiam passim confluens, siccum nigerrimum, cupulis varie formatis, sed semper conniventibus, extus papillis nigris tam dense obsitum (fere *Exosporium Tiliæ* referens), madefactum disco pallido-rubo, atro-pulverulento.

Hab. in lignis denudatis siccis, fibris insidens, primum subiectum, Bethlehem præsertim in *Castaneis* sepium.

92. **Cenangium Castaneæ** Schw. Syn. Am. n. 2004. — Simplex, pro-rumpens, epidermide cinctum et pulvere quodam spurio ex cortice insidens, sessile subrepando-undulatum, marginibus tumidulis, centro discoideo irregulariter depresso, rugoso, subpulverulento, demum ruptum et intus grumosum, fuscum, extus nigro-fuscum, 2 mm. diam. non excedens.

Hab. ramulis junioribus *Castaneæ* rarius insidens, Bethlehem.

93. **Cenangium Betulæ** Fr. S. M. II, p. 187. — Cæspitoso-connatum, 2382 sessile, coriaceo-membranaceum, nudum, atrum, disco concavo sub-compresso, atro-sanguineo.

Hab. erumpens e cortice *Betulæ albæ* in Kamtschatka. (WORMSK.) — Cupulæ majusculæ, subrotundæ, siccæ compresso-connientes, humidæ apertæ subrotundæ. *Cenango Abietis* paulo firmius.

94. **Cenangium compressum** Schw. Syn. Am. bor. n. 1987. — Transversim erumpens, cæspitosum, valde elevatum et ramum quasi aculeans, cupulis semper inflato-compressis, apice valde varie formatis, nigris, furfure cinereo adspersis, apice demum apertiusculis, 4 mm. sæpe altis, furfure externo detergibili (tum fungus nigro-pulverulentus evadit).

576 Discomyceteæ, Dermateæ, Hyalosporæ, Cenangium.

Hab. ex ramis Betulæ carpinifoliae (lentæ) propullans, Bethlehem et in New England Amer. bor.

95. **Cenangium molliusculum** Schw. Syn. Am. n. 2008. — Pezizoidem, aggregatum ac sparsum, saepè invicem adpresso et inde angulatum, statu madido molliusculum 2-6 mm. lat., nigro-olivaceum marginatum, sicco statu corneo-ceraceum, sessile, basi tamen contracta, cupulam marginatam subrepandam lobatam referens, extus olivaceum, disco convexo rugoso, punctato, nigro; intus substantia eleganter flavo-ferruginea.

Hab. eximia species in *Betula carpinifolia*, Mauch Chunk Amer. bor.

96. **Cenangium populinum** Schw. Syn. Am. n. 2003. — Sparsum, 2385 2-4 mm. latum, elevatum cum epidermide, a qua cinctum, primum clausum, nigrum, demum apertum, margine effuso, disco cinereo pulverulento.

Hab. sparsim erumpens ex *Populi italicæ* cortice, Bethlehem Amer. bor. — Clausum fuscescit; faciliter elabitur.

97. **Cenangium Schweinitzii** Sacc. *Dermea populea* Schw. Syn. Am. 2386 bor. n. 1972, nec *Cen. popul.* (Pers.). — Sessile, cæspitosum, coriaceo-membranaceum, minus, ovatum aut oblongum, fere semper clausum, extus cinereo-furfuraceum, intus nigrum; singulis cupulis, sphæriæformibus clavis, rarissime apertis, 1 mm. alt.

Hab. raro in *Populo italicæ*, Bethlehem Amer. bor.

98. **Cenangium fallax** B. et R. (nec Fr.). — Subglobosum, sparsum, 2387 dense viridi-pulveraceum.

Hab. in ramis *Taxodii distichi*, Santee Canal Carol. inf. Amer. bor. (RAVENEL). — Pulcrum sed fructificatio ignota.

99. **Cenangium chlorellum** Fr. Elench. II, pag. 21. — Cæspitosum, 2388 minutum, sessile, cupulare, æruginoso-pulverulentum, ore collabente.

Hab. ad cortices Abietis excelsæ in subalpinis Europæ mediæ (MOUG.). — Facillime dignoscitur pulchella hæc et distinctissima species cupulis, ut *Calicii chlorelli*, æruginoso-pulverulentis, quo color interior nigrescens obliteratur; individua 5-8 e cortice cæspitosim erumpunt, sed basi non in corpus solidum connata, ut in *Scler. Ribis* etc. Discus nigrescere videtur; equidem rite explicatum observari non potuit.

100. **Cenangium æruginosum** Schw. Syn. Am. n. 1999. — Crustæ 2389 atro-æruginosæ lignum obtegenti et tingenti insidens, cæspitosim aut in lineas flexuosas aggregatum; cupulis minutis applanatis, mar-

gine undulato-plicato tuberoso; disco applanato pulverulento unicolore atro-aeruginoso; vix 1 mm. alt.

Hab. in ligno udo decorticato aut sub cortice, Bethlehem Americae borealis.

101. **Cenangium nigro-fuscum** Schw. Syn. Am. n. 2006. — Sparsum 2390 aut subcæspitosum, gregarium, globosum aut turbinatum, liberum quasi in epidermide, ad 4 mm. latum, undique tuberosum et quasi fusco-barbatum pube brevi rigida; disco vix aperto, sed dehiscens in rimis varie tortis.

Hab. ingenti copia ad dejectos ramos ignotos, Bethlehem Amer. bor. — An hujus generis fungus distinctissimus?

102. **Cenangium tumorum** Schw. Syn. Am. n. 1997. — Prorumpens, 2391 primum tectum, demum epidermide stellatim fisco cinctum, sparsum nec cæspitosum, rotundatum, convexum, nigrum, extus squamu-losum, primum omnino clausum, demum rimis aut irregulariter rupto ore apertum.

Hab. in excrescentiis ramorum passim, Bethlehem Amer. bor.

103. **Cenangium episphærium** Schw. Syn. Amer. bor. n. 1988. — Dif- 2392 forme, subcylindrico-ventricosum extus tuberosum, inæquabile, apice truncato-impresso aut irregulariter aperto, nigro-fuscum, intus subgrumosum.

Hab. in lignis igni destinatis in Amer. bor. — Aliquo modo analogum *Hydno* aut *Radulo aterrimo*. Vulgare in *Diatrype stigmate* et *Eutypa limæformi*, præsertim non evolutis. Epidermidem valde elevat et ad dejectionem præparat.

Species inquirendæ.

104. **Cenangium pulvis** Reb. Neom. p. 388 (Pez.). — Gregarium, mi- 2393 nutissimum, substipitatum, cupulis subglobosis inapertis, olivaceis, dein nigris.

Hab. in ramis putrescentibus in Neomarkia.

105. **Cenangium hypodermium** DC. Fl. fr. 6, p. 19 (Pez.). — Cupulis 2394 sessilibus, subpulverulenti, adultis cylindricis, brevibus truncatis crassis.

Hab. sub epidermide Cerasi in Gallia. — An a *Cen. Cerasi* diversum?

106. **Cenangium? Thymi** Fuck. Symb. p. 271. — Ascomatibus . . . 2395 stylosporeis; sporulis allantoideis, 4 = 1,5.

Hab. in ramulis vivis Thymi Serpylli pr. Oestrich Rhenogoviae.

107. **Cenangium alniellum** Rehm in Hazsl. Einig. Disc. p. 10, t. III, ²³⁶ fig. 4. — Cupulis solitariis vel cæspitosis, e piriformi patellatis, $\frac{1}{2}$ -1 cm. lat. intus albis, farinosis, disco atro-brunneo; ascis clavatis, deorsum longe attenuatis; sporidiis cylindraceis, hyalinis, continuis, utrinque rotundatis, rarius clavatis, 15-18 \approx 3.

Hab. in ramis Alni incanæ prope Iglo Hungariæ.

108. **Cenangium alnicolum** Hazsl. Einig. Discom. p. 9, Magyar. Szab. ²³⁹⁷ Disc. t. VI, f. 14. — Cupulis minutis patellatis, atro-brunneis, solitariis vel cæspitosis, disco plano dein convexo, levi, nigro fere immarginato; ascis cylindraceis, aliis spermatiophoris, aliis sporia fusoidea, obtusa, brunnea, 14-16 μ . long. foventibus.

Hab. in ramis Alni prope Hrabko et Eperies Hungariæ. — An e sect. *Phæangii*?

109. **Cenangium ilicinum** Dur. et Lév. Expl. Sc. Alg. tab. 29, fig. 4 ²³⁹⁸ (absque diagnosi). — Ascomatibus cæspitosis, erumpentibus, cupulatis, exiguis, extus atro-cinereis, pruinosis, disco concaviusculo atro, nitido; ascis clavatis brevissime stipitatis, octosporis; sporidiis fusoideis, distichis, hyalinis; paraphysibus filiformi-clavulatis, pluries divisis.

Hab. in ramis corticatis Quercus Ilicis, in Algeria.

TYMPANIS Tode Meckl. I, p. 23, (1790), emend. Sacc. Gen. Disc. p. 8 [Etym. *tympanon*]. — Ascomata erumpenti-superficialia, saepius cæspitosa, mox scutellari-aperta, nuda vel furfuracea, coriaceo-cornea, firma, fusca vel atra. Ascii paraphysati, vel myriospori, vel myriospori simul et octospori. Sporidia (numerosa) spermatioidea oblonga perexigua, vel (octona) majora, oblonga, continua, rarissime (spurie?) 1-septata, hyalina.

1. **Tympinis conspersa** Fr. Syst. II, p. 175, Tul. Carp. III, p. 153, ²³⁹⁹ Ann. N. S. 1853, t. XVI, f. 15-16, *Peziza Piri* Pers. Syn. p. 671, *P. Aucuparie* et *sphæroides* Pers. Myc. I, p. 327, *P. conglomerata* Wahlenb., *P. sphæriæformis*, Rebent. *P. sphæroides* Roth Nees Syst. f. 281. — Ascomatibus subrotundo-cæspitosis, stipatis, mox epidermidem transversim scissam infarcentibus, erectis, atris nudis pulvereque albo adspersis, biformibus, aliis (spermogoniciis) cylindricis obtusis corniculatis, aliis e basi cylindrica obovatis in verticem umbilicato-depressum disciformem dilatatis, tenuissime rugulosis, dein discum nigricantem recludentibus; ascis tereti-clavatis, apice rotundatis, 115-180 \approx 18-20, myriosporis; spori-

diis spermatioideis, ovatis, flavidulis, $1-2 \approx 0,5$; paraphysibus furcatis, septatis, apice clavulatis, 5μ . cr. brunneolis.

Hab. ad ramulos languidos Sorbi Aucupariæ, Cratægi, Piri, Pruni, Alni, Populi, Betulæ aliarumque arborum in Germania, Gallia, Suecia, Italia, Belgio, Britannia, Amer. bor. — Etiam in hac specie forma octospora indicatur. — Var. **rugulosa** Fuck. Symb. p. 272: cupulis rugulosis, opacis, fuscis. In ramulis Sambuci pr. Bonn Germaniæ.

2. **Tympanis Aucupariæ** (Pers.) Wallr. Crypt. n. 2348, *Cenangium* 2400 *Aucupariæ* Fr. Syst. II, p. 181, *Peziza Aucupariæ* Spr., *Dothidea cæspitosa* Fr. *Sphæria Aucupariæ* Pers. Syn. p. 51, *Sph. cæspitosa* Tod. Meckl. I, p. 41, t. 14, f. 113. — Ascomatibus cæspitosis, nigris, albido-pulverulentis, e basi implexa elongata subcylindrica in verticem angustum primum clausum dein apertum desinentibus; ascis clavatis, myriosporis; sporidiis perexiguis; paraphysibus filiformi-clavulatis apice brunneolis, conglutinatis.

Hab. ad ramos Sorbi Aucupariæ in Germania, Suecia, Britanua, in Piro americana in Amer. bor. — Teste Rehm vix vel non diversa a *T. conspersa*.

3. **Tympanis polygonia** (Fuck.) Sacc. *Cen. polygonium* Fuck. Symb. 2401 App. II, p. 55. — Cupulis erumpentibus, raro sparsis, plerumque dense cæspitosis, basi in stipitem communem crassum connatis, coriaceis, cæspitulis 2-3 mm. lat., e 3-8 cupulis compositis, singulis 1 mm. lat., extus atris, 3-6-gonis, angulis obtusis, valde prominentibus, vertice conniventibus, inde marginem dentatum referentibus, siccis arcte clausis, disco minuto, carioso, vix concavo, humectato pallide-olivaceo, sicco atro; ascis clavatis, multisporis, $108 \approx 20$; sporidiis ovato-cylindraceis, subcurvatis, $3 \approx 1$, pallidissime flavescentibus; paraphysibus copiosis, simplicibus, anguste clavatis, 2-3-septatis.

Hab. in ramis adhuc stantibus Piri Mali prope Estrich Rhenoviæ. — Cfr. *Cenangium polygonium*.

4. **Tympanis Prunastri** (Fuck.) Rehm Disc. p. 266. — Gregaria in- 2402 nato-erumpens, cæspitulosa, demum brevissime crasse stipitata, urceolata dein patellaris, atro-nitida, cornea 0,4-1,2 mm. lat.; disco piano marginato; ascis tereti-clavatis, apice rotundatis, $130-150 \approx 12-15$, 8-sporis simul et myriosporis; sporidiis octonis ovoideis, (spurie?) 1-septatis, hyalinis 8-10 $\approx 3-3,5$ monostichis appendicula filiformi auctis; sporidiis spermatoideis oblongis 2-3 $\approx 1-1,5$ flavidulis; paraphysibus filiformi-clavulatis, apice 5μ . cr. brunneis.

Hab. in ramis siccis *Pruni spinosæ* in Germania media.

5. **Tympanis gyroza** B. et C. North Am. Fungi n. 761. — Fascicu. 2403
lata; cupulis arcte compressis, gyrosis; ascis nondum visis.

Hab. in ramis dejectis Piri Mali, Virginia, New England Amer. bor.

6. **Tympanis Rosæ** Karst. Symb. Myc. Fenn. XXIII, p. 6. — Cupu- 2404
lis sessilibus vel subsessilibus, solitariis vel cæspitulosis, nudis, ni-
gris, planiusculis, margine subacuto, subdenticulato-inæquali, ni-
tentibus, 0,4 mm. lat.; ascis cylindraceo-subclavatis, myriosporis,
75 = 9-10; sporidiis oblongis vel ellipsoideis, 2 = 0,5-1; paraphysi-
bus gracilibus apice conglutinatis.

Hab. in ramis Rosæ volvatæ in horto Mustiala Fenniæ.

7. **Tympanis plicato-crenata** (Schw.) Fr. Syst. Myc. II, pag. 175, 2405
Pez. plicato-crenata Schwein. Carol. n. 1257. — Substipitata,
opaca, atra, margine obtuso profunde crenato plicatoque, disco-
concaviusculo.

Hab. in cortice *Pruni* in Carolina superiore. — Sparsim e
cortice erumpunt greges minores, in siccis iterumque humectatis
aterimi, sed in vivis «discus cinerescens». Stipes in cortice omui-
no delitescit, quare sessilem potius dicerem. Optime dignoscitur
margine eleganter crenato et plicato-sulcato discoque compacto,
sed fatiscente nec polito. Cfr. *Phacidium plicatum* Fr. Elench.
II, pag. 132, ex quo videtur hæc diagnosis ad *Cen. Andromedæ*
spectare.

8. **Tympanis Ligustri** Tul. Carp. III, p. 154, Sacc. Fung. it. f. 1304, 2406
Cenangium Ligustri (Tul.) Fuck. Symb. pag. 268. — Erumpenti-
superficiale, elongatum, nitidulum, atrum, cæspitulosum, patellare,
disco concavo-plano marginato, 0,5-08 mm. lat.; ascis tereti-cla-
vatis, apice rotundatis, 95-115 = 12-15, myriosporis; sporidiis glo-
buloso-ellipsoideis, 2,5 = 2, chlorinis; paraphysibus furcatis, septa-
tis, apice clavulatis, 6 μ . cr. brunneolis.

Hab. in ramis *Ligustri vulgaris* in Germania, Gallia, Bri-
tannia. — Var. *japonica* Sacc. Mich. I, p. 58: ascis 100-110 = 18;
sporidiis botuliformibus 4 = 1, hyalinis. In ramis *Ligustri ja-
ponici* in Ital. bor.

9. **Tympanis Syringæ** Fuck. F. rhen. n. 2144, Rehm Disc. pag. 270, 2407
Cen. Syringæ Fuck. Symb. p. 270. — Cupulis sparsis, atris, 1-2
mm. diam., primo clausis, demum apertis, margine 4-7-lacerato,
disco atro; ascis elongatis, stipitatis, operculatis, 112 = 16; spori-
diis spermatiiformibus minutissimis, cylindraceis, curvatis, oscil-

lantibus 2-3 \approx 1,5; paraphysibus simplicibus, linearibus, ascis longitudine, apice 5 μ . cr. brunneis.

Hab. in ramis siccis *Syringæ vulgaris*, prope Hattenheim Rhenogoviæ.

10. **Tympanis Fraxini** (Schw.) Fr. Syst. Myc. II, p. 174, *Pez. Fraxini* Schwein. Carol. n. 1262. — Subsessilis, turbinato-truncata, nitida, atra, disco plano rugoso marginato 1 mm. lat.; ascis terti-clavatis, apice rotundatis, 90-100 \approx 10-12, myriosporis; sporidiis oblongis, rectis, chlorinis, 3 \approx 1-5; paraphysibus furcatis, septatis apice clavulatis, brunneolis, 5 μ . cr.

Hab. ad ramos exsiccatos fraxineos in Suecia, Germania, Britannia, Amer bor. — Specimina pauca fasciculatim erumpunt, e basi sursum ampliata, nitida; margine prominente obtuso flexuoso; disco opaco, solidiusculo, atro, punctato-rugoso. Cfr. Rehm Discom. pag. 266.

11. **Tympanis turbinata** Schw. Syn. Am. bor. n. 1969. — Substipitata, primum erumpens ex epidermide capitulo globoso, sparsim sita et quasi tuberculis echinata, fusco-nigra, demum turbinato-obconica, marginibus tumidulis, fere clausis, disco poriformi, tandem apertiusculo, extus striata 2 mm. alt.; ascis obtusis; sporidiis numerosis (Berk.)

Hab. in ramulis exoletis *Diervillæ trifidæ* et *Viburni prunifolii*, Bethlehem et Center Amer. bor.

12. **Tympanis fasciculata** Schw. Syn. Amer. bor. n. 1967, Rehm 2410 Ascom. n. 423. — Subsessilis, minor, fasciculatim erumpens, epidermide cincta; cupulis fusco-nigris, varie tortis difformibus, marginibus inflexis subuplicatis, disco excavato, junior clausa, subsphæriæformis; ascis clavatis, 100 \approx 5, myriosporis; sporidiis elongatis, obtusis, hyalinis 4 \approx 2; paraphysibus filiformibus epithecium virens formantibus.

Hab. in epidermide ramorum *Viburni dentati* in Pensylvania et New Jersey Amer. bor.

13. **Tympanis antarctica** Fung. Fueg. n. 346. — Dense cæspitosa, per rimas lineares transversas peridermii erumpens, 10-50 cupulis in quoque cæpituolo, minuta 250-350 μ . diam., primo sphæroideo-clausa dein irregulariter aperta extus atra subcarbonacea glabra, intus vix pallescens; ascis cylindraceo-clavatis, apice obtuse lateque rotundatis, deorsum brevissime crasseque noduloso-stipitatis, 75-80 \approx 10-14, myriosporis, paraphysibus filiformibus longioribus dense obvallatis; jodi ope nulla; sporidiis inordinatis, conglobatis,

muco immersis, minutis, cylindraceis, rectis vel allantoideis, 2-2,5
 \approx 0,7-1, utrinque plus minusve rotundatis, hyalinis.

Hab. ad ramos emortuos Fagi betuloidis in silvis prope Agaia Fuegiæ.

14. **Tympanis mutata** (Fuck.) Rehm Disc. pag. 270, *Cen. mutatum* 2412
 Fuck. Symb. p. 270. — Cupulis erumpentibus, in acervulos oblongos valde dense congestis, rarius sparsis, primo punctiformibus, globosis, mox explanatis, convexis deflexisque, 2 mm. latis, disco sicco aterrimo, humefacto pallide fusco sordidove, emarginato; ascis a corpusculis ut in *Tymp. pitya* etc. repletis.

Hab. in ramis siccis Betulæ albæ in silva Estrich Rhenogoviæ.

15. **Tympanis populina** (Fuck.) Sacc. Gen. Disc. pag. 8, *Cen. populinum* 2413
 Fuck. Symb. p. 268, Sacc. F. ital. f. 1305. — Cupulis superficialibus, sparsis cæspitosisve, aterrimis, 1-2 mm. latis, sessilibus, disco concavo, demum plano, concolore, repando-marginato; ascis oblongis, tunica crassa, 72-100 \approx 12-15; sporidiis minutissimis, cylindraceis, curvatis, 3-4 \approx 1, hyalinis.

Hab. ad corticem ramorum Populi tremulæ, fastigiatae, laurifoliæ etc. in Germania, Belgio, Sibiria, Italia. — Status spermogonici sunt *Phoma Crepini* Speg. et *Dothichiza populina* Sacc.

16. **Tympanis saligna** Tod. F. Meckl. I, p. 24, t. IV, f. 37, non *Cen. angium* 2414
 Fuck. Symb. p. 268, ? *Peziza Todeana* Spr. Syst. IV, p. 515. — Ascomatibus gregariis subcæspitosis, atris, nitidulis, primum globularibus dein elongatis, verticem obtusum truncatum clavum demum in discum concavum apertum informantibus, 0,5-1,5 mm. lat.; ascis tereti-fusoideis, 90-110 \approx 15-18, myriosporis; sporidiis elongatis rectis, chlorinis, 3 \approx 1,5; paraphysibus furcatis, supra clavatis, 5 μ . cr. brunneolis.

Hab. ad ramos salicinos siccios rarius in Germania, Suecia, Lapponia, Belgio.

17. **Tympanis alnea** (Pers.) Fr. Syst. Myc. II, p. 174, *Peziza alnea* 2415
 Pers. Syn. p. 673, Myc. Eur. p. 325, *Tymp. conspersa* f. *alnea* Karst. M. F. I, p. 227, *T. conspersa* f. *nitida* Fuck. Symb. p. 272. — Substipitata, opaca, umbrino-nigrescens, cornea; cupulis subflexuosis obsolete marginatis, in cæspitulos 3-4 mm. lat., 3-20 congestis, disco circulari saepe inæqualiter marginato, 0,3-0,5 mm. lat.; ascis tereti-clavatis, 150-160 \approx 15-20; sporidiis spermatioideis innumeris, allantoideis 3 \approx 1,5; paraphysibus filiformi-clavulatis, apice 6 μ . cr. brunneolis.

Hab. ad ramos Alni incanæ et viridis in Suecia, Germania, Britannia. — Parva, exsiccata disco excavata vel potius margine magis prominente distincta, non vero clausa.

18. **Tympanis hysteroides** Rehm Disc. p. 268. — Gregaria, saepe 2416 solitaria, innato-erumpens, parum emergens, scutellata, nitido-atra, 0,8-1,5 mm. lat., cornea, dura, stellata vel hysteroideo-contorta, disco oblongo, plano; ascis tereti-clavatis, myriosporis; sporidiis spermatioideis oblongis, curvulis, flaveolis, $3 \approx 1,5$; paraphysibus furcatis, septatis, apice 5 μ . cr. clavatis, brunneolis.

Hab. in ramis siccis Alni viridis, Ötzthal Tyrolis.

19. **Tympanis corylina** (Sacc.) Rehm Disc. p. 270, *Chilangium* Sacc. 2417 F. ital. f. 1303. — Gregaria, erumpens, atro-nitens, 1-1,2, scutellata, disco planiusculo; ascis fusoideis, myriosporis; sporidiis oblongis, obtusis, curvulis, continuis, hyalinis, $6-8 \approx 1$.

Hab. in ramis corticatis Coryli Avellanæ in Arduennis (LIBERT).

20. **Tympanis laricina** (Fuck.) Sacc. *Cen. laricinum* Fuck. Symb. 2418 p. 270, *Tymp. pinastri* Rehm Disc. p. 272, nec Tul. — Cupulis sparsis cæspitosive, erumpentibus ex urceolato patellaribus 0,5-1 mm. lat., totis atro-nitidulis; ascis tereti-clavatis, $90-130 \approx 9-12$, myriosporis; sporidiis spermatoideis elongatis, rectis, chlorinis, $2-2,5 \approx 1$; paraphysibus furcatis, apice clavulatis, 5 μ . cr. brunneolis.

Hab. in ramulis siccis Laricis europææ et Abietum in Germania, Italia, Britannia, Hungaria, Galicia. — Fungus pycnidicus est *Cenangium bullatum* Alb. et Schw. Consp. p. 344 seu *C. Pinastri* b. *monstrosum* Fr. Syst. Myc. II, p. 184: cupulis rotundatis, clausis, demum perforatis, nigris, rugulosis; stylosporis longis, filiformibus, tenuissimis, curvatis, in massa gelatinosa lutea profluentibus.

21. **Tympanis Abietis** Cr. Finist. p. 43. — Ascomate 1-2 mm. diam., 2419 brunneo-nigrescente, sessili vel turbinato, parum concavo; ascis cylindraceis; sporidiis sphæroideis numerosissimis; paraphysibus filiformibus, superne furcatis et parum incrassatis.

Hab. in cortice emortuo Abietis, Finistère Galliae.

22. **Tympanis pitya** (Fr.) Sacc., nec Karst. *Cenangium* Fr. Syst. Myc. 2420 II, p. 184, Fuck. Symb. p. 269, *Tryblidium* Fr. Summa p. 369. — Minutum, denudatum, nitidulum, cæspitulosum, subsessile, leve opacum, demum denticulato-dehiscens, disco nigro; ascis cylindraceo-clavatis, $100 \approx 18$; paraphysibus bacillaribus obvallatis; sporidiis spermatioideis, curvulis $3 \approx 1/2$, inæqualibus, hyalinis farctis.

Hab. in ramulis corticatis emortuis Pini silvestris, Abietum in Germania, Suecia, Italia. — Quoad pycnidium cfr. Fuck. loc. cit. — ***T. juniperina** Sacc. Mich. I, pag. 419: ascis 80-90 = 10-11, ceterum typo similibus, æqualiter paraphysatis et myriosporis; ascomatibus similibus vel paullo majoribus, opace nigris. In cortice trunci Juniperi communis in silva Cansiglio Italiæ borealis.

23. **Tympanis seriata** Schw. Syn. Amer. bor. n. 1968. — Stipitata, 2421 concrescens longis seriebus primum *Sphæriam* piriformem nigram majusculam rugosam in oram prolongatam referens, tum *Pezizam* ore contracto, stipite crasso cylindrico, marginibus tumidulis atrovirentibus æmulans; disco velato, velo pulverulento albescente-viridi, ipso atro-virente; ultimo statu valde dilatata disco tuberculato, margine viridi-pulverulento, lobato, undulato.

Hab. in ligno decorticato, longis seriebus, Bethlehem Amer. bor. — Ascomata 2 mm. vix superantia.

24. **Tympanis cinerescens** Schw. Syn. Amer. bor. n. 1966. — Ses- 2422 silis, majuscula, gregaria, prorumpens ex epidermide elevata, cupulas cingente; cupulis rotundis aut difformibus lobatisque et subconfluentibus; disco marginato, impresso aut umbonato et etiam tuberculoso, velo cinereo, primum clauso, demum evanido; tunc omnino nigra, pulvere seminali tecta.

Hab. in putridissimo ramo ignoto horti Easton Amer. bor.

AMEGHINIELLA Spieg. F. Fueg. n. 347 (1888) [Etym. a cl. F. AMEGHINO palæontologo austro-americ.]. — Ascomata primo parvula, tuberculum minutum cortice tectum efficientia, mox erumpentia ac explanato-expansa basi connata, sæpe confluentia, majora, tenuiuscula, membranacea-tenacella, flexilia, plicato-undulata, fusca v. atra. Ascii pusilli octospori, paraphysati; sporidia minuta cylindracea v. allantoidea, hyalina. — Genus singulare epithecii naturâ ac fabricâ inter *Rhizinam* ac *Bulgarieas* medium, vegetatione tamen ac fructificationis notis ad *Dermateas* accedens.

1. **Ameghiniella australis** Spieg. l. c. — Stromate parvulo, tuberculato- 2423 effuso, parvulo 3-5 mm. diam., primo peridermio tecto, dein transverse erumpente, atro; cupulis sessilibus primo parvulis, crassiusculis, rigidulis, mox explanato-expansis, majusculis 1-3 mm. diam., crassiuscule membranaceis 300-450 μ . crass., coriacellis, flexilibus, dense undulato-plicatis, margine acutis, repandis v. sublobulatis, atro-olivaceis, extus siccis, olivaceo-pruinulosis, udis viscosulis,

contextu interno rufesceni-ferrugineo; ascis pusillis, antice rotundato-subtruncatis, postice breviter crasseque pedicellatis 40-50 = 3-4,5, paraphysibus densis, fuscidulis, mucoso-diffluentibus obvallatis, octosporis; sporidiis cylindraceis v. allantoideis, utrinque obtusatis, 4,5-6 = 0,7-1 hyalinis (luce refracta chlorinis).

Hab. ad ramos emortuos Fagi betuloidis in Staten island et Fuegia tota. — Species summopere polymorpha, in juventute nunc *Cenangii* nunc *Tryblidii* cujusdam habitum simulans, per ætatem tamen mox dignoscenda: asci centrifuge evolvuntur (jam prima juventute) margine tandem semper sterili. Iodi ops nulla.

EPHELINA Sacc. (Etym. *ephelis* ustio a sole), *Ephelis* Phill. Disc. p. 358, nec Fr., nec Sacc., *Excipula* Fuck. et Berk. p. p. — Ascomata tuberculiformia, demum patellata, ceracea v. subcoriacea, sessilia, minuta, atra crusta effusa subinde cornea nigricante insidentia. Asci elongati, paraphysati, octospori. Sporidia oblonga continua, hyalina. — An *Pyrenop. Gentianæ* (n. 1496) huc pertineat, dubium.

1. **Ephelina Rhinanthi** (Phill.) Sacc. *Ephelis* Phill. Disc. p. 358, t. XI, 2424 f. 69, *Rhytisma radicalis* Cooke in Grev. VIII, p. 9 (immat.) — Cupulis tuberculatis, congestis vel gregariis, crusta cornea effusa atra insidentibus, denique expansis, patellatis, subceraceis, nigris; ascis tereti-clavatis, 8-sporis; sporidiis oblongis v. clavatis, 3-guttatis, 10 = 5; paraphysibus filiformi-clavulatis.

Hab. in caule imo et radice *Rhinanthi Cristæ-galli*, Aberdeen, Forres, Whitchurch Britanniæ. — Ascomata $\frac{1}{2}$ mm. lat. extus rugulosa, intus pallide grisea. Adest st. stylosporeus; sporulis fusoideis 2-pluriseptatis, 70 = 5.

2. **Ephelina stromatica** (Fuck.) Sacc. *Excipula* Fuck. Symb. p. 400. 2425 — Ascomatibus in stromate spurio, tenuissimo, atro insidentibus, subconfluentibus, orbicularibus, planis, umbilicatis, astomis, nigris; ascis ovato-oblongis, curvatis, 8-sporis, 28 = 12; sporidiis ovato-clavatis, simplicibus, 10 = 4 hyalinis.

Hab. in caulis siccis *Silenes nutantis*, Oestrich Rheno-goviæ et in Jura (MORTHIER).

3. **Ephelina Viburni** (Fuck.) Sacc. *Excipula* Fuck. Symb. Myc. p. 400. 2426 — Ascomatibus sparsis, punctiformibus, in macula grisea, indeterminata, hemisphæricis, mox cupulæformibus, astomis, atris; ascis et sporidiis maturis nondum visis.

Hab. in foliis siccis Viburni Lantanæ, Vollrads Rhenogoviae et in Jura (MORTHIER).

4. **Ephelina gregaria** (Berk.) Sacc. *Excipula* Berk. Flor. New Zeal. 2427 vol. II, p. 202, Hook. Handb. New Zeal. Flor. p. 631. — Ascomatibus minutis, maculis brunneolis insidentibus, extus nigris, intus pallidis; ascis latis, clavatis; sporidiis obovato-oblongis, 10 μ . longis, plasmate demum obscurō bipartito.

Hab. ad paginam superiorem foliorum Gnaphalii in Nova Zelandia septentr. (COLENZO).

5. **Ephelina Galii** (Lasch.) Sacc. *Excipula* Lasch. in Kl. H. M. n. 678, 2428 Fuck. Symb. p. 401 et App., Sacc. Syll. III, p. 669. — Erumpens elliptica v. oblonga, rugulosa, atra, disco pallido.

Hab. in caule Galii Molluginis et G. silvatici in Rhenogovia et Borussia. — Ascii nondum maturi observati. *Phacidio vernali* affinis dicitur.

LAQUEARIA Fries S. v. Sc. p. 366 (1849) [Etym. *laqueare*]. — Ascomata subcoriaceo-subcornea, dimidiata, infra deficientia, decidua, pertusa discum tegentia. Discus ceraceus persistens. Ascii octospori, clavati. Sporidia ellipsoidea, continua, hyalina. — Genus, quoad locum, dubium. Cl. Rehm rectius forte in familiola propria (*Ostropæ*) cum *Robergea* et *Ostropa* enumerat.

1. **Laquearia sphæralis** Fr. Summa pag. 366, *Stictis* Fr. Syst. Myc. II, 2429 pag. 194. — Erumpens, hemisphærica, fusco-atra, urceolata, ore contracto integro, disco plano nigro; ascis oblongis, sessilibus, 8-sporis; sporidiis distichis minutis, oblongo-ellipticis, hyalinis, continuis (Fuck.)

Hab. ad ramos exsiccatos decorticatos Fraxini in Suecia, Germania, Britannia, ut videtur, rarissime. — Habitus *Tremato-sphæriæ* pertusæ, sed quoad magnitudinem, colorem et oris formationem cum *Godronia Urceolo* potissimum comparanda. Gregaria erumpenti-superficialis, limbo crasso persistente in urceoli formam contracto, ore regulari inflexo punctiformi-pertuso. Hymenium humectatum tumet, basin innatam modo tegit, unde patet limbum cupulæ v. peritheciæ forma gaudentem esse accessoriū. Transversim dissecta intus livido-nigra, demum elabens foveolam nigram in ligno relinquit.

2. **Laquearia Nylanderii** Cr. Finist. p. 46. — Ascomatibus sparsis, 2430 1 mm. diam., nigris, sessilibus, glabris, lecanidiformibus; ascis

Discomyceteæ, Dermateæ, Didymosporæ, Cenangella.

587

clavatis, dehiscentibus, octosporis; sporidiis cylindraceis, guttulatis; paraphysibus filiformibus, superne ramosis.

Hab. in cortice vivo Fagi, Finistère Galliæ.

Sect. 2. Phæosporæ Sacc.

HYMENOBOLUS Mont. Ann. Sc. Nat. 3. IV, p. 359 (1845), Syll. Crypt.

p. 191 [Etym. *hymenium* et *ballo ejicio*). — Ascoma erumpens, coriaceum, fusco-atrum, primum clausum, dein ore lacero subcon-
nivente ruptum. Hymenium concolor, ceraceum, atro-pulverulentum,
tandem prorsus elabens. Ascii tubulosi, octospori, denique elastice
dissilientes. Sporidia continua, oblonga, fuliginea. Paraphyses fili-
formes. — Genus hinc *Phacidieis*, illinc *Ascoboleis* affine, quoad
locum adhuc dubium.

1. Hymenobolus Agaves Dur. et Mont. Cent. V, n. 90, et Fl. Alg. 2431
ined., Atlas, t. 28, f. 2. — Characteribus iisdem ac generis; ascis
cylindricis, p. s. 100-120 = 8-9 apice, tunica integra, obtusiusculis,
breve sensim stipitatis, paraphysibus bacillaribus hyalinis obvallatis
octosporis; sporidiis oblique monostichis, oblongo-ellipsoideis, 15-17
= 8, fuligineis, guttula crassâ pallidiore fœtis.

Hab. sub epidermide foliorum Agaves americanæ emor-
tuorum in collibus Babel-Oued et Alger Africæ sept. (DURIEU et
TRABUT). — Innato-erumpens. Cupulæ sparsæ aut seriatæ, sub
epidermide rimosa foliorum enatæ, e cortice tandem erumpentes,
primitus clausæ, sessiles, extus intusque fuscæ, subtus planiusculæ
adnatæ, demum apertæ 2-4 mm. latæ, semimillimetrum in medio
crassæ, coriaceæ, siccæ fragiles, acute marginatæ, ore inflexo, laci-
niis epidermidis stellatim ruptæ patentibus aut strictis cinctæ.

Sect. 3. Didymosporæ Sacc.

CENANGELLA Sacc. Cons. gen. Disc. p. 9 (1884), Rehm Disc. 229

(Etym. demin. a *Cenangio*). — Ascomata erumpenti-superficialia,
subinde basi crassa substipitata, ex urceolato patellata, coriacea,
rarius subcornea, brunnea vel atra, sicca varie contorta. Ascii elongati v. clavati, 4-8-16-spori, paraphysati. Sporidia oblonga v. fu-
soidea, 1-septata, subhyalina, rarius colorata. — *Niptera Rhododen-*
dri (Ces.) De Not. (supra n. 1997), [= *Peziza* Ces. Bot. Zeit. 1854.
p. 186, = *Velutaria* Rehm Ascom. n. 253], præeunte ipso Rehm Disc.
p. 230, potius *Cenangellæ* speciem sistit.

I. *Eu-Cenangella*. Sporidia subhyalina.1. **Cenangella Pinastri** (Tul.) Sacc. *Tymp. Pinastri* Tul. Sel. Fung. 2432

Carp. III, p. 151, t. 19, f. 10-12. — Cupulis cæspitosis, raro solitariis substipitatis, nigris, primitus clausis et sphæroideis demum apertis concavis, margine obtuso vel evanescente, disco tandem dilutiore, nudis, circiter 0,5 mm. lat.; ascis elongato-clavatis, obtusissimis, subsessilibus, 100 \approx 11, 8-16-sporis; sporidiis ovoideis, uniseptatis, hyalinis, 8-9 \approx 3,5; paraphysibus filiformibus, simplicibus, apice clavatis fuscescentibus et conglutinatis.

Hab. in corticibus Abietis excelsæ et pectinatæ recens demortuæ in Fennia, Gallia, Germania, Italia, Belgio, Amer. bor. — Hypothecium (lamina tenui) fuscescens. Spermogonia vulgo cæspitosa, modo segregata, nunc ascomatibus mixta, primitus subsphæroidea, mucronulata, demum ovoidea, nigra, cavitate simplici confossa; sporulæ elongatae, simples, rectæ, 3,5 μ . long.; sterig mata filiformia, ramis strictissimis multibrachiata, 35 μ . long.

2. **Cenangella pitya** (Karst.) Sacc. *Typanis hypopodia* Nyl. Obs. 2433

p. 72 (teste Karst.), *Typanis pitya* Karst. M. F. I, p. 228. — Ascomatibus erumpentibus, solitariis vel in cæspitulos sociatis (2-4 insimul), demum stipitatis, nigris nudis, primitus clausis et sphæroideis, deinde apertis et concavis, tandem sæpe convexis, margine demum acutiusculo, subdenticulato-inæquali, circiter 0,6 mm. lat.; stipite crasso, sensim in cupulam abeunte, 0-1-0,5 mm. alt. circiter 0,3 mm. crass.; ascis elongato-clavatis, obtusissimis, subsessilibus, 8-16-sporis, 90 \approx 8-10; sporidiis ovoideo-oblongatis, uniseptatis, incoloribus, 8-10 \approx 2-3; paraphysibus filiformibus, gracilibus, adspersis, apice clavatis fuscescentibus et conglutinatis.

Hab. ad corticem levem Pini silvestris in Fennia et in Lapponia. — Spermogonia erumpentia, gregaria, solitaria, rarius subcæspitosa, subsphæroidea, cavitate simplici confossa, nigra, minuta; sporidia elongata, recta vel subrecta, incoloria, simplicia 2 \approx 0,5.

3. **Cenangella Piceæ** (Pers.) Sacc. *Peziza* Pers. M. E. I, p. 269, *Mollisia* Quél. Ench. p. 318, *Peziza abieticola* Nyl. Pez. fenn. p. 52,

Dermatea Piceæ Rehm Disc. pag. 257, *Cenangium farinaceum* Fuck. Symb. p. 269. — Cupulis sparsis, erumpentibus, sessilibus, 1-2 mm. latis, primo clausis subglobosis, demum apertis, concavis, disco pallide fusco, extus fuscis, albo-farinaceis subglabrisve, margine incurvo; ascis clavatis, stipitatis, 8-sporis, 90-112 \approx 10-15;

sporidiis in ascis superiori parte inordinatis, lanceolatis, plasmate medio diviso, 2-cellularibus, hyalinis, 16-18 \approx 5-6.

Hab. in pag. inf. foliorum Abietis pectinatae in Rhenogovia, Tyroli, Helvetia, Fennia.

4. **Cenangella dolosa** Sacc. *Consp. gen. Disc.* p. 9, *Cenangium dolosum* Sacc. et Sp. Mich. I, p. 58. — Cupulis sparsis corticolis, innato-erumpentibus, initio et siccis clausis, udis urceolato-apertis fibrosis, coriaceis, duris, aterrimis $1\frac{1}{2}$ - $3\frac{1}{4}$ mm. diam.; ascis clavatis breve crassiuscule stipitatis, 80-100 \approx 10-13; paraphysibus filiformibus obvallatis, octosporis; sporidiis subdistichis, biconico-fusoideis 22-25 \approx 6-7, medio constricto 1-septatis, rectis curvulatis, 4-guttulatis, hyalinis.

Hab. in ramulis corticatis Coryli Avellanæ, Conegliano Ital. bor. (SPEGAZZINI).

5. **Cenangella stegioides** (Speg.) Sacc., *Dermatea stegioides* Speg. 2436 Mich. I, p. 471. — Ascomatibus primo subcutaneis, clausis, dein epidermide operculatim circumscissa sed permanente subvelatis, disciformibus, extus cinnamomeis, granuloso-tomentellis, disco cinereo v. subroseo, concavo, prominulo, $1\frac{1}{2}$ -1 mm. diam.; ascis cylindraceis; paraphysibus filiformibus obvallatis, 35-40 \approx 2-3, octosporis; sporidiis distichis, cylindraceis v. clavulatis, non v. tenuissime 1-septatis, ad septum non constrictis, hyalinis, 5-6 \approx 1- $1\frac{1}{2}$.

Hab. ad ramulos emortuos Quercus sessilifloræ Colfosco, Treviso Ital. bor.

6. **Cenangella Ravenelii** (Berk.) Sacc. *Tympinis Ravenelii* Berk. 2437 North Am. Fungi n. 760. — Sparsa vel fasciculata; cupulis breviter pedicellatis marginatis, disco cinereo; sporidiis biconicis commissura medioque constrictis, 34-36 μ . long.; paraphysibus clava obovata (conidio?) terminatis.

Hab. in ramis Carpinis et Ilicis in Carolina inf. et Alabama Amer. bor. — Prope septum sporidiorum subinde adest nucleus peculiaris.

7. **Cenangella Carpini** (Hazsl.) Sacc. *Cenangium Carpini* Hazsl. 2438 Einig. Disc. p. 10 et Magyar. Szab. Disc. t. VI, f. 15. — Eruinpens, obconica, subcaespitosa, dein dehiscendo discoidea, atro-brunnea, glabra; ascis fusoideis, tenuibus, octosporis; sporidiis anguste fusoideis, hyalinis, uniseptatis, latitudine 6-8-plo longioribus; paraphysibus tenuibus, filiformibus, apice brunneis.

Hab. in cortice Carpinis prope Eperies Hungariæ.

8. **Cenangella vernicosa** (Fuck.) Sacc. *Cenangium vernicosum* Fuck. 2439

Symb. p. 268, *Dermatea* Rehm Disc. p. 262. — Cupulis gregariis, innato-erumpentibus, dein subsessilibus, concavis, marginatis, aterrimis, nitidis, 1-2 mm. diam., disco nigro, opaco; ascis clavatis, apice rotundatis incrassatis, 70-80 = 10-12, 4-8-sporis; sporidiis fuscoideis, rectiusculis, dein 2-locularibus, 12-15 = 4-5, hyalinis; paraphysibus furcatis, septatis, apice clavulatis, brunneo-flavis 4 μ . crass.

Hab. in ramis corticatis Pruni Padi prope Biebrich Rhenogoviae et in Arduennis. — Fungus pycnidicus est *Sphæronœma polymorphum* Auersw.: cylindraceum, obtusum, $\frac{1}{2}$ mm. altum, aterrimum, nitidum, gregarium; stylosporis cylindraceis, rectis, minutis, hyalinis.

9. **Cenangella Ericæ** (Niessl) Rehm. Disc. p. 232, *Cenangium Ericæ* 2440 Niessl Beitr. p. 61, t. VII, f. 4-6. — Solitaria, nunc sessilis basi contracta, nunc substipitata, corrugata rugulosave, ore compresso-connivente, in statu humido aperta, nigra, 0,5-1 mm. lat., membranaceo-coriacea, disco griseo-albescente; ascis clavatis in stipitem brevem attenuatis, apice rotundatis, 8-sporis, 70-90 = 8-10; sporidiis distichis, oblongo-lanceolatis, utrinque attenuato-rotundatis, medio 1-septatis constrictisque, hyalinis, guttulatis, 15-18 = 3-4; paraphysibus ascorum longitudine, apice incrassatis interdum ramosis, hyalinis.

Hab. in ramulis foliisque siccis Callunæ vulgaris pr. Liezen et Voitsberg Stiriae et in montibus dell'Oropa Pedemontii. — Status pycnidicus, teste Niessl, est *Pestalozzia Callunæ* Ces.

10. **Cenangella Rehmiana** Sacc. *Dermatea Rhododendri* Rehm Disc. 2441 p. 248 *Helotium* Rehm Ascom. n. 109. — Ascomatibus per epidermidem fissam erumpentibus, plerumque gregariis, sessilibus v. brevissime stipitatis, primitus planis, orbicularibus, distincte marginatis, siccis subcorrugatis, extus fuscis, scabris, disco flavido vel viridi, contextu parenchymatico, fusco, margine pseudoprosenymatico, 0,5-2 mm. diam.; ascis clavatis, in stipitem elongatis, 8-sporis, 80-90 = 10; sporidiis ellipticis vel elongato-ovatis, 1-cellularibus, plerumque 1-2 nucleolis praeditis, denique bicellularibus, hyalinis, distichis, 15-18 = 4,5-5; paraphysibus filiformibus, haud clavatis, hyalinis, 2,5 μ .

Hab. ad ramulos sicos Rhododendri ferruginei in alpibus editis Tyrolensibus. — Nomen mutavi, quia generi præsenti adscribenda quoque homonyma *Cenangella Rhododendri* (Ces.) Rehm.

11. **Cenangella Fraxini** Sacc. Conspectus gen. Disc. pag. 9, *Fung. Ital.* 2442

f. 1437, (*Tympinis*) et forte *C. Carpini* Tul. Ann. Sc. nat. 1853 p. 140, (asci 8-spori, sporidia bilocularia!). — Erumpens, gregaria vel subcæspitosa, ex urceolato patellata, fuligineo-atra, marginata, 0,5-0,7 mm. lat., subcoriacea; ascis tereti-clavatis, breve stipitatis, filiformi-paraphysatis, 108-115 \approx 18-21, octosporis; sporidiis subdisticis, fusoideis, tenuiter 1-septatis, hyalinis, 26-29 \approx 10-12.

Hab. in cortice emortuo *Fraxini Orni*, Rouen Galliæ (LE-TENDRE). — Sporidia subinde extra ascos (abnormiter?) 3-septata fuscella.

12. **Cenangella urceolata** (Ellis) Sacc. *Cenangium urceolatum* Ellis 2443 in Grev. VI, p. 9. — Erumpens, prominens, olivaceo-atra; cupulis urceolatis, ad basin attenuatis, extus pilis brevibus flaccidis adspersis, ore contracto, disco pallide-fuligineo; ascis clavato-cylindraceis; sporidiis arcte fusiformibus, hyalinis, uniseptatis, 15 \approx 4.

Hab. ad ramos Clethræ alnifoliæ, New Jersey Amer. bor. — Ascomata 0,3-0,5 mm. lat. *Sphæronæma clethrincolum* Ell. videtur spermogonium.

13. **Cenangella æquinoctialis** (Mont.) Sacc. *Cenangium æquinoctiale* 2444 Mont. Guy., n. 262, in appendice et n. 452. — Gregaria, erumpens; excipulo coriaceo initio globoso clauso, helvolo-furfuraceo, tandem subcupulari-explanato; disco livido-fusco, pulvere copioso sulphureo velato; ascis cylindricis 8-sporis, 250 μ . long.; sporidiis ovoides, inæqualiter bilocularibus, monostichis, 30 \approx 15, limbo lato cinctis.

Hab. ad cortices in Guyana (LEPRIEUR). — Ascomata gregaria vel cæspitoso-confluentia, primo sphærica clausa, semen Cannabis magnitudine æquantia, dein hemisphærica aut pressione mutua irregularia, 1 ad $1\frac{1}{2}$ mm.; discus planus, marginem crassum interdum flexuosum æquans.

14. **Cenangella flavo-cinerea** (Phill.) Sacc. *Dermatea* Phill. in Grev. VII, p. 23. — Cupulis gregariis, sessilibus, initio hemisphæricis dein expansis; disco flavidò vel cyaneo-griseo, margine pilis brunneis brevibus fimbriato, extus avellaneo-brunnea; ascis clavatis 8-sporis; sporidiis fusoideis rectis curvulisve, 1-septatis vel 3-nucleatis, 10-20 \approx 4-6; paraphysibus numerosis tenuissimis.

Hab. in fragmentis ligneis in California.

15. **Cenangella radulicola** (Fuck.) Rehm Disc. p. 231, *Cenangium* 2446 Fuck. F. R. n. 2073, *Dermatea* Fuck. Symb. p. 278. — Cupulis sparsis, primo cylindraceis, clausis, demum apertis, 1-3 mm. latis, 1-1,5 mm. alt., calyciformibus, in stipitem crassum seorsum attenua-

tis, margine involuto, fuscis, striato-rugulosis, disco atro-olivaceo; ascis stipitatis, elongatis, 8-sporis, 128-150 \approx 16-18; sporidiis lanceolato-oblongis, rectis curvatisve, continuis, 2-4-guttulatis, dein 1-septatis, hyalinis, 18-26 \approx 6-8; paraphysibus simplicibus linearibus.

Hab. in ramulis Betulæ, *Radulo aterrimo* tectis, in Rhenogovia.

16. **Cenangella parasitica** (Fuck.) Rehm Disc. p. 232, *C. parasiticum* 2447

Fuck. Symb. App. I, p. 43. — Cupulis demum in pycnidii cæspitulæ ortis, orbicularibus seu irregularibus, subsessilibus, 1-2 mm. latis altisque, primo turbinatis, subclausis, demum perforatis, aperatis, tandem explanatis, extus margineque valde elevato, involuto, densissime, brevissime velutinis, pallide sordido-fuscis argillaceisve vel quandoque margine parum pallidiori, disco griseo, carnosæ; ascis linearibus, 8-sporis; sporidiis monostichis, ovatis, utrinque obtusis, didymis, medio constrictis, pallide flavis, 6-8 \approx 3-4.

Hab. parasitica in Sphaeronæmate cæspitoso ad Acer platanoides in Rhenogovia. — Sec. Fuck. *Sphaeronematis* species est pycnidium.

II. *Phœangella* Sacc. Sporidia colorata.

17. **Cenangella Aceris** (Hazsl.) Sacc. *Cenangium Aceris* Hazsl. Einig. 2448

Disc. p. 9, Magyar. Szab. Disc. t. XII, f. 13. — Gregaria, minuta, discoidea, erumpenti-superficialis, disco crasso, pallide brunneo, superficie atro-brunneo; ascis clavatis, 90-98 μ . long. breve stipitatis, octosporis; sporidiis elongatis vel ovato-oblongis, uniseptatis, brunneolis, 14-20 \approx 4; paraphysibus filiformibus.

Hab. in ramus Aceris tatarici in Hungaria (HAZSL) et Slavonia (SCHULZER).

18. **Cenangella Morthieri** (Fuck.) Sacc. *Cenangium Morthieri* Fuck. 2449

Symb. p. 272, t. IV, f. 47. — Cupulis cæspitosis seriatisque, primo tenuibus, valde excavatis, extus albo-farinaceis, demum disco crasso, plano, atro, distincte marginato; ascis oblongis, breviter stipitatis, 4-sporis, 80 \approx 10; sporidiis monostichis oblongis, utrinque parum attenuatis, subcurvatis, continuis uniseptatisve, 2-guttulatis, fuscis, 16 \approx 4.

Hab. in ramis corticatis Rhamni alpinæ in Jura Helvetiæ (MORTHIER). — Status pycnidicus est *Micula Maugeotii* Duby.

19. **Cenangella Ulicis** (Cooke) Sacc. *Dermatea* Cooke in Grev. III, 2450

p. 186. — Cæspitosa vel sparsa substipitata; cupulis initio turbinatis dein expansis, extus rufo-brunneis furfuraceis, 1-2 mm. lat.,

disco concavo obscuriore; ascis clavato-cylindraceis; sporidiis ellipsoideis 1-septatis, brunneis 12-5; paraphysibus subclavatis apice brunneolis.

Hab. in ramis Ulicis europæi, Shere Britanniæ. — Ascomma extus coffeicolor.

20. **Cenangella deformata** (Peck) Sacc. *Cenangium* Peck 28 Rep. 2451

Stat. Mus. p. 68. — Minuta, sparsa, vel congesta, initio inæqualis v. subspherica, dein apicè aperta discoidea, margine irregulari v. rupto, nigra; sporidiis stipatis ellipsoideis endochromate bipartito demum coloratis et 1-septatis, 26-32 μ . long.

Hab. in corticibus emortuis Juniperi virginianæ, Greenbush Amer. bor. — Species initio sphæriæformis, sæpius seriata.

21. **Cenangella Cesatii** Sacc. *Patellaria?* *Tympinis?* Ces. Born. p. 2452

11. — Tota aterrima, erumpens per greges lineares e ligno, primus turbinata clausa, serius patellaris, margine incrassato interdum crenulato; fructificatione ascospora compacta, sed ceterum incerta; sporidiis dipodiæformibus.

Hab. in corticibus Etich River in ditione Sarawak.

22. **Cenangella quercina** (Hazsl.) Sacc. *Cenangium quercinum* Hazsl. 2453

Ein. Discom. p. 7, Magyar. szab. Disc. t. VI, f. 10. — Sparsa, subcutaneo-erumpens, ex urceolato patellata, minuta; ascis linearis-fusoideis, paraphysatis; sporidiis fusoideis, uniseptatis, leniter constrictis, flavo-brunneis.

Hab. in ramis quercinæ prope Eperies Hungariæ. — Habitat *Peziculæ quercinæ*.

23. **Cenangella Potentillæ** (Hazsl.) Sacc. *Cenangium Potentillæ* Hazsl. 2454

Einig. Discom. p. 8, Magyar. szab. Disc. t. V, f. 11. — Erumpens, nigra, opaca, disco convexo pulvereo, margine prominulo, contextu griseo-flavo; ascis tereti-fusoideis, paraphysatis; sporidiis oblongo-fusoideis, uniseptatis, opace atro-brunneis.

Hab. in ramis Potentillæ fruticosæ, Eperies Hungariæ.

— Paraphyses crassæ pluriseptatæ.

24. **Cenangella aggregata** (Hazsl.) Sacc. *Cenangium* Hazsl. Ein. Disc. 2455

p. 9, nec Lasch. — Sparsa, urceolata, dein circulariter vel rimose dehiscens, demum discoidea; ascis clavatis, breve stipitatis, octosporis; sporidiis oblongis v. ovoideis, bilocularibus, brunneis; paraphysibus filiformi-clavulatis, coloratis.

Hab. in caulibus emortuis putrescentibus Odontites prope Bartfeld Hungariæ.

Sect. 4. **Phragmosporæ** Sacc.

SCLERODERRIS Fr. Syst. Myc. II, p. 178 (1822), ut subgen., De Not. Disc. p. 29 (Etym. *scleros* durus et *derris* pellis), *Cenangii*, *Tympanidis* etc. spec. Auct. — Ascomata erumpenti-superficialia, sæpius cæspitosa, ex urceolato patellata, sæpe margine lacero-fimbriata, pruinosa vel levia, coriacea vel cornea, brunnea vel atra raro læte colorata. Asci paraphysati, typice octospori. Sporidia elongata vel cylindracea, demum 3-pluriseptata, hyalina, rarius colorata. — Est mihi *Cenangium phragmosporum*.

I. *Eu-Scleroderris*. Sporidia subhyalina.

1. **Scleroderris ribesia** (Pers.) Karst. M. F. I, p. 215, *Peziza* Pers. 2456 Teut. p. 35, *Cenangium Ribis* Fr. Syst. Myc. II, pag. 179, Tul. Ann. Sc. Nat. 1853, t. 16, f. 9-11, Carp. III, pag. 164, t. XIX, f. 1-9. — Stromate ligno innato, primitus hypophloideo, sphæroideo, solido, intus pallido vel luteolo-virente, extus fulvo, e lateribus imis conidia ellipsoidea vel ovoidea, recta, simplicia, hyalina, 3,5-5 \approx 3 fundente, dein per epidermidem erumpente gignenteque ascomata et spermogonia; ascomatibus cæspitosis, 1-4 mm. lat., 1,5 mm. alt., breviter stipitatis primitus sphæroideis vertice depresso vel umbilicatis, clausis fuscis vel fulvescenti-fuscis, dein obverse turbinatis, ore connivente fimbriato dehiscentibus, rugulosis, subnudis, fusco-nigrescentibus, disco cinereo-albido; ascis cylindraceis, obtusis, deorsum longe attenuatis, 100-120 \approx 7-8, octosporis; sporidiis congregatis aciculari-elongatis vel aciculari-filiformibus, hinc obtusis, illinc acutis, 4-8-guttulatis dein tenuiter 4-septatis, chlorino-hyalinis, 30-36 \approx 3,5-4,5; paraphysibus filiformibus, simplicibus, apice vix clavato nonnihil corrugatis.

Hab. in ramis emortuis Ribis rubri, tristis et nigri in Suecia, Fennia, Britannia, Gallia, Germania, Belgio, Sibiria. — Stat. pycnidicus est *Fuckelia Ribis* Bon.; st. macrostylosporifer teste Fuck., est *Mastomyces Friesii* Mont.

2. **Scleroderris repanda** (Fr.) Rehm Disc. p. 209 (ut var.), *Cen. repandum* Fr. Elench. II, p. 22. — Gregaria, turbinata, fusco-pruinosa, demum nuda, levi, cervino-nigrescens, ore repando lacinatoque infexo, disco fusco; ascis sporidiisque ut in *Scl. ribesia*.

Hab. in ramis exsiccatis Ribis petræi in Vogesis (MOUGEOT). — Affinis *C. Ribis*, a qua differt cupulis tenuioribus, magis expli-

catis, ore repando laciniatoque in velum haud continuato; cupula demum nuda, trita subnitente discoque fuscescente.

3. **Scleroderris seriata** (Fr.) Rehm Disc. pag. 211, *Cenangium* Fr. 2458
S. M. II, p. 185, *Trybl. seriatum* Fries exs. n. 161, *Pez. truncatula* Rebent. Neom. p. 388 non Pers., *Dermatea ser.* Tul. Carp. III, pag. 160. — Gregaria, globoso-depressa, substipitata rugosa, nuda, atra, in lacinias obtusas dehiscens, disco fuligineo-atro; ascis ovato-elongatis, apice obtusis, longissime stipitatis 110 ± 12, 8-sporis; sporidiis anguste fusoideis, acutis rectis curvulisve, 55-80 ± 2-3, pseudo-multiseptatis, hyalinis.

Hab. in cortice Betulæ albæ in Suecia, Germania, Britannia, Belgio, Gallia. — Adsunt spermogonia et pycnidia. Cfr. Phill. Disc. p. 348. Per rimas corticis transversales erumpit, series 7-35 mm. longas lineares plus minus confertas sistens. Stipes brevis, inter corticem nidulat. Cupulæ opacæ; clausæ centro depresso, obtuse marginatæ, demum apertæ, in lacinias 4-6 fragiles hiantes.

4. **Scleroderris fuliginosa** (Pers.) Karst. M. F. I, p. 216, *Cenangium* 2459
Fr. Elench. II, p. 23, *Cen. difforme* Fr. in Moug. et Nestl. Stirp. vog. n. 889, *Sphæria fuliginosa* Pers. Syn. p. 52, *Tymp. fuliginosa* β. Wallr. Crypt. p. 430. — Cupulis in plagas latas effusas stipatis, difformibus, fuliginosis, disco pallido, subiculo late ambiente-effuso innatis; ascis tereti-clavatis, apice rotundatis 100-130 ± 9-10, 8-sporis; sporidiis cylindraceis utrinque acutis, 60-65 ± 2,5-3,5, rectis, 8-locularibus, hyalinis subparallelis; paraphysisibus filiformibus, hyalinis.

Hab. ad ramos emortuos Salicis capreæ in Suecia, Norvegia, Germania, Gallia. — Optime dignoscitur stromate sub epidermide cortici interiori innato, late ambiente effuso, humido nigrescente, sicco superne lutescente aut olivaceo; cupulæ simplices vel cæspitosæ, per epidermidem erumpentes, liberatæ, difformes, in perfectissimis tenues, coriaceo-membranaceæ, sublaciniatæ, pruinosæ, ferruginosæ, disco sublutescentes, in imperfectis minores, nigrescentes etc.

5. **Scleroderris concinna** (B. et C.) Sacc. *Cenangium concinnum* B. 2460
et C. North Am. Fungi n. 767. — Cupulis sessilibus subtiliter pulvрulentis marginatis; disco plano nigro; sporidiis biserialibus, oblongis, tri septatis, 15-16 μ. longis.

Hab. in cortice Quercus falcatae, Lauri Benzoin et Sassafras in Carolina et Alabama Amer. bor.

6. **Scleroderris bacillifera** (Karst.) Sacc. *Tymanis* Karst. M. F. I, 2461

p. 229. — Cupulis erumpentibus, sessilibus vel subsessilibus, solitariis vel saepius in cæspitulos sociatis (2-5 insimul), nudis, nigris, primitus clausis et sphæroideis, deinde a pertis et concavis, tandem planiusculis, margine obtuso, disco nitido, 0,5-0,6 mm. lat.; ascis clavatis, obtusis, subpedicellatis, 65-88 = 10-11, octosporis; sporidiis conglobatis, bacillaribus, vulgo apicibus attenuatis, flexuosis, simplicibus vel spurie tenuiter 5-7-septatis, 20-52 = 3-4,5; paraphysibus filiformibus, gracilibus, apice incrassatis fuscescentibus et conglutinatis.

Hab. supra corticem crassiorem truncorum Abietis excelsæ stantium, in Fennia saltem meridionali passim. — Spermogonia verrucariiformia, nigra, minuta; sporulæ oblongatae, simplices, incolores, 4 = 1,5.

7. **Scleroderris pitya** (B. et C.) Sacc. *Cenangium pityum* B. et C. 246
North Am. Fungi n. 765 nec Fr. — Erumpens, brunnea, margine obtuso inflexo subtiliter pulverulento; sporidiis lineari-oblongis, curvatis, 4-septatis, 20 μ . longis.

Hab. in cortice Abietum, New England Amer. bor. (SPRAGUE).

8. **Scleroderris Sequoiæ** (Plow.) Sacc. *Cenangium* Plow. in Grev. 246
VII, p. 23. — Cupulis gregariis turbinatis, nigris, margine connivente, disco nigricante intus cinereo; ascis late clavatis, octosporis; sporidiis ovoideis vel ovato-oblongis, simplicibus dein triseptatis, enucleatis, 25-30 = 7-15; paraphysibus furcatis tenuibus.

Hab. in trunco Sequoiæ giganteæ in California.

9. **Scleroderris sepiaria** (Karst.) Sacc. *Tympanis sepiaria* Karst. 246
Fragm. mycol. IX, p. 2. — Ascomatibus erumpenti-superficialibus sessilibus, interdum solitariis, subinde in cæspitulos sociatis (2-5 insimul), primitus clausis et subsphæroideis, deinde apertis, concavis, subinde ellipsoideis, obtusis marginatis, nudis subtiliter punctulatis, nigris, 0,3-0,4 mm. lat.; ascis clavatis, obtusis, subpedicellatis, 70-80 = 8-9,5 mm., octosporis; sporidiis oblique mono- vel subdistichis, ovoideo-oblongatis, simplicibus raro 1, rarissime 3-5 septatis, 12-18 = 3,5-4,5; paraphysibus filiformibus, 1,5 mm. crass., interdum ramosis.

Hab. ad lignum pineum prope Mustiala Fenniæ.

10. **Scleroderris amphibola** (Mass.) Gill. Disc. p. 198, *Tympanis am-* 246
phibola (Mass.) Karst. Mycol. Fenn. I, p. 230, *Pragmopora am-*
phibola Mass., *Cenangium sphæroides* Niessl herb. — Ascomatibus erumpenti-superficialibus, nigris, rugulosis, demum convexulis, immarginatis; ascis cylindraceo-clavatis, brevissime stipitatis,

apice tunica integra, rotundatis, 70-75 \approx 8-10, paraphysibus asco longioribus sursum paullum incrassatis guttulatis obvallatis, 8-sporis; sporidiis distichis vel oblique monostichis, fusoideis utrinque acutiusculis, 18-20 \approx 3 $\frac{1}{2}$ -4 $\frac{1}{2}$, 3-septatis leniter curvulis, hyalinis.

Hab. ad corticem Pini silvestris in Fennia, Gallia, Britannia, Italia. — Sporidia subinde 5-septata. Spermogonia nota.

1. **Scleroderris amphiboloides** (Nyl.) Sacc. *Tympinis* Nyl. Obs. 2466 p. 71, Karst. M. F. I, p. 230. — Sessilis, gregaria, subsolitaria, e sphæroideo demum patellata, concava dein plana, obtuse marginata, nuda, nigra, rugulosa, intus obscura, 0,4-0,5 mm. lat.; ascis elongato-clavatis, obtusissimis, sessilibus, 85-90 \approx 11-15, 8-16-sporis; sporidiis fusoideo-elongatis, spuriæ (?) 5-7-septatis, hyalinis, 15-25 \approx 2,5-4,5; paraphysibus filiformibus, gracilibus, apice incrassatulis dilute lutescentibus.

Hab. in ligno abietino vetusto in Fennia meridionali.

12. **Scleroderris hypopodiza** (Nyl.) Sacc. *Tympinis* Nyl. Obs. p. 71, 2467 Karst. M. F. I, p. 231, Rehm Asc. Lojk. n. 61. — Cupulis cæspitosis, rarius solitariis basi substipitato-constrictis, primitus clavis, deinde apertis, concavis, tandem convexulis et immarginatis, nudis, nigris, subrugulosis, inæqualibus, intus concoloribus, 0,5-0,8 mm. lat.; ascis elongato-clavatis, obtusissimis, 80 \approx 15, 8-24-sporis; sporidiis fusoideo-elongatis, spuriæ tenuiter pluri- (vulgo 7-) septatis, incoloribus 18-23 \approx 3,5-4; paraphysibus gracilibus, sursum leviter incrassatis.

Hab. ad lignum pineum (et quercinum?) vetustum in Fennia et Hungaria. — Spermogonia vulgo cæspitosa, segregata vel cupulis immixta, obovoidea vel vertice conoideo-attenuata, nigra; sporulæ elongatæ 3-4 \approx 0,5. Sec. Rehm asci 100 \approx 10-15, 8-spori; sporidia 18-24 \approx 3, disticha. Rehm Asc. n. 358 sistit formam myrioporum, an hujus speciei?

13. **Scleroderris aggregata** (Lasch) Rehm Disc. pag. 212, *Sphæria* 2468 Lasch Bot. Zeit. 1857, p. 428, *Cenangium* Fuck. Symb. p. 271. — Cæspitoso-seriata, coriaceo-cornea erumpens, atro-nitida urceolata, demum quadrifido-dehiscens 0,5-1 mm. lat.; disco pallido, concavo; ascis clavatis, 75-80 \approx 7-9, octosporis; sporidiis elongato-fusoideis, obtusiusculis, curvulis, 3-septatis, 4-guttatis, 30-33 \approx 3-4, distichis, hyalinis; paraphysibus filiformibus 2 μ . cr. hyalinis.

Hab. in caulis siccis Euphrasiæ officinalis in Germania media. — Teste Fuck., adsunt stylosporæ teretes, curvæ, pluriguttatæ, 1-septatæ, 18 \approx 4, hyalinæ.

14. **Scleroderris nigro-rufa** (Berk.) Sacc. *Excipula nigro-rufa* Berk. 2469

Fl. New Zeal., vol. II, p. 202, t. 106, fig. 11, Hook. Handb. New Zeal. Flor. p. 630. — Subglobosa, extus nigra, intus rubra; ostiolo infexo; ascis latis, clavatis, obtusis; sporidiis hyalinis, obovato-oblongis, obtusis, obliquis, 4-septatis.

Hab. ad paginam inferiorem foliorum Pittospori crassifolii, in Nova Zelandia septentr. (COLENZO).

15. **Scleroderris pallidula** (Cooke) Sacc. *Dermatea pallidula* Cooke 2470

Grev. XVI, pag. 70. — Erumpens; cupulis convexis, subcæspitosis (2-4), testaceo-carneis, demum pallescentibus 1-2 mm. lat., deorsum in stipite brevi communi attenuatis; margine obsoleto; ascis clavatis, octosporis; sporidiis ellipsoideo-lanceolatis, utrinque obtusis, triseptatis, hyalinis, 25 = 9; paraphysibus sursum incrassatis, brunneis.

Hab. ad ramos Rhois venenatæ New Jersy Amer. bor. (J. B. ELLIS).

16. **Scleroderris Eucalypti** (C. et H.) Sacc. *Dermatea Eucalypti* 2471

Cooke et Hark. Grevillea pag. 130. — Parva, erumpens, carnea; cupulis subsessilibus, solitariis, extus obscurioribus $\frac{1}{3}$ mm.; ascis clavatis; sporidiis subfusoideis utrinque obtusis, 20-30 = 10-12, quadrinucleatis, dein pseudo-triseptatis, hyalinis; paraphysibus flavidis.

Hab. in cortice Eucalypti in California.

17. **Scleroderris vermicularis** (Ces.) Sacc. *Tympanis vermicularis* 2472

Ces. Myc. Born. p. 11. — Cæspitibus nigrescentibus, usque 6 mm. diam. (*Myriangium* sat bene revocantibus); cupulis subcarnosis stromate discoideo tenui applanato nunc circinnatim, nunc gregatim emergentibus, obverse piriformibus, nonnunquam compressi sculis 1,5 mm. alt. e purpurascenti brunneis, sub lente minute granulosis vel pulverulentis, ostiolo punctiformi impresso; nucleo pallidiori; ascis longe pedicellatis, e cylindraceo clavatis, 64-70 μ . longis, tetrasporis; sporidiis vermicularibus, saepius uno alteroque fine reduncis, pluriseptatis, hyalinis, 24 μ . long.

Hab. ad lignum decorticatum Sarawak Bornei (BECCARI).

18. **Scleroderris fumosa** (C. et Ph.) Sacc. *Dermatea fumosa* C. et 2473

Ph. Grev. VIII, pag. 64. — Sparsa, sessilis; cupulis demum expansis 2-4 mm., extus pallidis, granulosis, margine elevato, flexuoso; hymenio murinaceo; ascis clavatis, longe stipitatis; sporidiis lanceolatis, rectis vel curvulis, endochromate quinque-divisis, 25-35 = 4-6.

Hab. ad ligna putrescentia, Dunedin Novæ Zelandiæ.

19. **Scleroderris aterrima** (Hazsl.) Sacc. *Encœlia aterrima* Hazsl. 2474

Einig. Disc. p. 10, Magyar szab. Disc. t. VIII, f. 17. — Sessilibus, opace nigra, margine nitidulo, subcupulata, 1 cm. lat.; ascis clavatis, stipitatis, parte superiore octosporis; sporidiis distichis, fusoideis, hyalinis, triseptatis.

Hab. in radice putri Pinuum prope Bade Czemeti Hungariæ.

20. **Scleroderris Syringæ** (Wint.) Sacc. *Cenangium Syringæ* Wint. 2475

Hedw. 1874, p. 57 c. ic. nec Rehm. — Ascomatibus ut in *Tympanide Syringæ* (quam cfr.); ascis octosporis; sporidiis fusoideis utrinque rotundatis, $31 \approx 5$, 8-10-guttulato-septatis, chlorino-hyalinis.

Hab. in ramis Syringæ vulgaris in Germania. — An forma octospora *Tymp. Syringæ?* Teste Minks in hac specie adsunt sporidia bilocularia. An species vere diversæ licet habitu simillimo?

II. *Phæoderris* Sacc. Sporidia colorata.

21. **Scleroderris betulina** (Peck) Sacc. *Cenangium betulinum* Peck 2476

35 Rep. p. 143. — Ascomatibus cæspitosis vel subseriatis, rima brevi corticali erumpentibus, primo sphæriæformibus, dein apertis atris, disco plano, tenuiter marginato sæpe mutua pressione irregulari, $170-350 \mu$. diam.; ascis oblongis vel subclavatis, $40 \approx 10$; sporidiis stipatis, oblongis, obtusis, pallide coloratis, triseptatis, $10-12 \approx 4$.

Hab. in cortice emortuo Betulæ populifoliæ, Center Americanæ borealis.

22. **Scleroderris pulveracea** (Hazsl.) Sacc. *Pezicula? pulveracea* 2477

Hazsl. Einig. Disc. p. 11, Magyar. szab. Disc. t. II, f. 19. — Gregaria, stromati nigricanti inserta, obconica, atra, intus albida, minuta, disco circulari, subconvexo; ascis clavatis, 4-8-sporis; sporidiis oblongo-cylindraceis, $14 \approx 2,5$, triseptatis, episporio crassuloso brunneo; paraphysibus filiformibus.

Hab. in cortice Betulæ in Hungaria.

23. **Scleroderris acerina** (Peck) Sacc. *Tymanis* Peck 31 Rep. Stat. 2478

Mus. p. 48. — Cupulis subcæspitosis, obconicis, erumpentibus, nigris, margine distincto sæpe flexuoso, disco concavo; ascis oblongo-cylindraceis; sporidiis oblongis, $13-20 \mu$. long. intus granulosis denique 4-nucleatis vel 3-septatis coloratis.

Hab. in cortice acerino, Adirondack et Catskill Mount. Amer.